Social Networks & Recommendation Systems

II. Historical overview of the complex network science. Examples of the real-life networks.

Grzegorz Siudem

Warsaw University of Technology



Warsaw University of Technology



MSc program in Data Science has been developed as a part of task 10 of the project "NERW PW. Science - Education - Development - Cooperation" co-funded by European Union from European Social Fund.

Before classes


Remember: graph representation methods

Adjacency matrix

$$A = \left[\begin{array}{rrr} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right].$$

Adjacency list

$$L = \{\{1, 2\}, \{2\}, \{1, 3\}\}.$$




Remember: vertex degree

Vertex degree

Number of incoming or outcoming edges.

$$k_{out} = \{4, 2, 3, 2, 4\},$$


$$k_{in} = \{3, 3, 3, 3, 3\}.$$



Remember: discrete probability distributions

Discrete probability distributions – examples

- binomial distribution,
- · Poissona distribution,
- · Zipf distribution.



$$P(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$P(k) = \frac{\lambda^k e^{-\lambda}}{k!}$$

$$P(k) = \frac{1/k^{s}}{H_{N,s}}$$

Reminder

Exercise to think about - continuous distributions


In complex networks, you often replace discrete distributions with continuous ones (this is the way we'll think during class). Find continuous analogs of distributions from the previous slide.

Lecture


Why are networks/graphs so useful?

Graphs are an illustration of a relationship

What relations are illustrated in the following graphs?



What is the difference between graphs and networks?



Properties of the real networks

· distributions with fat tails.

Properties of the real networks

- · distributions with fat tails.
- · heterognecity.

Properties of the real networks


- · distributions with fat tails.
- heterognecity.
- · small-world phenomenon (six steps?).

Properties of the real networks

- · distributions with fat tails.
- heterognecity.
- · small-world phenomenon (six steps?).
- · vertices are strongly correlated (see clustering coefficient).

Properties of the real networks

- · distributions with fat tails.
- · heterognecity.
- · small-world phenomenon (six steps?).
- · vertices are strongly correlated (see clustering coefficient).



Which of these graphs represents the real network?

Network Science is a composition of

Network Science is a composition of

· mathematics (Graph Theory, Probability Theory, Statistcs),

Network Science is a composition of

- · mathematics (Graph Theory, Probability Theory, Statistcs),
- physics (statistical physics methods, phase transitions),

Network Science is a composition of

- · mathematics (Graph Theory, Probability Theory, Statistcs),
- physics (statistical physics methods, phase transitions),
- computer science (graph visualization, storage and processing of large networks),

Network Science is a composition of

- · mathematics (Graph Theory, Probability Theory, Statistcs),
- physics (statistical physics methods, phase transitions),
- computer science (graph visualization, storage and processing of large networks),

· data science,

Network Science is a composition of

- · mathematics (Graph Theory, Probability Theory, Statistcs),
- physics (statistical physics methods, phase transitions),
- computer science (graph visualization, storage and processing of large networks),
- · data science,


with expert knowledge from

- sociology,
- · economics,
- · biology,
- · medicine,
- · engineering,
- · and many other...


The story of network science – Seven Bridges of Königsberg (1736)



The story of network science – Seven Bridges of Königsberg



The story of network science – Seven Bridges of Königsberg



SNARS

Source: 12

Great interest of sociologists in researching social networks

The most cited work from this period:

M. S. Granovetter. The strength of weak ties. American Journal of Sociology, 78: 1360, 1973.

Great interest of sociologists in researching social networks

The most cited work from this period:

M. S. Granovetter. The strength of weak ties. American Journal of Sociology, 78: 1360, 1973.

· mainly qualitative research,

Great interest of sociologists in researching social networks

The most cited work from this period:

M. S. Granovetter. The strength of weak ties. American Journal of Sociology, 78: 1360, 1973.

- · mainly qualitative research,
- independent of the research of mathematicians in graph theory,

Great interest of sociologists in researching social networks

The most cited work from this period:

M. S. Granovetter. The strength of weak ties. American Journal of Sociology, 78: 1360, 1973.

- · mainly qualitative research,
- · independent of the research of mathematicians in graph theory,
- · we do not focus on this line of research (with one exception!).

The first quantitative model of random networks

P. Erdős, A. Rényi. On random graphs. Publicationes Mathematicae, 6: 290, 1959.

· simple model in which we connect vertices randomly.

The first quantitative model of random networks

P. Erdős, A. Rényi. On random graphs. Publicationes Mathematicae, 6: 290, 1959.

- · simple model in which we connect vertices randomly.
- there are two types:

The first quantitative model of random networks

P. Erdős, A. Rényi. On random graphs. Publicationes Mathematicae, 6: 290, 1959.

- · simple model in which we connect vertices randomly.
- · there are two types:
 - with E edges and N vertices attached randomly.

The first quantitative model of random networks

P. Erdős, A. Rényi. On random graphs. Publicationes Mathematicae, 6: 290, 1959.

- · simple model in which we connect vertices randomly.
- · there are two types:
 - · with E edges and N vertices attached randomly.
 - \cdot with N vertices and probability of connection per edge equal to p.

The first quantitative model of random networks


P. Erdős, A. Rényi. On random graphs. Publicationes Mathematicae, 6: 290, 1959.

- · simple model in which we connect vertices randomly.
- · there are two types:
 - with E edges and N vertices attached randomly.
 - \cdot with N vertices and probability of connection per edge equal to p.

More details on Lecture 5.

Degree distribution

P(k) = fraction of vertices with degree k



Is this a real network?

The story of network science – Milgram experiment (1967)

Attention!

Do not confuse the *correspondence* experiment with the famous experiment of obedience to authorities!

The story of network science – Milgram experiment (1967)

Attention!

Do not confuse the *correspondence* experiment with the famous experiment of obedience to authorities!

The famous six handshakes

Milgram, Stanley, The Small World Problem, Psychology Today, 1967.

Attention!

Do not confuse the *correspondence* experiment with the famous experiment of obedience to authorities!

The famous six handshakes

Milgram, Stanley, The Small World Problem, Psychology Today, 1967.

Experiment description

 Milgram mails to 160 of randomly selected people a letter describing the experiment.

Attention!

Do not confuse the *correspondence* experiment with the famous experiment of obedience to authorities!

The famous six handshakes

Milgram, Stanley, The Small World Problem, Psychology Today, 1967.

Experiment description

- Milgram mails to 160 of randomly selected people a letter describing the experiment.
- The aim of the experiment was to deliver the letter to a certain, random person, whose photo, address and personal data were included in the letter.

Attention!

Do not confuse the *correspondence* experiment with the famous experiment of obedience to authorities!

The famous six handshakes

Milgram, Stanley, The Small World Problem, Psychology Today, 1967.

Experiment description

- Milgram mails to 160 of randomly selected people a letter describing the experiment.
- The aim of the experiment was to deliver the letter to a certain, random person, whose photo, address and personal data were included in the letter.
- If the recipient of the letter knew the addressee personally, he/she should deliver it to him/her, otherwise he/she should send it to another person who she/he suspected might know the addressee.

Conclusions

· Most of the letters were lost.

Conclusions

- · Most of the letters were lost.
- Only 42 out of 160 sent reached their destination.

Conclusions

- · Most of the letters were lost.
- · Only 42 out of 160 sent reached their destination.
- Few of the needed only two intermediaries to get to their destination!

Conclusions

- · Most of the letters were lost.
- Only 42 out of 160 sent reached their destination.
- Few of the needed only two intermediaries to get to their destination!
- · On average, the received letters were only forwarded six times.

Fun facts:

 Erdös number of scientist is the number of works which separates her/him from Paul Erdös.

Conclusions

- · Most of the letters were lost.
- · Only 42 out of 160 sent reached their destination.
- Few of the needed only two intermediaries to get to their destination!
- · On average, the received letters were only forwarded six times.

Fun facts:

- Erdös number of scientist is the number of works which separates her/him from Paul Erdös.
- Bacon number of actor/actress is the number of movies which separates him/her from Kevin Bacon.

Conclusions

- · Most of the letters were lost.
- Only 42 out of 160 sent reached their destination.
- Few of the needed only two intermediaries to get to their destination!
- · On average, the received letters were only forwarded six times.

Fun facts:

- Erdös number of scientist is the number of works which separates her/him from Paul Erdös.
- Bacon number of actor/actress is the number of movies which separates him/her from Kevin Bacon.

More details during Lecture 4.

Problem:

Poissonian graphs do not have small world property.

WS Model

D.J. Watts, S.H. Strogatz, Collective dynamics of 'small-world' networks, Nature. 393 440–442 (1998).

Problem:

Poissonian graphs do not have small world property.

WS Model

D.J. Watts, S.H. Strogatz, Collective dynamics of 'small-world' networks, Nature. 393 440–442 (1998).

Description:

 We start with a decorated regular network with periodic boundary conditions.

Problem:

Poissonian graphs do not have small world property.

WS Model

D.J. Watts, S.H. Strogatz, Collective dynamics of 'small-world' networks, Nature. 393 440–442 (1998).

Description:

- We start with a decorated regular network with periodic boundary conditions.
- · With probability *p* we switch every of nodes.

Problem:

Poissonian graphs do not have small world property.


WS Model

D.J. Watts, S.H. Strogatz, Collective dynamics of 'small-world' networks, Nature. 393 440–442 (1998).

Description:


- We start with a decorated regular network with periodic boundary conditions.
- With probability *p* we switch every of nodes.


More details during Lecture 5.



Problem:

Real-world networks usually have power law like distributions.





What does it means?

- · No typical scale.
- · Fat tails.
- Fast spreading epidemics...

BA model

A.-L. Barabási, R. Albert, Emergence on scaling in random networks, Science, 286:509-512, 1999.

A.-L. Barabási, R. Albert, H. Jeong, Mean-field theory for scale-free random networks, Physica A, 272: 173-187, 1999.

BA model

A.-L. Barabási, R. Albert, Emergence on scaling in random networks, Science, 286:509-512, 1999.

A.-L. Barabási, R. Albert, H. Jeong, Mean-field theory for scale-free random networks, Physica A, 272: 173-187, 1999.

Main assumptions:

 The network evolves i.e. with every step we add new vertex and few edges.

BA model

A.-L. Barabási, R. Albert, Emergence on scaling in random networks, Science, 286:509-512, 1999.

A.-L. Barabási, R. Albert, H. Jeong, Mean-field theory for scale-free random networks, Physica A, 272: 173-187, 1999.

Main assumptions:

- The network evolves i.e. with every step we add new vertex and few edges.
- Edges are added due to the preferential attachement rule (compare *rich get richer rule*).

BA model

A.-L. Barabási, R. Albert, Emergence on scaling in random networks, Science, 286:509-512, 1999.

A.-L. Barabási, R. Albert, H. Jeong, Mean-field theory for scale-free random networks, Physica A, 272: 173-187, 1999.

Main assumptions:

- The network evolves i.e. with every step we add new vertex and few edges.
- Edges are added due to the preferential attachement rule (compare *rich get richer rule*).

Both assumptions are necessary!

Construction procedure:

• We start with the small complete graph.

Construction procedure:

- We start with the small complete graph.
- In every step we add new vertex.

Construction procedure:

- · We start with the small complete graph.
- · In every step we add new vertex.
- Edges are connected due to the preferential attachment rule.

Construction procedure:

- · We start with the small complete graph.
- · In every step we add new vertex.
- Edges are connected due to the preferential attachment rule.
- We repeat the two points above.

Construction procedure:

- · We start with the small complete graph.
- · In every step we add new vertex.
- Edges are connected due to the preferential attachment rule.
- · We repeat the two points above.



Construction procedure:

- · We start with the small complete graph.
- · In every step we add new vertex.
- Edges are connected due to the preferential attachment rule.
- We repeat the two points above.



Summary:

• Evolving network wit power law degree distribution.

Summary:

- Evolving network wit power law degree distribution.
- Starting point of the modern complex network science.


Summary:

- Evolving network wit power law degree distribution.
- · Starting point of the modern complex network science.

Importance of so-called hubs.

Summary:


- Evolving network wit power law degree distribution.
- · Starting point of the modern complex network science.
- · Importance of so-called hubs.



23

If you are interested in the history of complex networks read





The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).
 - measuring with them real networks (Lectures 4,8, 13-15).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).
 - measuring with them real networks (Lectures 4,8, 13-15).
 - Description of the dynamical processes within real networks (e.g. epidemics, news spreading, etc.) (Lecture 8,11-15).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).
 - measuring with them real networks (Lectures 4,8, 13-15).
 - Description of the dynamical processes within real networks (e.g. epidemics, news spreading, etc.) (Lecture 8,11-15).

• Theoretical complex networks models (Lectures 5,6,9,10).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).
 - measuring with them real networks (Lectures 4,8, 13-15).
 - Description of the dynamical processes within real networks (e.g. epidemics, news spreading, etc.) (Lecture 8,11-15).
- Theoretical complex networks models (Lectures 5,6,9,10).
- Theoretical models of the dynamics in networks (Lectures 11,12).

The main task of network science is to model real phenomena and processes using complex networks, which are appropriate graphs.

Typical tools:

- · Quantitative description of the network, including
 - proposing new network metrics (Lecture 4).
 - · discussion of their properties (Lectures 4,8).
 - measuring with them real networks (Lectures 4,8, 13-15).
 - Description of the dynamical processes within real networks (e.g. epidemics, news spreading, etc.) (Lecture 8,11-15).
- Theoretical complex networks models (Lectures 5,6,9,10).
- Theoretical models of the dynamics in networks (Lectures 11,12).

· Visualization of networks (both theoretical and real) (Lecture 3).

Thank you for your attention!

Warsaw University of Technology



MSc program in Data Science has been developed as a part of task 10 of the project "NERW PW. Science - Education - Development - Cooperation" co-funded by European Union from European Social Fund.