

Social Networks & Recommendation Systems

V. Static random graphs.

Grzegorz Siudem

Warsaw University of Technology



**European
Funds**
Knowledge Education Development

**Warsaw University
of Technology**

European Union
European Social Fund



MSc program in Data Science has been developed
as a part of task 10 of the project
„NERW PW. Science - Education - Development - Cooperation”
co-funded by European Union from European Social Fund.

Before classes

Reminder

- Bernoulli distribution $\mathbb{P}(X = k) = \binom{N}{k} p^k (1 - p)^{N-k}$,
- Poisson distribution $\mathbb{P}(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}$,
- convergence of one to another (in what sense?)
- handshake lemma $\sum_i k_i = 2E$ ($\langle k \rangle = 2 \frac{E}{N}$).

Question:

Why does the 2 factor appear in the formula?

Lecture

A reminder of what complex network science is

complex networks = data + metrics + models + ...

complex networks = data + metrics + models + ...

Why do we need models?

- for generating networks under controlled conditions,

complex networks = data + metrics + models + ...

Why do we need models?

- for generating networks under controlled conditions,
- allow to extract network features important from a given point of view,

complex networks = data + metrics + models + ...

Why do we need models?

- for generating networks under controlled conditions,
- allow to extract network features important from a given point of view,
- give a chance to build good statistics while analyzing the dynamics on networks (classes 11 and 12),

complex networks = data + metrics + models + ...

Why do we need models?





- for generating networks under controlled conditions,
- allow to extract network features important from a given point of view,
- give a chance to build good statistics while analyzing the dynamics on networks (classes 11 and 12),
- give us an insight into the mechanisms behind real processes (see next classes and BA networks),

complex networks = data + metrics + models + ...

Why do we need models?

- for generating networks under controlled conditions,
- allow to extract network features important from a given point of view,
- give a chance to build good statistics while analyzing the dynamics on networks (classes 11 and 12),
- give us an insight into the mechanisms behind real processes (see next classes and BA networks),
- it is a very beautiful field of applied mathematics.

Attention! ER graphs are *normal*



SNARS And really poissonian, as we will see in a moment...

Erdős-Rényi model

On random graphs I.

Dedicated to O. Varga, at the occasion of his 50th birthday.

By P. ERDŐS and A. RÉNYI (Budapest).

Let us consider a "random graph" $\Gamma_{n,N}$ having n possible (labelled) vertices and N edges; in other words, let us choose at random (with equal probabilities) one of the $\binom{\binom{n}{2}}{N}$ possible graphs which can be formed from the n (labelled) vertices P_1, P_2, \dots, P_n by selecting N edges from the $\binom{n}{2}$ possible edges $\overline{P_i P_j}$ ($1 \leq i < j \leq n$). Thus the effective number of vertices of $\Gamma_{n,N}$ may be less than n , as some points P_i may be not connected in $\Gamma_{n,N}$ with any other point P_j ; we shall call such points P_i *isolated points*. We consider the isolated points also as belonging to $\Gamma_{n,N}$. $\Gamma_{n,N}$ is called completely connected if it effectively contains all points P_1, P_2, \dots, P_n (i. e. if it has no isolated points) and is connected in the ordinary sense. In the present paper we consider asymptotic statistical properties of random graphs for $n \rightarrow +\infty$. We shall deal with the following questions:

1. What is the probability of $\Gamma_{n,N}$ being completely connected?
2. What is the probability that the greatest connected component (sub-graph) of $\Gamma_{n,N}$ should have effectively $n-k$ points? ($k=0, 1, \dots$).
3. What is the probability that $\Gamma_{n,N}$ should consist of exactly $k+1$ connected components? ($k=0, 1, \dots$).
4. If the edges of a graph with n vertices are chosen successively so that after each step every edge which has not yet been chosen has the same probability to be chosen as the next, and if we continue this process until the graph becomes completely connected, what is the probability that the number of necessary steps r will be equal to a given number l ?


As (partial) answers to the above questions we prove the following four theorems. In Theorems 1, 2, and 3 we use the notation

$$(1) \quad N_c = \left\lfloor \frac{1}{2} n \log n + \epsilon n \right\rfloor$$


Two variants of the model:

- $G_{N,E}$ (Erdős-Rényi),
- $G_{N,p}$ (E. Gilbert)


Erdős-Rényi model in the $G_{N,E}$ form




Erdős-Rényi model in the $G_{N,E}$ form




Erdős-Rényi model in the $G_{N,E}$ form




Erdős-Rényi model in the $G_{N,p}$ form



Erdős-Rényi model in the $G_{N,p}$ form



Erdős-Rényi model in the $G_{N,p}$ form




Let us focus on the $G_{N,p}$ model


Assumptions

We analyze the ensemble described by two parameters:

- N number of vertices,
- p probability of two vertices connection.




Limiting case: $p = 0$



Conclusion:

$p = 0$ is a trivial case.


Limiting case: $p = 1$



Conclusion 1:

This is a very unsuccessful visualization...

Limiting case: $p = 1$




Conclusion 1:

~~This is a very unsuccessful visualization...~~

Conclusion 2:

SNARS $p = 1$ is a complete graphs.

ER graphs – dependence on the parameter p



Solution of the $G_{N,p}$ model

Question:

How many edges *on average* will appear?

Solution of the $G_{N,p}$ model

Question:

How many edges *on average* will appear?

Let's count

$$\langle E \rangle = p \times \frac{N(N-1)}{2},$$

which with the handshake lemma leads to

$$\langle k \rangle = p \frac{N(N-1)}{2} \frac{2}{N} = p(N-1) \approx pN.$$

Solution of the $G_{N,p}$ model

Question:

How many edges *on average* will appear?

Let's count

$$\langle E \rangle = p \times \frac{N(N-1)}{2},$$

which with the handshake lemma leads to

$$\langle k \rangle = p \frac{N(N-1)}{2} \frac{2}{N} = p(N-1) \approx pN.$$

So we know the average degree

What about the degree distribution of the vertices?

Solution of the $G_{N,p}$ model

Random variable describing vertex degree K

$$K = \sum_{i=1}^{N-1} X_i,$$

where X_i are iid variables with $\mathbb{P}(X_i = 1) = p$ $\mathbb{P}(X_i = 0) = 1 - p$.

Solution of the $G_{N,p}$ model

Random variable describing vertex degree K

$$K = \sum_{i=1}^{N-1} X_i,$$

where X_i are iid variables with $\mathbb{P}(X_i = 1) = p$ $\mathbb{P}(X_i = 0) = 1 - p$.

K has binomial distribution

$$\mathcal{P}(k) = \mathbb{P}(K = k) = \binom{N-1}{k} p^k (1-p)^{N-1-k}$$

Solution of the $G_{N,p}$ model

Random variable describing vertex degree K

$$K = \sum_{i=1}^{N-1} X_i,$$

where X_i are iid variables with $\mathbb{P}(X_i = 1) = p$ $\mathbb{P}(X_i = 0) = 1 - p$.

K has binomial distribution

$$\mathcal{P}(k) = \mathbb{P}(K = k) = \binom{N-1}{k} p^k (1-p)^{N-1-k}$$

Poissonian approximation

$$\mathcal{P}(k) \approx \frac{e^{-\langle k \rangle} \langle k \rangle^k}{k!},$$

because $\langle k \rangle \approx Np$.

Let's calculate variance for the poissonian case

$$\mathbb{E}(K) = \sum_{k=0}^{\infty} \frac{k e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle,$$

$$\mathbb{E}(K^2) = \sum_{k=0}^{\infty} \frac{k^2 e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle + \langle k \rangle^2.$$

$$\text{Var}(K) = \mathbb{E}(K^2) - [\mathbb{E}(K)]^2 = \langle k \rangle$$

Let's calculate variance for the poissonian case

$$\mathbb{E}(K) = \sum_{k=0}^{\infty} \frac{k e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle,$$

$$\mathbb{E}(K^2) = \sum_{k=0}^{\infty} \frac{k^2 e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle + \langle k \rangle^2.$$

$$\text{Var}(K) = \mathbb{E}(K^2) - [\mathbb{E}(K)]^2 = \langle k \rangle$$

Details?

During the project.

Let's calculate variance for the poissonian case

$$\mathbb{E}(K) = \sum_{k=0}^{\infty} \frac{k e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle,$$

$$\mathbb{E}(K^2) = \sum_{k=0}^{\infty} \frac{k^2 e^{-\langle k \rangle} \langle k \rangle^k}{k!} = \dots = \langle k \rangle + \langle k \rangle^2.$$

$$\text{Var}(K) = \mathbb{E}(K^2) - [\mathbb{E}(K)]^2 = \langle k \rangle$$

Details?

During the project.

Such small variance is not observed in the real life networks...

Clustering coefficient: reminder

We are looking for the value of the clustering coefficient for ER graphs

$$C_i = \frac{2E_i}{k_i(k_i - 1)}.$$

Non-physicality of ER graphs

Clustering coefficient: reminder

We are looking for the value of the clustering coefficient for ER graphs



$$C_i = \frac{2E_i}{k_i(k_i - 1)}.$$

Answer:

$$\langle C \rangle = \frac{p \langle k \rangle (\langle k \rangle - 1)}{\langle k \rangle (\langle k \rangle - 1)} = p.$$

Such small clustering coefficient is not observed in the real life networks...

Non-physicality of ER graphs




SNARS In the following classes, a more realistic BA model. ↗

Configuration model

Idea:


Let's build a graph from the bricks we have.



Configuration model

Idea:


Let's build a graph from the bricks we have.



Configuration model

Idea:


Let's build a graph from the bricks we have.



Configuration model

Idea:

Let's build a graph from the bricks we have.



What if you replace the actual bricks with selected *a priori*?

Configuration model

Let's choose the degree configuration according to the expected distribution:

An example:


$$\{1, 1, 1, 2, 2, 3\}$$

Configuration model

Let's choose the degree configuration according to the expected distribution:

An example:

$$\{1, 1, 1, 2, 2, 3\}$$



The only thing left to do is put these bricks together ...

Is it always possible?

Configuration model

Let's choose the degree configuration according to the expected distribution:

An example:

$$\{1, 1, 1, 2, 2, 3\}$$

The only thing left to do is put these bricks together ...

Is it always possible?

Question:

Find a (small) counterexample that the proposed procedure may not always be completed.


Generalization of the ER graphs

$$\begin{bmatrix} [p_{11}] & [p_{12}] & \dots & [p_{1N}] \\ [p_{21}] & [p_{22}] & \dots & [p_{2N}] \\ \dots & \dots & \ddots & \dots \\ [p_{N1}] & [p_{N2}] & \dots & [p_{NN}] \end{bmatrix}$$


Stochastic block model

Generalization of the ER graphs

$$\begin{bmatrix} [p_{11}] & [p_{12}] & \dots & [p_{1N}] \\ [p_{21}] & [p_{22}] & \dots & [p_{2N}] \\ \dots & \dots & \ddots & \dots \\ [p_{N1}] & [p_{N2}] & \dots & [p_{NN}] \end{bmatrix}$$



Watts-Strogatz model



Networks with given hamiltonian

Let us consider the space of every possible graph with N vertices
i.e. set $M_N = \mathbb{M}^{N \times N}(\{0, 1\})$. We want to define a probability
distribution on it.

Networks with given hamiltonian

Let us consider the space of every possible graph with N vertices
i.e. set $M_N = \mathbb{M}^{N \times N}(\{0, 1\})$. We want to define a probability
distribution on it.

We maximize entropy:

$$- \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G),$$

Networks with given hamiltonian

Let us consider the space of every possible graph with N vertices
i.e. set $M_N = \mathbb{M}^{N \times N}(\{0, 1\})$. We want to define a probability
distribution on it.

We maximize entropy:

$$- \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G),$$

Under certain condition $f(\mathcal{P}(G)) = 0$,

Networks with given hamiltonian

Let us consider the space of every possible graph with N vertices
i.e. set $M_N = \mathbb{M}^{N \times N}(\{0, 1\})$. We want to define a probability
distribution on it.

We maximize entropy:

$$- \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G),$$

Under certain condition $f(\mathcal{P}(G)) = 0$,

Which leads us to Lagrange multipliers

$$\mathcal{L}[\mathcal{P}(G)] = - \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G) + \lambda f(\mathcal{P}(G))$$

Networks with given hamiltonian

Let us consider the space of every possible graph with N vertices
i.e. set $M_N = \mathbb{M}^{N \times N}(\{0, 1\})$. We want to define a probability
distribution on it.

We maximize entropy:

$$- \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G),$$

Under certain condition $f(\mathcal{P}(G)) = 0$,

Which leads us to Lagrange multipliers

$$\mathcal{L}[\mathcal{P}(G)] = - \sum_{G \in M_N} \mathcal{P}(G) \ln \mathcal{P}(G) + \lambda f(\mathcal{P}(G))$$

It *only* remains to solve this equation

$$\frac{\partial \mathcal{L}}{\partial \mathcal{P}(G)} = 0.$$

Summary

Homework:

Read about:

- Dulbecco rule,
- Matthew effect,
- rich get richer rule,
- Yule processes.

Suggested source:

- M. Perc, *Journal of The Royal Society Interface* **11** (2014)



**European
Funds**
Knowledge Education Development

**Warsaw University
of Technology**

European Union
European Social Fund



MSc program in Data Science has been developed
as a part of task 10 of the project
„NERW PW. Science - Education - Development - Cooperation”
co-funded by European Union from European Social Fund.

Thank you for your attention!