

Missing Numbers



Numeros the Artist had two lists that were permutations of one another. He was very proud. Unfortunately, while transporting them from one exhibition to another, some numbers were lost out of the first list. Can you find the missing numbers?

Notes

- If a number occurs multiple times in the lists, you must ensure that the frequency of that number in both lists is the same. If that is not the case, then it is also a missing number.
- You have to print all the missing numbers in ascending order.
- Print each missing number once, even if it is missing multiple times.
- The difference between maximum and minimum number in the second list is less than or equal to 100.

Function Description

Complete the *missingNumbers* function in the editor below. It should return a sorted array of missing numbers.

missingNumbers has the following parameter(s):

- *arr*: the array with missing numbers
- *brr*: the original array of numbers

Input Format

There will be four lines of input:

n - the size of the first list, *arr*

The next line contains *n* space-separated integers *arr*[*i*]

m - the size of the second list, *brr*

The next line contains *m* space-separated integers *brr*[*i*]

Constraints

- $1 \leq n, m \leq 2 \times 10^5$
- $n \leq m$
- $1 \leq x \leq 10^4, x \in B$
- $X_{max} - X_{min} < 101$

Output Format

Output the missing numbers in ascending order.

Sample Input

```
10
203 204 205 206 207 208 203 204 205 206
13
203 204 204 205 206 207 205 208 203 206 205 206 204
```

Sample Output

```
204 205 206
```

Explanation

204 is present in both arrays. Its frequency in *arr* is **2**, while its frequency in *brr* is **3**. Similarly, **205** and **206** occur twice in *arr*, but three times in *brr*. The rest of the numbers have the same frequencies in both lists.