

An **XOR** operation on a list is defined here as the  $xor (\oplus)$  of all its elements (e.g.:  $XOR(\{A, B, C\}) = A \oplus B \oplus C$ ).

The **XorSum** of set **arr** is defined here as the sum of the **XORs** of all non-empty subsets of **arr** known as **arr'**. The set **arr'** can be expressed as:

$$XorSum(arr) = \sum_{i=1}^{2^n-1} XOR(arr'_i) = XOR(arr'_1) + XOR(arr'_2) + \dots + XOR(arr'_{2^n-2}) + XOR(arr'_{2^n-1})$$

**For example:** Given set  $arr = \{n_1, n_2, n_3\}$

- The set of possible non-empty subsets is:  
 $arr' = \{\{n_1\}, \{n_2\}, \{n_3\}, \{n_1, n_2\}, \{n_1, n_3\}, \{n_2, n_3\}, \{n_1, n_2, n_3\}\}$
- The **XorSum** of these non-empty subsets is then calculated as follows:  
 $XorSum(arr) = n_1 + n_2 + n_3 + (n_1 \oplus n_2) + (n_1 \oplus n_3) + (n_2 \oplus n_3) + (n_1 \oplus n_2 \oplus n_3)$

Given a list of **n** space-separated integers, determine and print  $XorSum \% (10^9 + 7)$ .

For example,  $arr = \{3, 4\}$ . There are three possible subsets,  $arr' = \{\{3\}, \{4\}, \{3, 4\}\}$ . The XOR of  $arr'[1] = 3$ , of  $arr'[2] = 4$  and of  $arr[3] = 3 \oplus 4 = 7$ . The XorSum is the sum of these:  $3 + 4 + 7 = 14$  and  $14 \% (10^9 + 7) = 14$ .

**Note:** The cardinality of **powerset**(**n**) is  $2^n$ , so the set of non-empty subsets of set **arr** of size **n** contains  $2^n - 1$  subsets.

## Function Description

Complete the **xoringNinja** function in the editor below. It should return an integer that represents the XorSum of the input array, modulo  $(10^9 + 7)$ .

xoringNinja has the following parameter(s):

- arr**: an integer array

## Input Format

The first line contains an integer **T**, the number of test cases.

Each test case consists of two lines:

- The first line contains an integer **n**, the size of the set **arr**.
- The second line contains **n** space-separated integers  $arr[i]$ .

## Constraints

$$1 \leq T \leq 5$$

$$1 \leq n \leq 10^5$$

$$0 \leq arr[i] \leq 10^9, 1 \leq i \leq n$$

## Output Format

For each test case, print its  $XorSum \% (10^9 + 7)$  on a new line. The  $i^{th}$  line should contain the output for the  $i^{th}$  test case.

Sample Input 0

```
1
3
1 2 3
```

Sample Output 0

```
12
```

Explanation 0

The input set,  $S = \{1, 2, 3\}$ , has 7 possible non-empty subsets:  $S' = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 3\}, \{1, 2, 3\}\}$ .

We then determine the *XOR* of each subset in  $S'$ :

$XOR(\{1\}) = 1$   
 $XOR(\{2\}) = 2$   
 $XOR(\{3\}) = 3$   
 $XOR(\{1, 2\}) = 1 \oplus 2 = 3$   
 $XOR(\{2, 3\}) = 2 \oplus 3 = 1$   
 $XOR(\{1, 3\}) = 1 \oplus 3 = 2$   
 $XOR(\{1, 2, 3\}) = 1 \oplus 2 \oplus 3 = 0$

Then sum the results of the *XOR* of each individual subset in  $S'$ , resulting in  $XorSum = 12$  and  $12 \% (10^9 + 7) = 12$ .

Sample Input 1

```
2
4
1 2 4 8
5
1 2 3 5 100
```

Sample Output 1

```
120
1648
```