

Hotelling T squared: OLS versus CV (quadratic spline degree 2; knots 0; l_21_mid vs t_vis_2_strong, gamma_CV = 0.4)

$T^2 = \beta[J]' V[J,J]^{-1} \beta[J]$; F approximation = $T^2 / \text{rank}(V[J,J])$; df2 = 1949; decisions use alpha = 0.05; direct causes in red

Covariate	Terms	OLS T2	OLS p	OLS H0	CV T2	CV p	CV H0
red	3	820.308	<1e-04	reject	418.062	<1e-04	reject
green	3	678.311	<1e-04	reject	384.308	<1e-04	reject
l_21	3	228.939	<1e-04	reject	167.706	<1e-04	reject
blue	3	89.005	<1e-04	reject	55.305	<1e-04	reject
vis_2	3	168.436	<1e-04	reject	36.228	<1e-04	reject
l_22	3	12.178	0.00691	reject	9.137	0.0278	reject
l_12	3	5.658	0.13	do not reject	6.281	0.0991	do not reject
current	3	10.805	0.013	reject	6.054	0.109	do not reject
vis_1	3	10.010	0.0187	reject	5.336	0.149	do not reject
ir_3	3	2.649	0.449	do not reject	3.938	0.269	do not reject
ir_1	3	3.765	0.288	do not reject	3.657	0.301	do not reject
vis_3	3	0.577	0.902	do not reject	3.332	0.344	do not reject
pol_2	3	4.792	0.188	do not reject	2.720	0.437	do not reject
l_11	3	3.423	0.331	do not reject	1.218	0.749	do not reject
pol_1	3	0.140	0.987	do not reject	0.926	0.819	do not reject
l_31	3	0.697	0.874	do not reject	0.677	0.879	do not reject
l_32	3	4.737	0.192	do not reject	0.402	0.94	do not reject