

Validation of a Mathematical Model Describing the Dynamics of Chemotherapy for Chronic Lymphocytic Leukemia In Vivo

Alessia Guadagnin Pattaro¹, Giovanni Plazzotta¹, Annarita Zanon¹

Abstract

To be added

¹ Master's degree in Quantitative and Computational Biology, University of Trento

1. Introduction

1.1 Chronic Lymphocytic Leukemia

Chronic lymphocytic leukemia (CLL) is the most common type of blood cancer in the Western world. It involves an accumulation of lymphocytes B in secondary lymphoid organs, spleen, peripheral blood, and bone marrow.[2, 3] There is no known cause for this disease even if it is suspected to have a genetic basis. Mutations in *IGHV* (immunoglobulin heavy variable) genes are thought to help distinguishing different types of clinical behaviours of CLL [1].

2. Methods

3. Results and Discussion

4. Conclusions

References

1. A., X., A., A. & I., K. Immunoglobulin heavy variable (*IGHV*) genes and alleles: new entities, new names and implications for research and prognostication in chronic lymphocytic leukaemia. *Immunogenetics* **67**, 61–66 (2015).
2. A., B. J. Treatment of Chronic Lymphocytic Leukemia. *The New England Journal of Medicine* **383**, 460–473 (2020).
3. Ciril, R. & Emilio, M. Chronic Lymphocytic Leukemia. *The New England Journal of Medicine* **333**, 1052–1057 (1995).