

① convergenza della serie

$$\sum_{n=1}^{\infty} \frac{\sin(nx)}{n^2+x^2}$$

$$\lim_{n \rightarrow \infty} \frac{\sin(nx)}{n^2+x^2} = 0 \text{ può convergere}$$

così totale

$$\left| \frac{\sin(nx)}{n^2+x^2} \right| \leq \frac{1}{n^2+x^2} \leq \left( \frac{1}{n^2} \right)$$

teorema generale serie armonica generalizzata  
convergente

$$\textcircled{2} \quad f(x, y) = \frac{(x+y)^4}{4} + \frac{(x-y)^4}{4} - x^2$$

$$f_x = (x+y)^3 + (x-y)^3 - 2x$$

$$f_y = (x+y)^3 - (x-y)^3$$

$$\begin{cases} 2x^3 - 2x = 0 \Rightarrow 2x(x^2 - 1) = 0 \\ (x+y)^3 = (x-y)^3 \Rightarrow x+y = x-y \Rightarrow y=0 \end{cases}$$

$$(0, 0) \quad (1, 0) \quad (-1, 0)$$

$$f_{xx} = 3(x+y)^2 + 3(x-y)^2 - 2$$

$$f_{yy} = 3(x+y)^2 + 3(x-y)^2$$

$$f_{xy} = f_{yx} = 3(x+y)^2 - 3(x-y)^2$$

$$Hf(0, 0) = \begin{vmatrix} -2 & 0 \\ 0 & 0 \end{vmatrix} = 0$$

$$Hf(\pm 1, 0) = \begin{vmatrix} 4 & 0 \\ 0 & 6 \end{vmatrix} = 24 \quad \begin{matrix} \text{Semi minimum} \\ f_{xx} > 0 \end{matrix}$$


$$f(x, y) - f(0, 0) = \frac{(x+y)^4}{4} + \frac{(x-y)^4}{4} - x^2 = f(x, y)$$

$$f(0, y) = \frac{y^4}{4} + \frac{y^4}{4} > 0 \quad \forall y$$

$$f(x, 0) = \frac{x^4}{2} - x^2 = x^2 \left( \frac{x^2}{2} - 1 \right) > 0$$

$$x < -\sqrt{2} \vee x > \sqrt{2}$$

no max no min in  $(0,0)$



③  $\omega(x,y) = \frac{2x-y}{x^2+y^2} dx + \frac{2y+x}{x^2+y^2} dy$   
 e' esoterico  $\int \omega$  da  $\gamma$  e  $\gamma: \{x^2+y^2=1\}$   
 $D = \mathbb{R}^2 \setminus \{(0,0)\}$

e' chiusa?

$$\frac{\partial a}{\partial y} = \frac{-x^2 - y^2 - 2y(2x-y)}{(x^2+y^2)^2} = \frac{-x^2 + y^2 - 4xy}{(x^2+y^2)^2}$$

$$\frac{\partial b}{\partial x} = \frac{x^2 + y^2 - 2x(2y+x)}{(x^2+y^2)^2} = \frac{-x^2 + y^2 - 4xy}{(x^2+y^2)^2}$$

e' chiusa

$$\begin{aligned} \int \frac{2x-y}{x^2+y^2} dx &= \int \frac{2x}{x^2+y^2} dx - \int \frac{y}{x^2+y^2} dx \\ &= \log(x^2+y^2) - \int \frac{y}{y^2 \left(1 + \left(\frac{x}{y}\right)^2\right)} dx \\ &= \log(x^2+y^2) - \arctan\left(\frac{x}{y}\right) + g(y) \end{aligned}$$

$$F_y = \frac{2y}{x^2+y^2} + \frac{x}{y^2 \left(1 + \frac{x^2}{y^2}\right)} + g'(y) = \frac{2y+x}{x^2+y^2}$$

$$\frac{2y}{x^2+y^2} + \frac{x}{x^2+y^2} \theta' = \frac{2y+x}{x^2+y^2} \Rightarrow \theta(y) = \cos t$$

$$F(x, y) = \log(x^2+y^2) - \arctan\left(\frac{x}{y}\right) + \cos t$$

$$\text{Dom } F = \{y \neq 0\} \neq \text{Dom } \omega = (\mathbb{R}^2 \setminus \{0,0\})$$

$$\Rightarrow \int_{C(0,1)} \omega = 2\pi \quad (\text{onde } \cos t \text{ é uma non sense}). \quad (*)$$

$$\gamma: \begin{cases} x = \cos \theta \\ y = \sin \theta \end{cases} \quad \theta \in [0, 2\pi]$$

$$\int_0^{2\pi} \frac{2\cos\theta - \sin\theta}{1} (-\sin\theta) + \frac{2\sin\theta + \cos\theta}{1} (\cos\theta)$$

$$\int_0^{2\pi} -2\sin\theta\cos\theta + \sin^2\theta + 2\sin\theta\cos\theta + \cos^2\theta$$

$$\int_0^{2\pi} 1 d\theta = 2\pi$$

$\Rightarrow$  o integral de uma curva chiusa  
 não é 0  $\Rightarrow$  não pode ser exata.

$$④ \quad y'' - 2y' + y = e^x \sin x$$

$$y'' - 2y' + y = 0$$

$$\lambda^2 - 2\lambda + 1 = 0 \Rightarrow (\lambda - 1)^2 = 0 \Rightarrow \lambda_{1,2} = 1 \text{ double}$$

$$y_1 = e^x \quad y_2 = x e^x$$

$$\lambda + i\mu = i \quad \text{non real}$$

$$y_p(x) = e^x (A \sin x + B \cos x)$$

$$\begin{aligned} y'_p &= e^x (A \sin x + B \cos x) + e^x (A \cos x - B \sin x) \\ &= e^x ((A - B) \sin x + (A + B) \cos x) \end{aligned}$$

$$\begin{aligned} y''_p &= e^x ((A - B) \sin x + (A + B) \cos x) + \\ &\quad e^x ((A - B) \cos x - (A + B) \sin x) \end{aligned}$$

$$\begin{aligned} &e^x [-2B \sin x + 2A \cos x] - 2e^x [(A - B) \sin x + \\ &+ (A + B) \cos x] + e^x [A \sin x + B \cos x] \\ &= e^x \sin x \end{aligned}$$

$$(-2B - 2A + 2B + A) \sin x + (2A - 2A - 2B + B) \cos x = \sin x$$

$$\cos x = \sin x$$

$$-A \sin x - B \cos x = \sin x \Rightarrow \begin{aligned} A &= -1 \\ B &= 0 \end{aligned}$$

$$y_p = -e^x \sin x$$

$$y(x) = C_1 e^x + C_2 x e^x - e^x \sin x$$

$$(5) \iiint_E f \, dx \, dy \, dz$$

$$f(x, y, z) = [4 - x^2 + 2x - 1 - y^2] z$$

$$E = \{ (x-1)^2 + y^2 \leq 4, y \geq 0, 0 \leq z \leq 2 \}$$

$$\iint_D \left[ \int_0^2 (4 - x^2 + 2x - 1 - y^2) z \, dz \right]$$

$$\iint_D 2(3 - x^2 + 2x - y^2) \, dx \, dy$$

$$\begin{cases} x = 1 + \rho \cos \theta \\ y = \rho \sin \theta \end{cases} \quad \begin{matrix} \rho \in [0, 2] \\ \theta \in [0, \pi] \end{matrix}$$

$$\int_0^2 \rho \, d\rho \int_0^\pi 2\rho [3 - 1 - \rho^2 \cos^2 \theta - \cancel{2\rho \cos \theta} + \cancel{2\rho \cos \theta} - \rho^2 \sin^2 \theta + z] \, d\theta$$

$$\begin{aligned} \int_0^2 (8\rho - 2\rho^3) \pi \, d\rho &= \pi \left( 4\rho^2 - \frac{\rho^4}{2} \right) \Big|_0^2 \\ &= \pi (16 - 8) = 8\pi \end{aligned}$$

metodo variazione delle costanti es. 5

$$y_p = C_1(x)e^x + C_2(x)xe^x$$

$$\begin{cases} C_1'(x)e^x + C_2'(x)xe^x = 0 \end{cases}$$

$$\begin{cases} C_1'(x)e^x + C_2'(x)(e^x + xe^x) = e^x \sin x \end{cases}$$

$$W = \begin{vmatrix} e^x & xe^x \\ e^x & e^x + xe^x \end{vmatrix} = e^{2x}(1+x) - xe^{2x} = e^{2x}$$

$$C_1' = \frac{\begin{vmatrix} 0 & xe^x \\ e^x \sin x & e^x(1+x) \end{vmatrix}}{e^{2x}} = \frac{-xe^{2x} \sin x}{e^{2x}}$$

$$C_2' = \frac{\begin{vmatrix} e^x & 0 \\ e^x & e^x \sin x \end{vmatrix}}{e^{2x}} = \frac{e^{2x} \sin x}{e^{2x}} = \sin x$$

$$C_1(x) = \int -x \sin x \, dx = x \cos x - \int \cos x \, dx = x \cos x - \sin x$$

$$C_2(x) = \int \sin x \, dx = -\cos x$$

$$y_p(x) = (x \cos x - \sin x)e^x - xe^x \cos x$$

$$g_p(x) = -e^x \sin x$$

$$\Rightarrow y(x) = c_1 e^x + c_2 x e^x - e^x \sin x,$$

(i due metodi sono equivalenti).