Fixed $m_G L$ campaign

Theory, methods and preliminary results

Current status and scope

The canonical analysis contains 49 physical ensembles and 200 independent chains: 1,904,000 measured sweeps, 1,796,340 after independent thermalization cuts, and 1,786,070 used in complete blocks. There are 49 numerical mass estimates and 10 usable preliminary coupling interpolations. All three targets, $c = 0.5, 1, 2$, have $N_t = 2, 3, 4$; $c = 1$ also has $N_t = 6$. Every new bootstrap operation uses exactly 500 replicas. These counts describe the analyzed register; new simulation batches are tracked separately below. Raw histories remain on qbig.

The register now also includes three $N_t = 1$ matched-coupling ensembles: twelve independent 8,000-sweep chains completed on lnode19 A100 GPUs on 14 October. Their couplings are the three current $N_t = 2$ tuning estimates. For these rows, the folder's c_{ref} denotes that source target; the actual $N_t = 1$ mass is measured without imposing $m_G N_t = c_{\mathrm{ref}}$. They are excluded from fixed-c interpolation and continuum fits.

Eight matched-y ensembles, $y = \beta/N_t$, add thirty-two 8,000-sweep chains. The four completed on 16 October extend $y = 1.5$ to $N_t = 3, 4, 6$ and check the transverse volume at $N_t = 4, \beta = 6.2$. Four $N_t = 1, N_s = 128$ ensembles completed on 20 October. These rows have `target_c=null`, with $c_{\mathrm{ref}} = 0.5$ identifying their source family. They enter correlator and scaling comparisons. Fixed-c tuning and its continuum inputs remain unchanged.

The four 80,000-sweep chains at $c = 0.5, N_t = 3, N_s = 80, \beta = 4.4$ have the explicit flag `length_outlier`. Their length differs from the 8,000-sweep standard; the flag does not establish a physics bias. They are pooled with four independent 8,000-sweep chains of identical parameters. Eight 10,000-sweep chains are explicitly labelled prior inputs from the fixed-β/N_t campaign, at $N_t = 2, \beta = 3.0$ and 2.1.

All defined estimates remain available for visual review. Flags identify window drift, incomplete error plateaus, limited block counts, chain differences, numerical covariance problems and interpolation sensitivity. They propagate to preliminary continuum results. The editable `analysis/mg_nt_fixed/review.csv` records manual window choices and inclusion decisions. No flag is silently converted into a rejection.

How to read the results

Each ensemble lives at `analysis/mg_nt_fixed/c_<c>/Nt_<Nt>/Ns_<Ns>/beta_<beta>/`. It contains correlators, all fit windows, chain statistics and compact bootstrap arrays. Its `plots/` directory shows correlator fits, residuals and the full mass-window scan; `plots/first_stage/` shows every chain's thermalization cut, bootstrap error versus block size, the selected block and block counts. Blocking examples are kept outside this PDF. Comparisons sit at the nearest common physics level. The CSV column guide is `docs/mg_nt_fixed/csv_columns.md`. Figures retain axis labels and legends, with no displayed titles or footer notes. Their paths, CSVs and this logbook supply the physics parameters and diagnostic context. The top-level correlator, ensemble, chain, autocorrelation and fit-window tables include every analyzed simulation, including both untuned diagnostic batches.

Theory: what is being held fixed?

The model is three-dimensional SU(2) Yang–Mills theory with Wilson action

$$S_W = -\frac{\beta}{2} \sum_p \mathrm{Re}\,\mathrm{Tr}\, U_p + \text{constant}.$$

Both transverse directions are periodic. Tangential links on two temporal Dirichlet surfaces are fixed to the identity [1]. The separation is $L = aN_t$; the transverse box size is $L_s = aN_s$. Current simulations store $N_t + 1$ spatial-link slices and N_t temporal-link layers, with no closing layer. Thus N_t counts links between the fixed surfaces. Historical wrapping-sector outputs retain their original executable provenance. With fixed surface links their action factorizes into the physical slab and an independent auxiliary sector, so removing that sector preserves physical slab expectation values.

For integer transverse separation r, define

$$G(\mathbf{x}) = \prod_{t=0}^{N_t-1} U_t(\mathbf{x}, t),$$
$$C_G(r) = \left\langle \tfrac{1}{2}\,\mathrm{Re}\,\mathrm{Tr}[G(\mathbf{x})G^\dagger(\mathbf{x} + r\hat{e})] \right\rangle.$$

Origins and both transverse axes are averaged. The normalization gives $C_G(0) = 1$. The residual boundary transformations act as $G \mapsto V_L G V_R^\dagger$; in $GG^\dagger$, the V_R factors cancel and the trace removes the remaining V_L conjugation. This is the $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ symmetry of the principal chiral model (PCM). At $N_t = 1$ the slab plaquettes reduce exactly to nearest-neighbor PCM interactions [1].

The asymptotic decay of C_G measures the lightest transverse screening scale with overlap in this holonomy channel, under single-state dominance. With $\widehat{m} = am_G$, the trajectory is

$$c = m_G L = \widehat{m} N_t \in \{0.5, 1, 2\}, \qquad \xi_{\mathrm{exp}}/a = 1/\widehat{m} = N_t/c, \qquad \xi_{\mathrm{exp}}/L = 1/c.$$

The correlation length is twice, equal to, or half the slab thickness. These are geometry-to-correlation-length ratios; identifying a phase or a limiting effective theory requires additional evidence. The smallest target has only a factor-two length separation. A controlled PCM long-distance interpretation would benefit from $m_G L \ll 1$ and agreement of further dimensionless observables. The present scope is the holonomy mass.

Increasing N_t at fixed c gives, step by step,

$$a/L = 1/N_t \to 0, \qquad \widehat{m} = c/N_t \to 0, \qquad \xi_{\mathrm{exp}}/a = N_t/c \to \infty.$$

Thus the same physical thickness is increasingly well resolved in lattice units, using the measured screening length as the ruler. The transverse volume is separately controlled through $m_G L_s = \widehat{m} N_s$.

From correlator to continuum mass

At each N_t, nearby couplings are simulated and their masses determine $\beta_\star$ satisfying $\widehat{m}(\beta_\star, N_t)N_t = c$. The tuning condition fixes $m_G L$ by construction. The nontrivial continuum information is the coupling required to satisfy it. In the isotropic Wilson convention $\beta = 4/(ag_{3,0}^2)$ [2],

$$g_{3,0}^2 L = (ag_{3,0}^2)N_t = 4N_t/\beta_\star,$$
$$Q(N_t) \equiv m_G/g_{3,0}^2 = \frac{am_G}{ag_{3,0}^2} = \frac{\widehat{m}\beta_\star}{4} = \frac{c\beta_\star}{4N_t}.$$

The plots fit $Q(N_t) = Q_0 + d/N_t$ only ($p = 1$). The equivalent $\beta_\star/N_t$ plots are in each target's comparison directory. At finite spacing these are bare-coupling coordinates; interpretation in continuum coupling units assumes the usual continuum matching. At fixed boundary prescription and large transverse volume, the result determines the local inverse relation between $m_G L$ and $g_3^2 L$.

The $p = 1$ form is a restricted cutoff ansatz. Fits using all resolutions and omitting the coarsest resolution show sensitivity to this choice. Two points determine a line with zero residual degrees of freedom and provide no residual test of the ansatz. Statistical bands are conditional on the chosen mass windows, interpolation windows and volumes. The CSV also reports a separate conservative linear-response bound on the continuum intercept from independently signed two-point interpolation shifts at each resolution. This bound is not added in quadrature to the statistical error.

Mass analysis and correlated uncertainty

Each chain is cut independently with MSER-5 and a 500-sweep minimum; the largest cut across monitored observables is used. Profiles at cadence five enter as equilibration monitors. Integrated autocorrelation times use the self-consistent Sokal window and are reported in sweeps [3]. One composite sweep is a heatbath update and five overrelaxation updates. Holonomy observables are measured every sweep. The second-moment length is monitored using ensemble structure factors,

$$\xi_{2\mathrm{nd}}/a = \frac{1}{2\sin(\pi/N_s)}\sqrt{\frac{\widetilde{C}_G(0)}{\widetilde{C}_G(p_{\min})} - 1}, \qquad ap_{\min} = 2\pi/N_s.$$

The same joint replicas propagate its error. Ratios are formed after ensemble averaging; unresolved replicas are recorded explicitly.

Complete nonoverlapping blocks are formed inside each chain. Candidate lengths scan the whole post-cut history, starting at five sweeps. The chosen length exceeds twice the largest measured autocorrelation time and uses the earliest common three-length error plateau when available. Each curve point is a separate 500-replica bootstrap SE. Plateau tolerances are 5% or two approximate SE uncertainties, whichever is larger; the latter use Welch–Satterthwaite degrees of freedom and the Monte Carlo SE of 500 draws. This is a diagnostic stability criterion. Missing plateaus are flagged. Fewer than 30 complete blocks is a warning; 50 is the target.

Pooling, fitting and window review

If chain k contributes n_k complete blocks of B_k measurements, $w_k = n_k B_k / \sum_j n_j B_j$. In each of exactly 500 replicas, draw n_k block vectors with replacement within chain k, then form $C^{(b)} = \sum_k w_k \overline{C}_k^{(b)}$ [4]. The draw acts on the entire separation vector. Chain identities and weights are fixed. The ensemble mean is $\sum_k w_k \overline{C}_k$. Independent chains retain their own cuts, autocorrelations and block boundaries. Split-history and between-chain discrepancies are reported separately; maxima over many correlated observables are review diagnostics.

The asymptotic two-image model is [1, Eq. (4.14)]

$$C_G(r) \simeq A\{K_0(\widehat{m}r) + K_0[\widehat{m}(N_s - r)]\}.$$

Its interpretation requires a stable long-distance window and finite-volume control. Additional images, excited contributions and lattice dispersion may matter outside that regime. Each central and bootstrap fit minimizes the diagonal-error objective. Parameter errors are the SD of the same fit over the 500 vector replicas. The complete covariance is used to report $\chi^2_{\text{full}} = r^T \,\text{Cov}^{-1}\, r$ after fitting. This statistic is descriptive: the fitted parameters do not minimize it, so a nominal analytic chi-square p-value is not appropriate. No p-value is computed and no chi-square gate is imposed. Singular covariance leaves this diagnostic undefined while retaining a defined diagonal fit.

For the fixed-c and matched-y ensembles, all contiguous windows with at least five points are fitted, with every positive $R_{\min}$ and $R_{\max} = N_s/2, N_s/2 - 2, \ldots \geq \lceil N_s/3 \rceil$. The recommendation is the earliest stable 3×3 patch in adjacent endpoints, preferring the largest $R_{\max}$. Pairwise masses must differ by at most two combined bootstrap SEs. Overlapping windows are correlated; this rule is a heuristic for visual review. If no such patch exists, the least-drifting numerical patch is flagged and retained. Failed refit slots remain NaN in the saved 500-slot arrays; success counts and failure reasons are in CSV. Relative error above 25%, low success fraction and a missing mass-error blocking plateau produce flags.

For the new $N_t = 1$ boxes with $N_s = 512, 2048$, the user approved a bounded endpoint scan: seven allowed upper endpoints and logarithmically spaced lower endpoints, followed by dense adjacent checks near three candidate plateaus. Every integer separation inside each window and all 500 refits are retained. The recommendation is the earliest stable patch among tested windows; it does not establish the exhaustive earliest patch. The $N_s = 128$ scan remains exhaustive. A covariance formed from 500 centered replicas has rank at most 499; windows with 500 or more separations therefore have undefined full-covariance χ^2. Fewer independent block vectors can reduce the rank further. No inverse regularization is used to manufacture a goodness-of-fit statistic.

The collector enforces the additional block-count bound $\text{rank(Cov)} \leq \min[499, \sum_k (n_k - 1)]$: within each chain, centered resampling multiplicities sum to zero, leaving at most $n_k - 1$ independent directions. Wider windows have undefined full χ^2 even if roundoff allows Cholesky factorization. Diagonal-fit masses and their 500 refits are unchanged by this diagnostic correction.

Coupling windows and volume choices

For each c, N_t, all contiguous three-coupling windows are examined using $\log(\widehat{m}N_t) = a_0 + b\beta$. Only centrally bracketed targets yield an interpolation; undefined slopes or unbracketed targets are recorded. Each aligned mass replica gives $\beta_\star^{(b)} = (\log c - a_0^{(b)})/b^{(b)}$. The median and 16–84% interval are quoted. The largest shift relative to a target-bracketing two-point interpolation is a separate systematic. The closest bracketing window is suggested; manual choices remain editable.

The larger $N_s = 96$ boxes are primary at $c = 1, N_t = 4, \beta = 4.95, 5.3$; the $N_s = 48$ versions remain volume diagnostics. At $c = 0.5, N_t = 4, \beta = 5.58394149$, $N_s = 96$ is primary and 128 is a matched-volume check. The $N_t = 4, c = 1$ confirmation ensemble now contributes to tuning. Its earlier independence applies to the historical interpolation only.

Coupling and continuum results

c	N_t	$\beta_\star$	16–84% interval	interp. shift	status
0.5	2	2.9895	[2.9888, 2.9903]	0.00276	flagged
0.5	3	4.3069	[4.3059, 4.3081]	0.000383	flagged
0.5	4	5.5846	[5.5821, 5.5868]	0.000685	flagged
1	2	2.5045	[2.5038, 2.5051]	0.000365	flagged
1	3	3.5985	[3.5968, 3.6002]	0.00201	flagged
1	4	4.6305	[4.6279, 4.6326]	0.00352	flagged
1	6	6.6664	[6.6636, 6.6691]	0.0108	flagged
2	2	1.9624	[1.9603, 1.964]	0.00269	flagged
2	3	2.9428	[2.941, 2.9442]	0.00187	flagged
2	4	3.773	[3.7704, 3.7754]	0.00349	flagged

All entries are preliminary and subject to visual review. The interpolation shift is kept separate from the statistical interval.

c	resolutions	Q_0	boot. SE	interp. bound	dof
0.5	2, 3, 4	0.16331	0.000124	0.000295	1
0.5	3, 4	0.15971	0.000331	0.000134	0
1	2, 3, 4, 6	0.26225	0.000158	0.000647	2
1	3, 4, 6	0.25572	0.000261	0.00108	1
2	2, 3, 4	0.45733	0.000768	0.00163	1
2	3, 4	0.41509	0.00153	0.00268	0

The second row for each target omits its coarsest resolution. Detailed flags, full-covariance residual statistics and bootstrap success counts are in the CSV tables. The chosen primary volumes are not a volume extrapolation.

Resolution dependence is the principal limitation of these extrapolations. Omitting $N_t = 2$ moves Q_0 from approximately 0.16331 to 0.15971 at $c = 0.5$, 0.26225 to 0.25572 at $c = 1$, and 0.45733 to 0.41509 at $c = 2$. These shifts are much larger than the conditional bootstrap errors. They are explicitly flagged using the paired-replica uncertainty of the difference between the two fits. The tables retain all requested preliminary fits; their small statistical errors do not establish a controlled continuum mass. Finer resolutions and visual range selection are the priority before a precision continuum claim.

Why earlier results changed

The original resampler included an outer draw of independent chain identities. The corrected method fixes those identities and resamples blocks within each chain. Earlier bootstrap counts and uncertainty prescriptions are preserved in the checksum-verified qbig archive; they are historical products. This reanalysis additionally pools matching short and long runs with fixed used-measurement weights and applies the requested 500-replica policy to all 38 physical ensembles.

Earlier acceptance also imposed fit p-value and window gates. In particular, the $c = 2, N_t = 3, N_s = 32, \beta = 3.25$ mass was unavailable after those gates, despite numerical windows that can be inspected for a plateau. The new analysis retains numerical windows, flags drift and lets visual review determine their use. The coupling fits previously rejected for local log-linearity are displayed in the comparison directories, including their residuals. Keeping these preliminary results increases coverage; their statistical errors remain conditional on the model and selected window.

Next simulations to consider after review

First inspect flagged mass and coupling windows, especially the $\beta = 3.25$ case and the smallest remaining block counts. Additional independent 8,000-sweep chains can improve precision and between-chain checks without adding another length outlier. At otherwise fixed conditions, doubling effective statistics reduces statistical error by approximately $\sqrt{2}$; halving it requires roughly four times the effective statistics. More sweeps cannot resolve a poor fit model or uncontrolled finite-volume effects. At a fixed block size B, obtaining 50 blocks within every chain requires at least its thermalization cut plus $50B$ sweeps: for a 500-sweep cut this is 10,500 sweeps at $B = 200$ and 12,500 at $B = 240$. Adding more 8,000-sweep chains improves combined statistics but leaves that per-chain count unchanged unless the selected block size changes.

The largest current dataset supports three c values, with four resolutions only at $c = 1$. The next coverage priority is an $N_t = 6$ bracket for $c = 0.5$ and $c = 2$, followed by independent confirmation of the tuned couplings and volume/timing checks. No new simulation is authorized by this report.

New $N_t = 1$ matched-coupling diagnostics

c_{ref}	β	N_s	am_G	boot. SE	$\xi_{2\mathrm{nd}}/a$	min. blocks
0.5	2.989543	2048	0.0067043	3.77e-05	146.7	6
1	2.504456	512	0.028794	6.3e-05	40.59	36
2	1.962387	128	0.14111	0.000219	7.348	37

All twelve chains completed with verified inputs, output counts, boundary conditions and checkpoints, using KLFT revision `f47421d895469b72d4af8a9b8479807f2b09a692` and its native A100 build. Total simulation time was 3.038 A100 GPU-hours. Holonomies were measured every sweep, profiles every five sweeps, and Wilson loops were disabled. The separate P100 validation job 290996 also passed all six CTests, CUDA memory checking and the benchmark; it launched no production.

The transverse sizes 128, 512 and 2048 were explicit candidate input choices. The saved planning records label them unvalidated and contain no quantitative derivation establishing their necessity. At the largest coupling, the first-stage estimate is $\xi_{2\mathrm{nd}}/a \simeq 147 \pm 34$, giving $N_s/\xi_{2\mathrm{nd}} \simeq 14$ at the central value. This is preliminary because independent cuts range from 825 to 3625 sweeps, the maximum estimated autocorrelation time is about 317 sweeps, and $B = 635$ leaves only 6–11 complete blocks per chain. The common blocking plateau is unresolved and split-history/between-chain differences require review. At $N_s = 512$, $\xi_{2\mathrm{nd}}/a \simeq 40.6 \pm 2.9$ and $B = 200$ leaves 36–37 blocks; at $N_s = 128$, $\xi_{2\mathrm{nd}}/a \simeq 7.35 \pm 0.23$ and $B = 200$ leaves 37 blocks. These measured lengths explain why matched couplings cannot be interpreted as the source fixed-c targets at $N_t = 1$.

The largest-volume statistics are the immediate concern. At the present block length, 50 complete blocks require 31,750 post-cut sweeps per chain; the most conservative observed cut would bring this to 35,375 total sweeps. This is a planning estimate: autocorrelation and thermalization must be rechecked after extending statistics. A smaller-volume comparison is needed before replacing 2048 by a smaller production size. Adding these untuned points does not extend the fixed-c continuum resolution set.

$N_t = 1$ **near** $\beta/N_t = 1.5$

On 20 October, the user approved four additional $N_t = 1, N_s = 128$ ensembles at $\beta = 1.45, 1.50, 4.6/3, 1.55$, each with four independent 8,000-sweep hot chains. P100 smoke job 291028 and timing array 291029 passed geometry, boundary, cadence and output-hash checks. The projected cost is 0.438 P100 GPU-hours. Production array 291033 completed all sixteen chains on lnode13 and lnode17 by 22:00:55 CEST: 128,000 sweeps and 0.1525 P100 GPU-hours. Output counts, input hashes and physical checkpoints passed verification. Analysis array 291051 uses the same independent-chain cuts, blocking, 500-replica estimator and exhaustive 550-window mass scan as the canonical campaign. These are untuned measurements with `target_c=null`. All four analyses completed successfully. The selected windows are $r = 5$–64 at $\beta = 1.45$ and $r = 4$–64 at the other three couplings; each has 500 successful refits.

β	am_G	boot. SE	$\xi_{2\mathrm{nd}}/a$	boot. SE	min. blocks
1.45	0.45461	0.00123	1.992	0.495	31
1.5	0.41618	0.000643	2.604	0.381	31
1.533333	0.38858	0.00052	3.272	0.307	93
1.55	0.37646	0.000465	2.489	0.413	31

Every chain uses a 500-sweep cut. The selected block lengths in increasing β order are $B = 240, 240, 80, 240$, retaining 31,31,93,31 complete blocks per chain. Three ensembles are below the 50-block target. At the present cut and $B = 240$, that target would require at least 12,500 total sweeps per chain and renewed autocorrelation and blocking checks. These statistics warnings accompany the plotted points. The common observable plateau is unresolved at $\beta = 1.45, 1.50$; $\beta = 1.45$ also has between-chain and split-history warnings. Later-window drift is flagged at $\beta = 1.45, 4.6/3, 1.55$.

Visual inspection at $\beta = 1.55$ shows a stable early-window mass and large-distance correlated excursions, including a negative tail for $r \gtrsim 52$. The selected mass remains preliminary, conditional on its window; increased statistics and tail/window review remain relevant. This observation is retained alongside the automatic split-history flag.

Nine scaling comparisons

All measured points are shown against $y = \beta/N_t$ in three representations, with $p = 0, 1, 3/2$:

$$\widehat{m}N_t^p, \qquad \frac{\xi_{2\mathrm{nd}}/a}{N_t^p}, \qquad \frac{(\xi_{2\mathrm{nd}}/a)^2}{N_t^{2p}}.$$

Thus the squared-length denominators are $1, N_t^2, N_t^3$. Mass and length errors use the saved 500-replica estimates; the squared-length uncertainty uses the structure-factor ratio evaluated on those same joint replicas. The fixed exact scaling factors multiply the corresponding errors. All points use open markers and the same color and shape for a given N_t. The legends contain only N_t; campaign roles and flags remain in the CSVs. The comparison index links the nine individual, grid-free 16:9 PDFs. Nine additional `_zoom.pdf` views restrict $1.2 \le \beta/N_t \le 1.6$, with vertical limits set by the displayed points. Full-range plots are retained; the zoom reuses the same estimates and errors without fitting or resampling. The trial exponent 3/2 is a display choice; no exponent fit is implied.

Matched-y extension near $\beta/N_t = 1.5$

N_t	N_s	β	am_G	boot. SE	$m_G L$	min. blocks
3	96	4.5	0.13865	0.000348	0.41594	93
4	160	6	0.094149	0.000193	0.3766	125
4	160	6.2	0.082128	0.000186	0.32851	93
6	320	9	0.056306	0.000126	0.33784	31

Each $N_t \geq 2$ ensemble has four independent 8,000-sweep hot-start chains, measured every sweep (profiles every five). This batch used 0.2745 A100 GPU-hours on lnode19, preserving one free GPU. Inputs, output counts, checkpoints and raw hashes passed validation. Array 291008 used the same 500-replica estimator and exhaustive window scan as the fixed-c analysis.

These four mass scans completed successfully. Their selected windows have 500 successful refits. The $N_t = 3$ ensemble has a split-history warning; $N_t = 4, \beta = 6$ has between-chain and later-window drift warnings. The $N_t = 6$ window is $r = 6$–160, with later-window drift retained for review. Its $B = 240$ leaves 31 blocks per chain. The mass-error plateau includes $B = 160, 200, 240$, while the common observable plateau remains unresolved. At the current 500-sweep cut, 50 blocks would require 12,500 total sweeps per chain, followed by renewed equilibration and autocorrelation checks. The full-covariance statistic is undefined because its rank is at most 120 for 155 fitted separations. The defined diagonal mass remains included. At $N_t = 4, \beta = 6.2$, the $N_s = 160$ minus 96 mass difference is -0.0003541 ± 0.0003835, or 0.92 bootstrap standard deviations, conditional on the chosen windows.

Visual review of the new $N_t = 6$ correlator shows a positive, flattening tail beyond roughly $r = 90$, above the selected two-image K_0 curve. Later lower-endpoint choices pull the fitted mass downward as its error grows. The origin of that tail remains unresolved. The quoted mass therefore requires window and equilibration review alongside increased statistics; the numerical result and every alternative window are retained.

The purpose is to test the proposed collapse where the screening mass is small, and to check transverse-volume sensitivity. The existing $N_t = 2, N_s = 64, \beta = 3$ point joins the three finer $y = 1.5$ points and the new $N_t = 1, \beta = 1.5$ point. From the Wilson coupling convention,

$$y = \beta/N_t = \frac{4}{a g_{3,0}^2 N_t} = \frac{4}{g_{3,0}^2 L}.$$


Thus fixed y holds the bare slab-coupling coordinate fixed while the resolution changes. Measuring $am_G N_t$ tests its approach to a finite physical screening scale. If that limit is nonzero, am_G ultimately scales as $1/N_t$, up to cutoff and coupling-matching corrections.

The displayed $am_G N_t^{1.5}$ collapse uses the user's trial exponent; no exponent was fitted. For successive resolutions we also tabulate

$$\alpha_{\text{eff}}(N_1, N_2) = \frac{\log[\widehat{m}(N_1)/\widehat{m}(N_2)]}{\log(N_2/N_1)}.$$


Substitution of $\widehat{m} = A N_t^{-\alpha}$ gives $\log[(N_2/N_1)^\alpha]/\log(N_2/N_1) = \alpha$. For measured finite-spacing masses this is a diagnostic log ratio, with errors propagated through the saved 500 replicas. Window, volume and thermalization uncertainties remain separate from these conditional statistical errors. The fixed-c continuum fit continues to use its existing independently tuned inputs. The measured adjacent-resolution powers are 0.7636 ± 0.0034 (1 to 2), 1.4054 ± 0.0077 (2 to 3), 1.3455 ± 0.0111 (3 to 4), and 1.2678 ± 0.0075 (4 to 6). Their drift shows that the trial exponent 1.5 is an approximate visual scaling choice over this range. These finite-spacing ratios inherit the window and statistics warnings; they do not determine an asymptotic exponent.

Lorenzo's trial collapse with all measured points




Reproduction of Lorenzo's supplied $am_G N_t^{3/2}$ versus β/N_t plot, including all seven $N_t = 1$ ensembles. Every marker is open; its color and shape identify N_t. The exponent 3/2 is an assumed illustrative choice. All defined flagged masses remain visible, with bootstrap standard errors conditional on their chosen windows.

Screening mass in slab units at all measured couplings




The ordinate is $m_G L = am_G N_t$, with $L = aN_t$. Each saved mass and its bootstrap standard error are multiplied by the exact temporal extent. All seven $N_t = 1$ ensembles retain their untuned role in the tables. The matched-$y = 1.5$ comparison now includes $N_t = 1, 2, 3, 4, 6$. These additional untuned points leave the fixed-c continuum fits unchanged.

Measured screening scale at $y = 1.5$




The measured resolutions are $N_t = 1, 2, 3, 4, 6$. No fixed-c target is imposed on this comparison and no additional continuum fit is performed.

Second-moment length in slab units




All measured couplings are shown, including all $N_t = 1$ ensembles. The logarithmic scale accommodates the length range. The coordinate is $\xi_{2\mathrm{nd}}/L = (\xi_{2\mathrm{nd}}/a)/N_t$; companion figures show its square and the unscaled and trial-$N_t^{3/2}$ forms, all versus β/N_t. Error bars use the saved 500 joint replicas and retain the volume and statistics limitations. For the squared estimator, evaluate $(\widetilde{C}(0)/\widetilde{C}(p_{\mathrm{min}}) - 1)/[4\sin^2(\pi/N_s)]$ on each replica before taking its SD. Negative noisy squared-moment estimates remain in that distribution. Both central values and errors are in the existing ensemble CSVs.

Transverse-volume check at $N_t = 4, \beta = 6.2$




Independent $N_s = 96$ and 160 ensembles are compared using their selected mass windows. The companion CSV reports the difference and its 500-replica uncertainty; other windows and correlator curves remain available for visual review.

$N_t = 1$: matched-coupling correlators




The three volumes correspond to different couplings. Their folder labels identify the source $N_t = 2$ tuning. Error bars use independent-chain, pooled-with-boundaries bootstrap replicas.

$N_t = 1$: preliminary screening masses




Conditional bootstrap standard errors are shown. Window, blocking, chain-consistency and volume flags remain in the per-ensemble CSVs. These masses are excluded from fixed-c continuum fits.

Continuum comparison: all targets




All three targets use the requested $p = 1$ ansatz. Flags remain attached to their input points.

$c = 0.5$: resolution sensitivity




The drop-coarsest line uses two resolutions and has zero residual degrees of freedom. Bands contain conditional statistical uncertainty only.

$c = 1$: four measured resolutions




The available $N_t = 6$ bracket extends the continuum comparison. Independent tuning confirmation and further cutoff control remain useful.

$c = 2$: preliminary retained trajectory




Mass-window and interpolation concerns remain visible in the CSV flags and per-ensemble plots.

Previously gated mass: $N_t = 3, N_s = 32, \beta = 3.25$



Every numerical mass window at this endpoint is shown. The band marks the suggested mass; overlapping-window stability is a diagnostic heuristic. Other endpoints and the correlator residuals are saved alongside this plot.

Example coupling interpolation



The previously gated mass is retained as a preliminary input. All coupling windows and their residual plots are available in the same comparison directory.

Fixed-c ensemble result register

c	N_t	N_s	β	am_G	SE	window	used sweeps
0.5	2	64	2.9	0.28458	0.000491	3–32	29,760
0.5	2	64	3	0.24513	0.000456	4–32	37,600
0.5	2	64	3.1	0.21427	0.000438	3–32	29,920
0.5	3	80	4.2	0.18478	0.000372	3–40	29,440
0.5	3	80	4.27	0.17315	0.000339	3–40	30,000
0.5	3	80	4.29	0.16951	0.000328	3–40	30,000
0.5	3	80	4.4	0.15159	0.000156	5–40	347,000
0.5	3	80	4.6	0.12704	0.000299	3–40	29,760
0.5	4	96	5.5	0.1334	0.000399	4–48	29,440
0.5	4	96	5.58394149	0.125	0.000371	4–48	29,600
0.5	4	96	5.6	0.12358	0.000333	4–48	29,760
0.5	4	96	5.9	0.10119	0.000325	4–48	30,000
0.5	4	96	6.2	0.082482	0.000326	4–48	29,760
0.5	4	128	5.58394149	0.12562	0.000255	4–64	29,760
1	2	32	2.35	0.6181	0.00139	2–16	29,920
1	2	32	2.5	0.50322	0.000599	1–16	30,000
1	2	32	2.65	0.40906	0.000958	2–16	29,760
1	3	48	3.5	0.36806	0.000895	3–24	30,000
1	3	48	3.7	0.3004	0.000733	3–24	29,920
1	3	48	3.9	0.2481	0.00064	3–24	30,000
1	4	48	4.6	0.25524	0.000661	3–24	29,920
1	4	48	4.61755576	0.25314	0.000718	3–24	30,000
1	4	48	4.95	0.19764	0.000578	3–24	29,760
1	4	48	5.3	0.15342	0.000614	3–24	29,760
1	4	96	4.95	0.19631	0.00042	4–48	30,000
1	4	96	5.3	0.15284	0.000385	4–48	30,000
1	6	96	6.4	0.19015	0.000364	4–48	29,760
1	6	96	6.7	0.16289	0.000486	5–48	30,000
1	6	96	7	0.14187	0.000362	4–48	29,760
2	2	32	1.9	1.0772	0.00403	2–16	29,440
2	2	32	2	0.95897	0.00322	2–16	29,920
2	2	48	2.1	0.83975	0.00392	3–24	37,920
2	3	32	2.85	0.73663	0.00198	2–16	29,760
2	3	32	3.05	0.59445	0.00143	2–16	29,920
2	3	32	3.25	0.47481	0.0019	3–16	30,000
2	4	32	3.7	0.53286	0.00134	2–16	29,920
2	4	32	4	0.40814	0.0016	3–16	30,000
2	4	32	4.3	0.323	0.00125	3–16	29,920

The untuned diagnostic registers are in their respective sections above. Uncertainties are bootstrap standard errors of am_G. The selected window and used statistics are included so a result can be traced to its CSV and raw-input hashes. The complete chain-level cuts, autocorrelation times, flags and block counts are in `chains.csv` and `autocorrelations.csv`; no chain histories are concatenated.

References

[1] M. Nebbia, *Effects of Dirichlet Boundary Conditions in Yang–Mills Theories*, MSc thesis (2024/2025), local reference `docs/marconebbia_tesi.pdf`.

[2] M. Teper, SU(N) Yang–Mills theory in $2 + 1$ dimensions, Phys. Rev. D 59, 014512 (1999), hep-lat/9804008.

[3] A. D. Sokal, Monte Carlo Methods in Statistical Mechanics: Foundations and New Algorithms, in *Functional Integration*, NATO ASI B 361 (1997), pp. 131–192.

[4] B. Efron and R. J. Tibshirani, *An Introduction to the Bootstrap*, Chapman & Hall (1993), chapters 6 and 8.