

Bilinear pseudodifferential operators and Leibniz-type rules

by

Alexander Thomson

B.S., Missouri State University, 2013

M.S., Kansas State University, 2015

---

AN ABSTRACT OF A DISSERTATION

submitted in partial fulfillment of the  
requirements for the degree

DOCTOR OF PHILOSOPHY

Department of Mathematics  
College of Arts and Sciences

KANSAS STATE UNIVERSITY  
Manhattan, Kansas

2019

# Abstract

Enter the text of your abstract in the abstract.tex file. Be sure to delete the text below before you submit your ETDR.

This template uses a separate file for each section of your ETDR: title page, abstract, preface, chapters, reference, etc. This makes it easier to organize and work with a lengthy document. The template is configured with page margins required by the Graduate School and will automatically create a table of contents, lists of tables and figures, and PDF bookmarks.

The file etdrtemplate.tex is the "master" file for the ETDR template. This is the file you need to process with PDFLaTeX in order to produce a PDF version of your ETDR. See the comments in the etdrtemplate.tex and other files for details on using the template. You are not required to use the template, but it can save time and effort in making sure your ETDR meets the Graduate School formatting requirements.

Although the template gives you a foundation for creating your ETDR, you will need a working knowledge of LaTeX in order to produce a final document. You should be familiar with LaTeX commands for formatting text, equations, tables, and other elements you will need to include in your ETDR.

Bilinear pseudodifferential operators and Leibniz-type rules.

by

Alexander Thomson

B.S., Missouri State University, 2013

M.S., Kansas State University, 2015

---

A DISSERTATION

submitted in partial fulfillment of the  
requirements for the degree

DOCTOR OF PHILOSOPHY

Department of Mathematics  
College of Arts and Sciences

KANSAS STATE UNIVERSITY  
Manhattan, Kansas

2019

Approved by:

Major Professor  
Enter Your Major Professor's Name

# Abstract

Enter the text of your abstract in the `abstract.tex` file. Be sure to delete the text below before you submit your ETDR.

This template uses a separate file for each section of your ETDR: title page, abstract, preface, chapters, reference, etc. This makes it easier to organize and work with a lengthy document. The template is configured with page margins required by the Graduate School and will automatically create a table of contents, lists of tables and figures, and PDF bookmarks.

The file `etdrtemplate.tex` is the "master" file for the ETDR template. This is the file you need to process with PDFLaTeX in order to produce a PDF version of your ETDR. See the comments in the `etdrtemplate.tex` and other files for details on using the template. You are not required to use the template, but it can save time and effort in making sure your ETDR meets the Graduate School formatting requirements.

Although the template gives you a foundation for creating your ETDR, you will need a working knowledge of LaTeX in order to produce a final document. You should be familiar with LaTeX commands for formatting text, equations, tables, and other elements you will need to include in your ETDR.

# Table of Contents

List of Figures . . . . .	vii
List of Tables . . . . .	viii
1 Introduction to Leibniz-type rules . . . . .	1
2 Weighted Leibniz-type rules with applications to scattering properties of PDEs . .	5
2.1 Definitions . . . . .	6
2.1.1 Coifman-Meyer Multipliers . . . . .	6
2.1.2 Function spaces . . . . .	8
2.2 Weighted Leibniz-type rules . . . . .	13
2.2.1 Homogeneous Leibniz-type rules . . . . .	13
2.2.2 Connection to Kato-Ponce inequalities . . . . .	16
2.2.3 Proof of Theorem <a href="#">2.2.1</a> . . . . .	17
2.2.4 Inhomogeneous Leibniz-type rules . . . . .	21
2.3 Leibniz rules in other functions spaces . . . . .	22
2.3.1 Leibniz-type rules in the setting of Lorentz-based Triebel-Lizorkin and Besov spaces . . . . .	23
2.3.2 Morrey spaces . . . . .	31
2.3.3 Variable Lebesgue spaces . . . . .	35
2.4 Applications to scattering properties of PDEs . . . . .	42
Bibliography . . . . .	43
A Title for This Appendix . . . . .	49

# List of Figures

# List of Tables

# Chapter 1

## Introduction to Leibniz-type rules

Leibniz-type rules have been extensively studied due to their connections to partial differential equations which model many real world situations such as shallow water waves and fluid flow. In this chapter we introduce some of the definitions and history of the development of Leibniz-type rules which motivated the results to be discussed in chapters 2 and 3 of this manuscript. First consider the Leibniz rule taught in Calculus courses which expresses the derivatives of a product of functions as a linear combination of derivatives of the functions involved; more specifically, for functions  $f$  and  $g$  sufficiently smooth, it holds that

$$\partial_x^\alpha(fg)(x) = \sum_{\alpha_1+\alpha_2=\alpha} C_{\alpha_1\alpha_2} \partial_x^{\alpha_1} f(x) \partial_x^{\alpha_2} g(x),$$

where  $\alpha \in \mathbb{N}_0^n$  and  $C_{\alpha_1\alpha_2}$  are appropriate constants. The definitions in of the function spaces used below and multiindices  $\alpha \in \mathbb{N}_0^n$  will be discussed in Appendix A [A](#). In particular one term has all the derivatives on  $f$  and another with  $\alpha$  derivatives on  $g$  and

$$\partial_x^\alpha(fg)(x) = \partial_x^\alpha f(x)g(x) + f(x)\partial_x^\alpha g(x) + \dots$$

In an analogous way, fractional Leibniz rules give estimates of the smoothness and size of a product of functions in terms of the smoothness and size of the factors. For instance, for  $f$



and  $g$  in the Schwartz class  $\mathcal{S}(\mathbb{R}^n)$ , it holds that

$$\|D^s(fg)\|_{L^r} \lesssim \|D^s f\|_{L^{p_1}} \|g\|_{L^{q_1}} + \|f\|_{L^{p_2}} \|D^s g\|_{L^{q_2}}, \quad (1.0.1)$$

where  $1/r = 1/p_1 + 1/q_1 = 1/p_2 + 1/q_2$ ,  $1 < p_1, p_2, q_1, q_2 \leq \infty$ ,  $1/2 < r \leq \infty$ , and  $s > N(1/\min(r, 1) - 1)$  or  $s$  is an even natural number. The homogeneous fractional differentiation operator of order  $s$ ,  $D^s$ , is defined as

$$D^s f(x) = \int_{\mathbb{R}^n} |\xi|^s \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi,$$

where  $\widehat{f}$  is the Fourier transform of  $f$ . For  $s > 0$ , the operator  $D^s$  is naturally understood as taking  $s$  derivatives of its argument. Indeed, in the case  $s = 2$ ,  $D^2 f = c\Delta f$ , where  $\Delta = \sum_{j=1}^n \partial_{x_j}^2$  is the Laplacian operator. Furthermore, if  $s$  is a positive integer,

$$\|D^s f\|_{L^p} \sim \sum_{|\alpha|=s} \|\partial^\alpha f\|_{L^p},$$

where  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$  for  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}_0^n$ .

Another version of (1.0.1) is obtained by using the inhomogeneous sth order fractional differentiation operator  $J^s$ :

$$\|J^s(fg)\|_{L^r} \lesssim \|J^s f\|_{L^{p_1}} \|g\|_{L^{q_1}} + \|f\|_{L^{p_2}} \|J^s g\|_{L^{q_2}}. \quad (1.0.2)$$

Similarly to its homeogenous counterpart, the operator  $J^s$  is defined through the Fourier transform as

$$J^s f(x) = \int_{\mathbb{R}^n} (1 + |\xi|^2)^{\frac{s}{2}} \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

and can be interpreted as taking derivatives up to order  $s$  of  $f$ .

The estimates (1.0.1) and (1.0.2) are also known as Kato-Ponce inequalities due to the foundational work of Kato-Ponce<sup>1</sup>, where the estimate (1.0.2) was proved in the case

$1 < r = p_1 = q_2 < \infty$  and  $p_2 = q_1 = \infty$ , with applications to the Cauchy problem for Euler and Navier-Stokes equations. This result was extended by Gulisashvili-Kon<sup>2</sup>, who showed (1.0.1) and (1.0.2) for the cases  $s > 0$ ,  $1 < r < \infty$ , and  $1 < p_1, p_2, q_1, q_2 \leq \infty$  in connection to smoothing properties of Schrödinger semigroups. Grafakos-Oh<sup>3</sup> and Muscalu-Schlag<sup>4</sup> established the cases for  $1/2 < r \leq 1$  and the case  $r = \infty$  was completed in the work of Bourgain-Li<sup>5</sup> and Grafakos-Maldonado-Naibo<sup>6</sup>. Applications of the estimates (1.0.1) and (1.0.2) to Korteweg-de Vries equations were studied by Christ-Weinstein<sup>7</sup> and Kenig-Ponce-Vega<sup>8</sup>.

In the estimates (1.0.1) and (1.0.2) the two functions  $f$  and  $g$  are related through pointwise multiplication. Throughout the rest of this manuscript we will consider estimates similar to (1.0.1) and (1.0.2) where the two functions are related through a pseudodifferential operator. Let  $\sigma(x, \xi, \eta)$  be a complex-valued, smooth function for  $x, \xi, \eta \in \mathbb{R}^n$ . We define the *bilinear pseudodifferential operator* associated to  $\sigma$ ,  $T_\sigma$ , by

$$T_\sigma(f, g)(x) = \int_{\mathbb{R}^{2n}} \sigma(x, \xi, \eta) \widehat{f}(\xi) \widehat{g}(\eta) e^{-2\pi i x \cdot (\xi + \eta)} d\xi d\eta.$$

We refer to  $\sigma$  as the symbol of the operator  $T_\sigma$ . When  $\sigma$  is independent of  $x$  we call  $\sigma$  the multiplier of the *bilinear multiplier operator*  $T_\sigma$ .

Throughout this manuscript we will study estimates related to (1.0.1) and (1.0.2) associated to bilinear pseudodifferential operators that are of the form

$$\|D^s T_\sigma(f, g)\|_Z \lesssim \|D^s f\|_{X_1} \|g\|_{Y_1} + \|f\|_{X_2} \|D^s g\|_{Y_2}, \quad (1.0.3)$$

$$\|J^s T_\sigma(f, g)\|_Z \lesssim \|J^s f\|_{X_1} \|g\|_{Y_1} + \|f\|_{X_2} \|J^s g\|_{Y_2}, \quad (1.0.4)$$

for a variety of function spaces  $X_1$ ,  $X_2$ ,  $Y_1$ ,  $Y_2$ , and  $Z$ .

The estimates (1.0.3) and (1.0.4) have been extensively studied in a variety of settings. In<sup>9</sup>, Brummer-Naibo studied Leibniz-type rules in function spaces that admit a molecular

decomposition and a  $\varphi$ -transform characterization in the sense of Frazier-Jawerth<sup>10;11</sup>. In the context of Lebesgue spaces and mixed Lebesgue spaces, estimates of the type (1.0.3) were studied in Hart-Torres-Wu<sup>12</sup> for bilinear multiplier operators with minimal smoothness assumptions on the multipliers. Related mapping properties for bilinear pseudodifferential operators with symbols in the bilinear Hörmander classes were studied by Bényi-Torres<sup>13</sup> and Bényi-Nahmod-Torres<sup>14</sup> in the setting of Sobolev spaces, by Bényi<sup>15</sup> in the setting of Besov spaces, and by Naibo<sup>16</sup> and Koezuka-Tomita<sup>17</sup> in the context of Triebel-Lizorkin spaces. Additionally, versions of (1.0.1) and (1.0.2) in weighted Lebesgue spaces were proved in Cruz-Uribe-Naibo<sup>18</sup>, while Brummer-Naibo<sup>19</sup> proved (1.0.3) and (1.0.4) in weighted Lebesgue spaces for Coifman-Meyer multiplier operators.

In chapter two we will present Leibniz-type rules in the settings of Besov and Triebel-Lizorkin spaces based in various function spaces. The techniques used are quite flexible and allow the method of proof to be adapted to many different function spaces.

In chapter three we will present Leibniz-type rules in Besov spaces for bilinear pseudodifferential operators symbols of critical order.

# Chapter 2

## Weighted Leibniz-type rules with applications to scattering properties of PDEs

In this chapter we discuss bilinear multiplier operators associated to Coifman-Meyer multipliers and Leibniz-type rules in the settings of weighted Triebel-Lizorkin and Besov spaces which generalize 1.0.1 and 1.0.2. Additionally we obtain applications of these results to scattering properties of certain systems of partial differential equations. One of the main results in this chapter is in the setting of Triebel-Lizorkin spaces based in weighted Lebesgue spaces and Hardy spaces. It is as follows.

**Theorem 2.0.1.** *For  $m \in \mathbb{R}$ , let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be a Coifman-Meyer multiplier of order  $m$ . Consider  $0 < p, p_1, p_2 \leq \infty$  such that  $1/p = 1/p_1 + 1/p_2$  and  $0 < q \leq \infty$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $0 < p, p_1, p_2 < \infty$  and  $s > \tau_{p,q}(w)$ , it holds that*

$$\|T_\sigma(f, g)\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{F}_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n). \quad (2.0.1)$$

If  $0 < p, p_1, p_2 \leq \infty$  and  $s > \tau_p(w)$ , it holds that

$$\|T_\sigma(f, g)\|_{\dot{B}_{p,q}^s(w)} \lesssim \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{B}_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.0.2)$$

where the Hardy spaces  $H^{p_1}(w_1)$  and  $H^{p_2}(w_2)$  must be replaced by  $L^\infty$  if  $p_1 = \infty$  or  $p_2 = \infty$ , respectively.

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand sides of (2.2.11) and (2.2.12); moreover, if  $w \in A_\infty$ , then

$$\|T_\sigma(f, g)\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p,q}^{s+m}(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{\dot{F}_{p,q}^{s+m}(w)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.0.3)$$

where  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,q}(w)$ .

In the following sections the function spaces and multipliers used in the hypotheses are defined and discussed. We will then remark on corollaries to Theorem (2.2.1) and their connection to the Leibniz-type rules in the previous chapter. theorem and the proof of Theorem (2.2.1). We will then see how the method to prove these results can be adapted for Triebel-Lizorkin and Besov spaces based in other function spaces. Last we obtain applications to scattering properties of certain systems of partial differential equations.

## 2.1 Definitions

### 2.1.1 Coifman-Meyer Multipliers

The symbols used in Theorem (2.2.1) and results later in this chapter are Coifman-Meyer multipliers. Such multipliers are defined as follows.

*Definition 2.1.1.* For  $m \in \mathbb{R}^n$ , a smooth function  $\sigma = \sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , is a *Coifman-Meyer multiplier* of order  $m$  if for all multi-indices  $\alpha, \beta \in \mathbb{N}_0^n$  there exists a positive constant  $C_{\alpha,\beta}$

such that

$$|\partial_\xi^\alpha \partial_\eta^\beta \sigma(\xi, \eta)| \leq C_{\alpha, \beta} (|\xi| + |\eta|)^{m - (|\alpha| + |\beta|)} \quad \forall (\xi, \eta) \in \mathbb{R}^{2n} \setminus \{(0, 0)\}. \quad (2.1.4)$$

We say  $\sigma = \sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , is an *inhomogeneous Coifman-Meyer multiplier* of order  $m$  if for all multi-indices  $\alpha, \beta \in \mathbb{N}_0^n$  there exists a positive constant  $C_{\alpha, \beta}$  such that

$$|\partial_\xi^\alpha \partial_\eta^\beta \sigma(\xi, \eta)| \leq C_{\alpha, \beta} (1 + |\xi| + |\eta|)^{m - (|\alpha| + |\beta|)} \quad \forall (\xi, \eta) \in \mathbb{R}^{2n}. \quad (2.1.5)$$

Operators associated to Coifman-Meyer multipliers have been well studied. For instance the estimate

$$\|T_\sigma(f, g)\|_{L^p} \lesssim \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}$$

holds where  $\sigma$  is a Coifman-Meyer multiplier,  $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ , and  $1 < p_1, p_2 < \infty$ . For more on these multipliers see Coifman-Meyer<sup>20</sup> for background information and  $L^2$  estimates, see also David- Journé<sup>7</sup> and Grafakos-Torres<sup>21</sup> for other work. Weighted estimates in other settings were studied in Grafakos-Torres<sup>22</sup> and Lerner et al.<sup>23</sup> in connection to a more general class of operators called Calderón-Zygmund operators. We also note that in particular Coifman-Meyer multipliers of order  $m$  belong to the bilinear Hörmander class  $\dot{B}S_{1,1}^m$ . These symbols and operators associated to them will be discussed in the following chapter.

For the proof of Theorem (2.2.1) we will use a the following decomposition of Coifman-Meyer multipliers. Let  $\sigma$  be a Coifman-Meyer multiplier of order  $m$ . Fix  $\Psi \in \mathcal{S}(\mathbb{R}^n)$  such that

$$\text{supp}(\widehat{\Psi}) \subseteq \{\xi \in \mathbb{R}^n : \tfrac{1}{2} < |\xi| < 2\} \quad \text{and} \quad \sum_{j \in \mathbb{Z}} \widehat{\Psi}(2^{-j}\xi) = 1 \quad \forall \xi \in \mathbb{R}^n \setminus \{0\};$$

define  $\Phi \in \mathcal{S}(\mathbb{R}^n)$  so that

$$\widehat{\Phi}(0) := 1, \quad \widehat{\Phi}(\xi) := \sum_{j \leq 0} \widehat{\Psi}(2^{-j}\xi) \quad \forall \xi \in \mathbb{R}^n \setminus \{0\}.$$

For  $a, b \in \mathbb{R}^n$ ,  $\Delta_j^{\tau_a \Psi} f$  and  $S_j^{\tau_a \Phi} f$  satisfy  $\widehat{\Delta_j^{\tau_a \Psi} f}(\xi) = \widehat{\tau_a \Psi}(2^{-j}\xi) \widehat{f}(\xi) = e^{2\pi i 2^{-j}\xi \cdot a} \widehat{\Psi}(2^{-j}\xi) \widehat{f}(\xi)$  and  $\widehat{S_j^{\tau_b \Phi} f}(\xi) = \widehat{\tau_b \Phi}(2^{-j}\xi) \widehat{f}(\xi) = e^{2\pi i 2^{-j}\xi \cdot b} \widehat{\Phi}(2^{-j}\xi) \widehat{f}(\xi)$ . By the work of Coifman and Meyer in [20](#), given  $N \in \mathbb{N}$  such that  $N > n$ , it follows that  $T_\sigma = T_\sigma^1 + T_\sigma^2$ , where, for  $f \in \mathcal{S}_0(\mathbb{R}^n)$  ( $f \in \mathcal{S}(\mathbb{R}^n)$  if  $m \geq 0$ ) and  $g \in \mathcal{S}(\mathbb{R}^n)$ ,

$$T_\sigma^1(f, g)(x) = \sum_{a, b \in \mathbb{Z}^n} \frac{1}{(1 + |a|^2 + |b|^2)^N} \sum_{j \in \mathbb{Z}} \mathcal{C}_j(a, b) (\Delta_j^{\tau_a \Psi} f)(x) (S_j^{\tau_b \Phi} g)(x) \quad \forall x \in \mathbb{R}^n, \quad (2.1.6)$$

the coefficients  $\mathcal{C}_j(a, b)$  satisfy

$$|\mathcal{C}_j(a, b)| \lesssim 2^{jm} \quad \forall a, b \in \mathbb{Z}, j \in \mathbb{Z}, \quad (2.1.7)$$

with the implicit constant depending on  $\sigma$ , and an analogous expression holds for  $T_\sigma^2$  with the roles of  $f$  and  $g$  interchanged.

If  $\sigma$  is an inhomogeneous Coifman–Meyer multiplier of order  $m$ , a similar decomposition to (2.1.6) follows with the summation in  $j \in \mathbb{N}_0$  rather than  $j \in \mathbb{Z}$ , with  $\Delta_0^{\tau_a \Psi}$  replaced by  $S_0^{\tau_a \Phi}$  and for  $f, g \in \mathcal{S}(\mathbb{R}^n)$ .

## 2.1.2 Function spaces

### Weighted spaces

*Definition 2.1.2.* A *weight*  $w(x)$  defined on  $\mathbb{R}^n$  is a nonnegative, locally integrable function such that  $0 < w(x) < \infty$  for almost every  $x \in \mathbb{R}^n$ .

Given a weight  $w(x)$  and  $0 < p < \infty$  we define the weighted Lebesgue space  $L^p(w)$  as

the space of all measurable functions satisfying

$$\|f\|_{L^p(w)} = \left( \int_{\mathbb{R}^n} |f(x)|^p w(x) dx \right)^{\frac{1}{p}} < \infty.$$

In the case that  $p = \infty$  we define  $L^\infty(w) = L^\infty$ .

The specific classes of weights in the hypotheses of our results are Muckenhoupt weights.

*Definition 2.1.3.* For  $1 < p < \infty$  the *Muckenhoupt class*  $A_p$  consists of all weights  $w$  on  $\mathbb{R}^n$  satisfying

$$\sup_B \left( \frac{1}{|B|} \int_B w(x) dx \right) \left( \frac{1}{|B|} \int_B w(x)^{-\frac{1}{p-1}} dx \right)^{p-1} < \infty, \quad (2.1.8)$$

where the supremum is taken over all Euclidean balls  $B \subset \mathbb{R}^n$  and  $|B|$  is the Lebesgue measure of  $B$ . For  $p = \infty$  we define  $A_\infty = \cup_{1 < p < \infty} A_p$ .

From this definition it follows that  $A_p \subset A_q$  when  $p \leq q$ . For a  $w \in A_\infty$  we set  $\tau_w = \inf\{\tau \in (1, \infty] : w \in A_\tau\}$ . The condition (2.1.8) is motivated by the following fact: for  $f \in L^p(w)$  the Hardy-Littlewood maximal function  $\mathcal{M}(f)(x)$  is bounded on  $L^p(w)$  if and only if  $w \in A_p$ . That is for  $1 < p < \infty$   $w \in A_p$  if and only if

$$\|\mathcal{M}(f)\|_{L^p(w)} \lesssim \|f\|_{L^p(w)}.$$

Later in this chapter we will use the maximal function  $\mathcal{M}_r(f) = (\mathcal{M}(|f|^r))^{\frac{1}{r}}$ . By the properties for the Hardy-Littlewood maximal function above it holds that for  $0 < r < p$   $\mathcal{M}_r$  is bounded on  $L^p(w)$  for  $w \in A_{p/r}$  and in this case  $0 < r < \frac{p}{\tau_w}$ . The following theorem is a vector valued version of the previous statement called a weighted Fefferman-Stein inequality.

**Theorem 2.1.1.** *If  $0 < p < \infty$ ,  $0 < q \leq \infty$ ,  $0 < r < \min(p, q)$  and  $w \in A_{p/r}$  (i.e.  $0 < r < \min(p/\tau_w, q)$ ), then for all sequences  $\{f_j\}_{j \in \mathbb{Z}}$  of locally integrable functions defined on  $\mathbb{R}^n$ , we have*

$$\left\| \left( \sum_{j \in \mathbb{Z}} |\mathcal{M}_r(f_j)|^q \right)^{\frac{1}{q}} \right\|_{L^p(w)} \lesssim \left\| \left( \sum_{j \in \mathbb{Z}} |f_j|^q \right)^{\frac{1}{q}} \right\|_{L^p(w)},$$



where the implicit constant depends on  $r$ ,  $p$ ,  $q$ , and  $w$  and the summation in  $j$  should be replaced by the supremum in  $j$  if  $q = \infty$ .

For more details on the Muckenhoupt classes see Grafakos<sup>24</sup>.

## **Triebel-Lizorkin and Besov spaces**

Here we describe the function spaces in which (2.2.1) is based and some properties of such spaces.

Let  $\psi$  and  $\varphi$  be functions in  $\mathcal{S}(\mathbb{R}^n)(\mathbb{R}^n)$  satisfying the following conditions:

- $\text{supp}(\widehat{\psi}) \subset \{\xi \in \mathbb{R}^n : \frac{1}{2} < |\xi| < 2\}$ ,
- $|\widehat{\psi}(\xi)| > c$  for all  $\xi$  such that  $\frac{3}{5} < |\xi| < \frac{5}{3}$  for some  $c > 0$ ,
- $\text{supp}(\widehat{\varphi}) \subset \{\xi \in \mathbb{R}^n : |\xi| < 2\}$ ,
- $|\widehat{\varphi}(\xi)| > c$ .

For  $\psi$  supported in an annulus and  $j \in \mathbb{Z}$  we define the operator  $\Delta_j^\psi(f)$  through its Fourier transform as

$$\widehat{\Delta_j^\psi(f)}(\xi) = \psi(2^{-j}\xi)\widehat{f}(\xi)$$

For such  $\psi$  we define the homogeneous Triebel-Lizorkin and Besov spaces as follows.

*Definition 2.1.4.* Let  $s \in \mathbb{R}$ ,  $0 < p < \infty$ , and  $0 < q \leq \infty$ .

- The weighted *homogeneous Triebel-Lizorkin space*  $\dot{F}_{p,q}^s(w)$  consists of all  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$  such that

$$\|f\|_{\dot{F}_{p,q}^s(w)} = \left\| \left( \sum_{j \in \mathbb{Z}} (2^{sj} |\Delta_j^\psi f|)^q \right)^{\frac{1}{q}} \right\|_{L^p(w)} < \infty.$$

- The weighted *homogeneous Besov space*  $\dot{B}_{p,q}^s(w)$  consists of all  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$  such that

$$\|f\|_{\dot{B}_{p,q}^s(w)} = \left( \sum_{j \in \mathbb{Z}} (2^{js} \|\Delta_j^\psi f\|_{L^p(w)})^q \right)^{\frac{1}{q}} < \infty. \quad (2.1.9)$$

Given  $\varphi$ ,  $\psi$ ,  $S_0^\varphi$ , and  $\Delta_j^\psi$  as above the weighted inhomogeneous Triebel-Lizorkin and Besov spaces are defined as follows.

*Definition 2.1.5.* Let  $s \in \mathbb{R}$ ,  $0 < p < \infty$ , and  $0 < q \leq \infty$ .

- The weighted *inhomogeneous Triebel-Lizorkin space*  $F_{p,q}^s(w)$  is the class of all  $f \in \mathcal{S}(\mathbb{R}^n)'$  such that

$$\|f\|_{F_{p,q}^s(w)} = \|S_0^\varphi f\|_{L^p(w)} + \left\| \left( \sum_{j \in \mathbb{N}_0} (2^{sj} |\Delta_j^\psi f|)^q \right)^{\frac{1}{q}} \right\|_{L^p(w)} < \infty.$$

- The weighted *inhomogeneous Besov space*  $\dot{B}_{p,q}^s(w)$  is the class of all  $f \in \mathcal{S}(\mathbb{R}^n)'$  such that

$$\|f\|_{\dot{B}_{p,q}^s(w)} = \|S_0^\varphi f\|_{L^p(w)} + \left( \sum_{j \in \mathbb{N}_0} (2^{js} \|\Delta_j^\psi f\|_{L^p(w)})^q \right)^{\frac{1}{q}} < \infty. \quad (2.1.10)$$

The definitions above are independent of the choice of  $\varphi$  and  $\psi$ . These spaces are generally quasi-Banach spaces and if  $1 \leq p, q < \infty$  they are Banach spaces. These spaces provide a framework to study a variety of other spaces such as Lebesgue, Hardy, Sobolev, and BMO spaces with a unified approach. For a detailed overview of the development of these spaces see Triebel<sup>25</sup>.

For certain  $s$ ,  $p$ , and  $q$  as in the definitions (2.1.4) and (2.1.5) Triebel-Lizorkin and Besov spaces coincide with other well known function spaces. For instance we have the following equivalences where the function spaces are equivalent in norm

$$F_{p,2}^0(w) \simeq H^p(w) \text{ for } 0 < p < \infty, \quad w \in A_\infty,$$

$$F_{p,2}^0(w) \simeq L^p(w) \simeq H^p(w) \text{ for } 1 < p < \infty, \quad w \in A_p,$$

$$F_{p,2}^s(w) \simeq \dot{W}^{s,p}(w) \text{ for } 1 < p < \infty, \quad w \in A_p.$$

Additionally by the lifting property of Triebel-Lizorkin and Besov spaces Theorem

(2.2.1) and other results in this manuscript can be seen as Leibniz-type rules as in Chapter 1. For weighted Triebel-Lizorkin spaces the lifting property is as follows: for  $s$ ,  $p$ , and  $q$  as in (2.1.5) and (2.1.4) and  $w \in A_\infty$  we have that

$$\|f\|_{\dot{F}_{p,q}^s(w)} \simeq \|D^s f\|_{\dot{F}_{p,q}^0(w)} \quad \text{and} \quad \|f\|_{F_{p,q}^s(w)} \simeq \|J^s f\|_{F_{p,q}^0(w)}.$$

The corresponding statement for Besov spaces is: for  $s$ ,  $p$ , and  $q$  as in (2.1.5) and (2.1.4) and  $w \in A_\infty$  we have that

$$\|f\|_{\dot{B}_{p,q}^s(w)} \simeq \|D^s f\|_{\dot{B}_{p,q}^0(w)} \quad \text{and} \quad \|f\|_{B_{p,q}^s(w)} \simeq \|J^s f\|_{B_{p,q}^0(w)}.$$

### Nikol'skij representations for weighted homogeneous and inhomogeneous Triebel-Lizorkin and Besov spaces

An important tool for the proof of Theorem (2.2.1) is the Nikol'skij representation for weighted Triebel-Lizorkin and Besov spaces. Here we state a weighted version of<sup>26</sup> Theorem 3.7 (see also<sup>25</sup> Section 2.5.2).

**Theorem 2.1.2.** *For  $D > 0$ , let  $\{u_j\}_{j \in \mathbb{Z}} \subset \mathcal{S}'(\mathbb{R}^n)$  be a sequence of tempered distributions such that*

$$\text{supp}(\hat{u}_j) \subset B(0, D 2^j) \quad \forall j \in \mathbb{Z}.$$

*If  $w \in A_\infty$ , then the following holds:*

- (i) *Let  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,q}(w)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{L^p(w)(\ell^q)} < \infty$ , then the series  $\sum_{j \in \mathbb{Z}} u_j$  converges in  $\dot{F}_{p,q}^s(w)$  (in  $\mathcal{S}'_0(\mathbb{R}^n)$  if  $q = \infty$ ) and*

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{\dot{F}_{p,q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{L^p(w)(\ell^q)},$$

*where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p$  and  $q$ . An analogous statement,*

with  $j \in \mathbb{N}_0$ , holds true for  $F_{p,q}^s(w)$  (when  $q = \infty$ , the convergence is in  $\mathcal{S}'(\mathbb{R}^n)$ ).

(ii) Let  $0 < p, q \leq \infty$  and  $s > \tau_p(w)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{\ell^q(L^p(w))} < \infty$ , then the series  $\sum_{j \in \mathbb{Z}} u_j$  converges in  $\dot{B}_{p,q}^s(w)$  (in  $\mathcal{S}'_0(\mathbb{R}^n)$  if  $q = \infty$ ) and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{\dot{B}_{p,q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{\ell^q(L^p(w))},$$

where the implicit constant depends only on  $n, D, s, p$  and  $q$ . An analogous statement, with  $j \in \mathbb{N}_0$ , holds true for  $B_{p,q}^s(w)$  (when  $q = \infty$ , the convergence is in  $\mathcal{S}'(\mathbb{R}^n)$ ).

## 2.2 Weighted Leibniz-type rules

### 2.2.1 Homogeneous Leibniz-type rules

In the setting of weighted homogeneous Besov and Triebel-Lizorkin spaces we obtain the following Leibniz-type rule. As we will see in the corollaries to this result it improves the Leibniz-type rule (1.0.2) and has extensions to weighted versions of (1.0.2).

**Theorem 2.2.1.** *For  $m \in \mathbb{R}$ , let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be a Coifman-Meyer multiplier of order  $m$ . Consider  $0 < p, p_1, p_2 \leq \infty$  such that  $1/p = 1/p_1 + 1/p_2$  and  $0 < q \leq \infty$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $0 < p, p_1, p_2 < \infty$  and  $s > \tau_{p,q}(w)$ , it holds that*

$$\|T_\sigma(f, g)\|_{\dot{B}_{p,q}^s(w)} \lesssim \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{B}_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n). \quad (2.2.11)$$

If  $0 < p, p_1, p_2 \leq \infty$  and  $s > \tau_p(w)$ , it holds that

$$\|T_\sigma(f, g)\|_{\dot{B}_{p,q}^s(w)} \lesssim \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{B}_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.2.12)$$

where the Hardy spaces  $H^{p_1}(w_1)$  and  $H^{p_2}(w_2)$  must be replaced by  $L^\infty$  if  $p_1 = \infty$  or  $p_2 = \infty$ , respectively.

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand sides of (2.2.11) and (2.2.12); moreover, if  $w \in A_\infty$ , then

$$\|T_\sigma(f, g)\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p,q}^{s+m}(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{\dot{F}_{p,q}^{s+m}(w)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.2.13)$$

where  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,q}(w)$ .

We note that if  $m \geq 0$  then the above estimates hold for any  $f, g \in \mathcal{S}(\mathbb{R}^n)(\mathbb{R}^n)$  when  $\mathcal{S}(\mathbb{R}^n)(\mathbb{R}^n)$  is a subspace of the function spaces on the right-hand side. This is the case when  $1 < p_1, p_2 < \infty$ ,  $w_1 \in A_{p_1}$ , and  $w_2 \in A_{p_2}$  in (2.2.11) and (2.2.12) and  $w \in A_p$  for (2.2.13).

By the lifting property of weighted Besov and Triebel-Lizorkin spaces in section and their relation to weighted Hardy spaces in section (2.1.2) the estimates (2.2.11) and (2.2.12) imply the following Leibniz-type rule for Coifman-Meyer multipliers of order zero.

**Corollary 2.2.2.** *Let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be a Coifman-Meyer multiplier of order 0. Consider  $0 < p, p_1, p_2 < \infty$  such that  $1/p = 1/p_1 + 1/p_2$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $s > \tau_p(w)$ , it holds that*

$$\|D^s(T_\sigma(f, g))\|_{H^p(w)} \lesssim \|D^s f\|_{H^{p_1}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|D^s g\|_{H^{p_2}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n). \quad (2.2.14)$$

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand side of (2.2.14); moreover, if  $w \in A_\infty$ , then

$$\|D^s(T_\sigma(f, g))\|_{H^p(w)} \lesssim \|D^s f\|_{H^p(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|D^s g\|_{H^p(w)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.2.15)$$

where  $0 < p < \infty$  and  $s > \tau_p(w)$ .

Corollary (2.2.2) gives estimates related to those in Brummer-Naibo<sup>19</sup> where using different methods the following result was proven

**Theorem 2.2.3.** *if  $\sigma$  is a Coifman-Meyer multiplier of order 0,  $1 < p_1, p_2 \leq \infty$ ,  $\frac{1}{2} < p < \infty$ ,  $1/p = 1/p_1 + 1/p_2$ ,  $w_1 \in A_{p_1}$ ,  $w_2 \in A_{p_2}$ ,  $w = w_1^{p/p_1} w_2^{p/p_2}$  and  $s > \tau_p$ , then for all  $f, g \in \mathcal{S}(\mathbb{R}^n)$  it holds that*

$$\|D^s(T_\sigma(f, g))\|_{L^p(w)} \lesssim \|D^s f\|_{L^{p_1}(w_1)} \|g\|_{L^{p_2}(w_2)} + \|f\|_{L^{p_1}(w_1)} \|D^s g\|_{L^{p_2}(w_2)}. \quad (2.2.16)$$

Moreover, if  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand side of (2.2.16).

Corollary (2.2.2) and Theorem overlap in the following ways.

- The estimate (2.2.14) allows for  $0 < p, p_1, p_2 < \infty$ ,  $w_1, w_2 \in A_\infty$ , and the  $H^P(w)$  on the left-hand side if  $s > \tau_p(w)$ . However (2.2.16) requires  $1 < p_1, p_2 \leq \infty$ ,  $w_1 \in A_{p_1}$ , and  $w_2 \in A_{p_2}$  but allows for the Lebesgue norm on left-hand side when  $s > \tau_p$ . So (2.2.14) is less restrictive than (2.2.16) in terms of the indices  $p$ ,  $p_1$ , and  $p_2$  and the classes that the weights  $w_1$  and  $w_2$  belong to. However because  $\tau_p \leq \tau_p(w)$  (2.2.14) is more restrictive in terms of the range of the regularity  $s$  than (2.2.16).
- If  $s > \tau_p(w)$ ,  $1/2 < p < \infty$ ,  $1 < p_1, p_2 < \infty$  such that  $1/p = 1/p_1 + 1/p_2$ ,  $w_1 \in A_{p_1}$  and  $w_2 \in A_{p_2}$  then (2.2.14) implies (2.2.16). However if  $\tau_p < \tau_p(w)$  then (2.2.14) does not imply (2.2.16) for  $\tau_p < s < \tau_p(w)$ . Here we give examples of weights  $w_1$  and  $w_2$  such that the corresponding weight  $w$  satisfies  $\tau_p < \tau_p(w)$ . Let  $1 < p_1 \leq p_2 < \infty$  and  $w_1(x) = w_2(x) = w(x) = |x|^a$  with  $n(r-1) < a < n(p_1-1)$  for some  $1 < r < p_1$ . Then  $w(x) \in A_{p_1}$ ,  $A_{p_1} \subset A_{p_2}$ , and  $w \notin A_r$ . This implies that  $1 < \tau_w$  which implies that  $\tau_p < \tau_p(w)$ .
- The estimate (2.2.16) implies (2.2.15) for  $1 < p < \infty$ ,  $w \in A_p$ , and  $s > \tau_p$  and gives the endpoint estimate

$$\|D^s(T_\sigma(f, g))\|_{L^p(w)} \lesssim \|D^s f\|_{L^\infty(w)} \|g\|_{L^p(w)} + \|f\|_{L^p(w)} \|D^s g\|_{L^\infty}.$$

However (2.2.15) allows  $0 < p < \infty$  and  $w \in A_\infty$  if  $s > \tau_p(w)$ .

## 2.2.2 Connection to Kato-Ponce inequalities

By setting  $\sigma \equiv 1$  we obtain the following Kato-Ponce inequality as a corollary to Theorem (2.2.1).

**Corollary 2.2.4.** *Consider  $0 < p, p_1, p_2 \leq \infty$  such that  $1/p = 1/p_1 + 1/p_2$  and  $0 < q \leq \infty$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $0 < p, p_1, p_2 < \infty$  and  $s > \tau_{p,q}(w)$ , it holds that*

$$\|fg\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p_1,q}^s(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{F}_{p_2,q}^s(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n). \quad (2.2.17)$$

If  $0 < p, p_1, p_2 \leq \infty$  and  $s > \tau_p(w)$ , it holds that

$$\|fg\|_{\dot{B}_{p,q}^s(w)} \lesssim \|f\|_{\dot{B}_{p_1,q}^s(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|g\|_{\dot{B}_{p_2,q}^s(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.2.18)$$

where the Hardy spaces  $H^{p_1}(w_1)$  and  $H^{p_2}(w_2)$  must be replaced by  $L^\infty$  if  $p_1 = \infty$  or  $p_2 = \infty$ , respectively.

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand sides of (2.2.17) and (2.2.18); moreover, if  $w \in A_\infty$ , then

$$\|fg\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p,q}^s(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{\dot{F}_{p,q}^s(w)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n), \quad (2.2.19)$$

where  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,q}(w)$ .

In particular of we set  $q = 2$  and use the connection between weighted Hardy spaces and weighted Triebel-Lizorkin spaces we obtain the following corollary.

**Corollary 2.2.5.** *Consider  $0 < p, p_1, p_2 < \infty$  such that  $1/p = 1/p_1 + 1/p_2$ ; let  $w_1, w_2 \in A_\infty$*

and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $s > \tau_p(w)$ , it holds that

$$\|D^s(fg)\|_{H^p(w)} \lesssim \|D^s f\|_{H^{p_1}(w_1)} \|g\|_{H^{p_2}(w_2)} + \|f\|_{H^{p_1}(w_1)} \|D^s g\|_{H^{p_2}(w_2)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n). \quad (2.2.20)$$

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand side of (2.2.20); moreover, if  $w \in A_\infty$ , then

$$\|D^s(fg)\|_{H^p(w)} \lesssim \|D^s f\|_{H^p(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|D^s g\|_{H^p(w)} \quad \forall f, g \in \mathcal{S}_0(\mathbb{R}^n),$$

where  $0 < p < \infty$  and  $s > \tau_p(w)$ .

We note that (2.2.20) extends and improves the inequality (1.0.1). The inequality (2.2.20) extends the range of  $p$ ,  $p_1$ , and  $p_2$  by allowing  $0 < p, p_1, p_2 < \infty$  while (1.0.1) requires that  $1 < p_1, p_2 < \infty$ . Additionally (2.2.20) allows for the  $H^p$  norm on the left-hand side which is generally larger than  $\|\cdot\|_{L^p}$ .

### 2.2.3 Proof of Theorem 2.2.1

The following lemma will be useful in the proof of Theorem 2.2.1 and other main results.

**Lemma 2.2.6.** *Let  $\phi_1, \phi_2 \in \mathcal{S}(\mathbb{R}^n)$  be such that  $\widehat{\phi_1}$  and  $\widehat{\phi_2}$  have compact supports and  $\widehat{\phi_1} \widehat{\phi_2} = \widehat{\phi_1}$ . If  $0 < r \leq 1$  and  $\varepsilon > 0$ , it holds that*

$$\left| P_j^{\tau_a \phi_1} f(x) \right| \lesssim (1 + |a|)^{\varepsilon + \frac{n}{r}} \mathcal{M}_r(P_j^{\phi_2} f)(x) \quad \forall x, a \in \mathbb{R}^n, j \in \mathbb{Z}, f \in \mathcal{S}(\mathbb{R}^n).$$

*Proof.* This estimate is a consequence of Lemma 2.3.2. In view of the supports of  $\widehat{\phi_1}$  and  $\widehat{\phi_2}$  we have  $P_j^{\tau_a \phi_1} f = P_j^{\tau_a \phi_1} P_j^{\phi_2} f$  for  $j \in \mathbb{Z}$  and  $f \in \mathcal{S}(\mathbb{R}^n)$ . Applying Lemma 2.3.2 with  $\phi(x) = 2^{nj} \tau_a \phi_1(2^j x)$ ,  $A = 2^j$ ,  $R \geq 1$  such that  $\text{supp}(\widehat{\phi_2}) \subset \{\xi \in \mathbb{R}^n : |\xi| \leq R\}$  and



$d = \varepsilon + n/r$ , we get

$$\begin{aligned} \left| P_j^{\tau_a \phi_1} f(x) \right| &\lesssim R^{n(\frac{1}{r}-1)} 2^{-jn} \left\| (1 + |2^j \cdot|)^{\varepsilon + \frac{n}{r}} 2^{nj} \tau_a \phi_1(2^j \cdot) \right\|_{L^\infty} \mathcal{M}_r(P_j^{\phi_2} f)(x) \\ &\sim \left\| (1 + |2^j \cdot|)^{\varepsilon + \frac{n}{r}} \tau_a \phi_1(2^j \cdot) \right\|_{L^\infty} \mathcal{M}_r(P_j^{\phi_2} f)(x) \quad \forall x, a \in \mathbb{R}^n, j \in \mathbb{Z}, f \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

Since  $\phi_1 \in \mathcal{S}(\mathbb{R}^n)$ ,

$$|\tau_a \phi_1(2^j x)| = |\phi_1(2^j x + a)| \lesssim \frac{(1 + |a|)^{\varepsilon + \frac{n}{r}}}{(1 + |2^j x|)^{\varepsilon + \frac{n}{r}}} \quad \forall x, a \in \mathbb{R}^n, j \in \mathbb{Z}.$$

Combining these two estimates completes the proof.  $\square$

We know proof Theorem 2.2.1.

*Proof of Theorem 2.2.1.* Consider  $\Phi, \Psi, T_\sigma^1, T_\sigma^2, \{\mathcal{C}_j(a, b)\}_{j \in \mathbb{Z}, a, b \in \mathbb{Z}^n}$  as in Section ???. Let  $m, \sigma, p, p_1, p_2, q, s, w_1, w_2$  and  $w$  be as in the hypotheses. For ease of notation,  $p_1$  and  $p_2$  will be assumed to be finite; the same proof applies for (2.2.12) if that is not the case, and for (2.2.13).

We next prove (2.2.11) and (2.2.12). By symmetry, it is enough to work with  $T_\sigma^1$  and prove that

$$\|T_\sigma^1(f, g)\|_{\dot{F}_{p,q}^s(w)} \lesssim \|f\|_{\dot{F}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)} \quad \text{and} \quad \|T_\sigma^1(f, g)\|_{\dot{B}_{p,q}^s(w)} \lesssim \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)}.$$

Moreover, since  $\|\sum f_j\|_{\dot{F}_{p,q}^s(w)}^{\min(p,q,1)} \lesssim \sum \|f_j\|_{\dot{F}_{p,q}^s(w)}^{\min(p,q,1)}$  and similarly for  $\dot{B}_{p,q}^s(w)$ , it suffices to prove that, given  $\varepsilon > 0$  there exist  $0 < r_1, r_2 \leq 1$  such that for all  $g \in \mathcal{S}(\mathbb{R}^n)$  and  $f \in \mathcal{S}_0(\mathbb{R}^n)$  ( $f \in \mathcal{S}(\mathbb{R}^n) \cap \dot{F}_{p,q}^s(w)$  or  $f \in \mathcal{S}(\mathbb{R}^n) \cap \dot{B}_{p,q}^s(w)$  if  $m \geq 0$ ), it holds that

$$\|T^{a,b}(f, g)\|_{\dot{F}_{p,q}^s(w)} \lesssim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \|f\|_{\dot{F}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)}, \quad (2.2.21)$$

$$\|T^{a,b}(f, g)\|_{\dot{B}_{p,q}^s(w)} \lesssim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)}, \quad (2.2.22)$$

where

$$T^{a,b}(f, g) := \sum_{j \in \mathbb{Z}} \mathcal{C}_j(a, b) (\Delta_j^{\tau_a \Psi} f) (S_j^{\tau_b \Phi} g)$$

and the implicit constants are independent of  $a$  and  $b$ . We will assume  $q$  finite; obvious changes apply if that is not the case.

In view of the supports of  $\Psi$  and  $\Phi$  we have that

$$\text{supp}(\mathcal{F}[\mathcal{C}_j(a, b) (\Delta_j^{\tau_a \Psi} f) (S_j^{\tau_b \Phi} g)]) \subset \{\xi \in \mathbb{R}^n : |\xi| \lesssim 2^j\} \quad \forall j \in \mathbb{Z}, a, b \in \mathbb{Z}^n.$$

For (2.2.21), Theorem 2.3.6(2.3.6), the bound (2.1.7) for  $\mathcal{C}_j(a, b)$ , and Hölder's inequality imply

$$\begin{aligned} \|T^{a,b}(f, g)\|_{\dot{F}_{p,q}^s(w)} &\lesssim \left\| \{2^{sj} \mathcal{C}_j(a, b) (\Delta_j^{\tau_a \Psi} f) (S_j^{\tau_b \Phi} g)\}_{j \in \mathbb{Z}} \right\|_{L^p(w)(\ell^q)} \\ &\lesssim \left\| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |(\Delta_j^{\tau_a \Psi} f)(x) (S_j^{\tau_b \Phi} g)|^q \right)^{\frac{1}{q}} \right\|_{L^p(w)} \\ &\leq \left\| \sup_{j \in \mathbb{Z}} |S_j^{\tau_b \Phi} g| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |(\Delta_j^{\tau_a \Psi} f)|^q \right)^{\frac{1}{q}} \right\|_{L^p(w)} \\ &\leq \left\| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |\Delta_j^{\tau_a \Psi} f|^q \right)^{\frac{1}{q}} \right\|_{L^{p_1}(w_1)} \left\| \sup_{j \in \mathbb{Z}} |S_j^{\tau_b \Phi} g| \right\|_{L^{p_2}(w_2)}. \end{aligned}$$

Consider  $\varphi, \psi \in \mathcal{S}(\mathbb{R}^n)$  as in Section 2.1.2 such that  $\widehat{\varphi} \equiv 1$  on  $\text{supp}(\widehat{\Phi})$  and  $\widehat{\psi} \equiv 1$  on  $\text{supp}(\widehat{\Psi})$ .

Let  $0 < r_1 < \min(1, p_1/\tau_{w_1}, q)$ ; by Lemma 2.2.6 and the weighted Fefferman-Stein inequality

we have that

$$\begin{aligned}
\left\| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |(\Delta_j^{\tau_a} \Psi f)|^q \right)^{\frac{1}{q}} \right\|_{L^{p_1}(w_1)} &\lesssim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} \left\| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |\mathcal{M}_{r_1}(\Delta_j^{\psi} f)|^q \right)^{\frac{1}{q}} \right\|_{L^{p_1}(w_1)} \\
&\lesssim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} \left\| \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} |\Delta_j^{\psi} f|^q \right)^{\frac{1}{q}} \right\|_{L^{p_1}(w_1)} \\
&\sim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} \|f\|_{\dot{F}_{p,q}^{s+m}(w_1)},
\end{aligned}$$

where the implicit constants are independent of  $a$  and  $f$ . Next, let  $0 < r_2 < \min(1, p_2/\tau_{w_2})$ ; by Lemma 2.2.6 and the boundedness properties of the Hardy-Littlewood maximal operator on weighted Lebesgue space we have that

$$\begin{aligned}
\left\| \sup_{j \in \mathbb{Z}} |S_j^{\tau_b \Phi} g| \right\|_{L^{p_2}(w_2)} &\lesssim (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \left\| \mathcal{M}_{r_2}(\sup_{j \in \mathbb{Z}} |S_j^{\varphi} g|) \right\|_{L^{p_2}(w_2)} \\
&\lesssim (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \left\| \sup_{j \in \mathbb{Z}} |S_j^{\varphi} g| \right\|_{L^{p_2}(w_2)} \\
&\sim (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \|g\|_{H^{p_2}(w_2)},
\end{aligned}$$

where the implicit constants are independent of  $b$  and  $g$ . Putting all together we obtain (2.2.21).

For (2.2.22), Theorem 2.3.6(2.3.6), the bound (2.1.7) for  $\mathcal{C}_j(a, b)$  and Hölder's inequality give

$$\begin{aligned}
\|T^{a,b}(f, g)\|_{\dot{B}_{p,q}^s(w)} &\lesssim \left\| \{2^{sj} \mathcal{C}_j(a, b) (\Delta_j^{\tau_a} \Psi f) (S_j^{\tau_b \Phi} g)\}_{j \in \mathbb{Z}} \right\|_{\ell^q(L^p(w))} \\
&\lesssim \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} \left\| (\Delta_j^{\tau_a} \Psi f) (S_j^{\tau_b \Phi} g) \right\|_{L^p(w)}^q \right)^{\frac{1}{q}} \\
&\leq \left( \sum_{j \in \mathbb{Z}} 2^{(s+m)qj} \left\| (\Delta_j^{\tau_a} \Psi f) \right\|_{L^{p_1}(w_1)}^q \right)^{\frac{1}{q}} \left\| \sup_{k \in \mathbb{Z}} |S_k^{\tau_b \Phi} g| \right\|_{L^{p_2}(w_2)} \\
&\lesssim (1 + |a|)^{\varepsilon + \frac{n}{r_1}} (1 + |b|)^{\varepsilon + \frac{n}{r_2}} \|f\|_{\dot{B}_{p_1,q}^{s+m}(w_1)} \|g\|_{H^{p_2}(w_2)},
\end{aligned}$$

where in the last inequality we have used Lemma 2.2.6 and the boundedness properties of  $\mathcal{M}$  with  $0 < r_j < \min(1, p_j/\tau_{w_j})$  for  $j = 1, 2$ .

It is clear from the proof above that if  $w_1 = w_2$ , then different pairs of  $p_1, p_2$  related to  $p$  through the Hölder condition can be used on the right-hand sides of (2.2.11) and (2.2.12); in such case  $w = w_1 = w_2$ .  $\square$

## 2.2.4 Inhomogeneous Leibniz-type rules

In this section we obtain Leibniz-type rules for Coifman-Meyer multiplier operators associated to inhomogeneous symbols. Our main result is the inhomogeneous counterpart to Theorem 2.2.1.

**Theorem 2.2.7.** *For  $m \in \mathbb{R}$ , let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be an inhomogeneous Coifman-Meyer multiplier of order  $m$ . Consider  $0 < p, p_1, p_2 \leq \infty$  such that  $1/p = 1/p_1 + 1/p_2$  and  $0 < q \leq \infty$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $0 < p, p_1, p_2 < \infty$  and  $s > \tau_{p,q}(w)$ , it holds that*

$$\|T_\sigma(f, g)\|_{F_{p,q}^s(w)} \lesssim \|f\|_{F_{p_1,q}^{s+m}(w_1)} \|g\|_{h^{p_2}(w_2)} + \|f\|_{h^{p_1}(w_1)} \|g\|_{F_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n). \quad (2.2.23)$$

If  $0 < p, p_1, p_2 \leq \infty$  and  $s > \tau_p(w)$ , it holds that

$$\|T_\sigma(f, g)\|_{B_{p,q}^s(w)} \lesssim \|f\|_{B_{p_1,q}^{s+m}(w_1)} \|g\|_{h^{p_2}(w_2)} + \|f\|_{h^{p_1}(w_1)} \|g\|_{B_{p_2,q}^{s+m}(w_2)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n), \quad (2.2.24)$$

where the local Hardy spaces  $h^{p_1}(w_1)$  and  $h^{p_2}(w_2)$  must be replaced by  $L^\infty$  if  $p_1 = \infty$  or  $p_2 = \infty$ , respectively.

If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand sides of (2.2.23) and (2.2.24); moreover, if  $w \in A_\infty$ , then

$$\|T_\sigma(f, g)\|_{F_{p,q}^s(w)} \lesssim \|f\|_{F_{p,q}^{s+m}(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{F_{p,q}^{s+m}(w)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n),$$

where  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,q}(w)$ .

The corollaries that follow are similar to those in the homogeneous setting. As an example we state the inhomogeneous counterpart to Corollary 2.2.2.

**Corollary 2.2.8.** *Let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be an inhomogeneous Coifman-Meyer multiplier of order 0. Consider  $0 < p, p_1, p_2 < \infty$  such that  $1/p = 1/p_1 + 1/p_2$ ; let  $w_1, w_2 \in A_\infty$  and set  $w = w_1^{p/p_1} w_2^{p/p_2}$ . If  $s > \tau_p(w)$ , it holds that*

$$\|J^s(T_\sigma(f, g))\|_{h^p(w)} \lesssim \|J^s f\|_{h^{p_1}(w_1)} \|g\|_{h^{p_2}(w_2)} + \|f\|_{h^{p_1}(w_1)} \|J^s g\|_{h^{p_2}(w_2)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n).$$

*If  $w_1 = w_2$  then different pairs of  $p_1, p_2$  can be used on the right-hand side of (2.2.14); moreover, if  $w \in A_\infty$ , then*

$$\|J^s(T_\sigma(f, g))\|_{h^p(w)} \lesssim \|J^s f\|_{h^p(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|J^s g\|_{h^p(w)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n),$$

*where  $0 < p < \infty$  and  $s > \tau_p(w)$ .*

## 2.3 Leibniz rules in other functions spaces

The method used to prove Theorem (2.2.1) is quite versatile and can be applied to Triebel-Lizorkin and Besov spaces that are based in other quasi-Banach spaces. Here we highlight the important features of Lebesgue spaces that are necessary for the proof of Theorem (2.2.1) to be adapted to other settings. We also define Triebel-Lizorkin and Besov spaces that are based in other quasi-Banach spaces.

The main features of weighted Triebel-Lizorkin and Besov spaces used in the proof of Theorem (2.2.1) are the following:

- (i) there exists  $r > 0$  such that  $\|f + g\|_{F_{p,q}^s(w)}^r \leq \|f\|_{F_{p,q}^s(w)}^r + \|g\|_{F_{p,q}^s(w)}^r$ ; similarly for the weighted inhomogeneous Besov spaces and the weighted homogeneous Triebel-Lizorkin and Besov spaces;

- (ii) Hölder's inequality in weighted Lebesgue spaces;
- (iii) the boundedness properties in weighted Lebesgue spaces of the Hardy–Littlewood maximal operator (for the Besov space setting) and the weighted Fefferman–Stein inequality (for the Triebel–Lizorkin space setting);
- (iv) Nikol'skij representations for weighted Triebel–Lizorkin and Besov spaces (Theorem 2.3.6).

In the following subsections we consider quasi-Banach spaces  $\mathcal{X}$  such that properties (2.3)–(2.3) hold for the homogeneous and inhomogeneous  $\mathcal{X}$ -based Triebel–Lizorkin and Besov spaces. Corresponding versions of Theorems (2.2.1) and 2.2.7 hold in Triebel–Lizorkin and Besov spaces based in these spaces. The homogeneous  $\chi$ -based Triebel–Lizorkin and Besov spaces denoted by  $\dot{F}_{\mathcal{X},q}^s$  and  $\dot{B}_{\mathcal{X},q}^s$  respectively are defined similarly to the weighted, homogeneous Triebel–Lizorkin and Besov spaces with the  $\|\cdot\|_{L^p(w)}$  quasi-norm replaced with the  $\|\cdot\|_{\mathcal{X}}$  quasi-norm. The inhomogeneous spaces are defined similarly.

### 2.3.1 Leibniz-type rules in the setting of Lorentz-based Triebel–Lizorkin and Besov spaces

For  $0 < p < \infty$  and  $0 < t \leq \infty$  or  $p = t = \infty$ , and an  $A_\infty$  weight  $w$  defined on  $\mathbb{R}^n$ , we denote by  $L^{p,t}(w)$  the weighted Lorentz space consisting of complex-valued, measurable functions  $f$  defined on  $\mathbb{R}^n$  such that

$$\|f\|_{L^{p,t}(w)} = \left( \int_0^\infty \left( \tau^{\frac{1}{p}} f_w^*(\tau) \right)^t \frac{d\tau}{\tau} \right)^{\frac{1}{t}} < \infty,$$

where  $f_w^*(\tau) = \inf\{\lambda \geq 0 : w_f(\lambda) \leq \tau\}$  with  $w_f(\lambda) = w(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})$ . In the case that  $t = \infty$ ,  $\|f\|_{L^{p,t}(w)} = \sup_{t>0} t^{\frac{1}{p}} f_w^*(t)$ . We note that for  $p = t$  we have  $L^{p,p}(w) = L^p(w)$  and  $L^{1,\infty}(w)$  is the weighted weak  $L^1(w)$  space. For more details on these spaces and their properties see Hunt<sup>27</sup>.

We now turn our attention to the analogues to properties (2.3)-(2.3) in the setting of weighted Lorentz-based Triebel-Lizorkin and Besov spaces.

Property (2.3) follows from the work of Hunt<sup>27</sup> that the quasi norm  $\|\cdot\|_{L^{p,t}(w)}$  is comparable to a quasi-norm  $|||\cdot|||_{L^{p,t}(w)}$  that is subadditive. Therefore we have that the norms  $|||\cdot|||_{\dot{F}_{(p,t),q}^s}$  and  $\|\cdot\|_{\dot{F}_{(p,t),q}^s}$  are comparable where  $|||\cdot|||_{\dot{F}_{(p,t),q}^s}$  is the quasi-norm with  $\|\cdot\|_{L^{p,t}(w)}$  replaced by  $|||\cdot|||_{L^{p,t}(w)}$  and for some  $r > 0$   $|||f+g|||_{\dot{F}_{(p,t),q}^s}^r \leq |||f|||_{\dot{F}_{(p,t),q}^s}^r + |||g|||_{\dot{F}_{(p,t),q}^s}^r$ .

By using the following version of Hölder's inequality we obtain property (2.3) [Theorem 4.5,<sup>27</sup>]: Let  $f, g \in L^{p,t}(w)$ . Then for  $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$  and  $\frac{1}{t} = \frac{1}{t_1} + \frac{1}{t_2}$  it holds that

$$\|fg\|_{L^{p,t}(w)} \lesssim \|f\|_{L^{p_1,t_1}(w)} \|g\|_{L^{p_2,t_2}(w)}.$$

The corresponding version of property (2.3) is: If  $0 < p < \infty$ ,  $0 < t, q \leq \infty$ ,  $0 < r < \min(p/\tau_w, q)$  and  $0 < r \leq t$ , it holds that

$$\left\| \left( \sum_{j \in \mathbb{Z}} |\mathcal{M}_r(f_j)|^q \right)^{\frac{1}{q}} \right\|_{L^{p,t}(w)} \lesssim \left\| \left( \sum_{j \in \mathbb{N}_0} |f_j|^q \right)^{\frac{1}{q}} \right\|_{L^{p,t}(w)} \quad \forall \{f_j\}_{j \in \mathbb{N}_0} \in L^{p,t}(w)(\ell^q); \quad (2.3.25)$$

in particular, if  $0 < r < p/\tau_w$  and  $0 < r \leq t$ , it holds that

$$\|\mathcal{M}_r(f)\|_{L^{p,t}(w)} \lesssim \|f\|_{L^{p,t}(w)} \quad \forall f \in L^{p,t}(w).$$

This is established by the extrapolation theorem<sup>28</sup> Theorem 4.10 and comments on page 70 when  $r = 1$ ,  $1 < p < \infty$ ,  $1 \leq t \leq \infty$  and  $1 < q \leq \infty$ . The remaining cases are shown using the previous cases and the following scaling property for Lorentz spaces: For  $0 < s < \infty$   $|||f|||_{L^{p,t}(w)}^s = |||f|||_{L^{sp,st}(w)}^s$ .

The substitute for property 2.3 is the following Nikols'kij representation in weighted Lorentz spaces.

**Theorem 2.3.1.** *For  $D > 0$ , let  $\{u_j\}_{j \in \mathbb{N}_0} \subset \mathcal{S}'(\mathbb{R}^n)$  be a sequence of tempered distributions*

such that

$$\text{supp}(\widehat{u}_j) \subset B(0, D 2^j) \quad \forall j \in \mathbb{N}_0.$$

If  $w \in A_\infty$ , then the following holds:

(i) Let  $0 < p < \infty$ ,  $0 < t, q \leq \infty$  and  $s > \left( \frac{1}{\min(p/\tau_w, t, q, 1)} - 1 \right)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{L^{(p,t)}(w)(\ell^q)} < \infty$ , then the series  $\sum_{j \in \mathbb{N}_0} u_j$  converges in  $F_{(p,t),q}^s(w)$  (in  $\mathcal{S}'(\mathbb{R}^n)$  if  $q = \infty$ ) and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{F_{(p,t),q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{L^{(p,t)}(w)(\ell^q)},$$

where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p$  and  $q$ .

(ii) Let  $0 < p, q \leq \infty$  and  $s > \tau_{p,t}(w)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(L^{(p,t)}(w))} < \infty$ , then the series  $\sum_{j \in \mathbb{N}_0} u_j$  converges in  $B_{(p,t),q}^s(w)$  (in  $\mathcal{S}'(\mathbb{R}^n)$  if  $q = \infty$ ) and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{B_{(p,t),q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(L^{(p,t)}(w))},$$

where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p$  and  $q$ .

For the proof of Theorem 2.3.1 we need the following lemmas.

**Lemma 2.3.2** (Particular case of Corollary 2.11 in <sup>26</sup>). Suppose  $0 < r \leq 1$ ,  $A > 0$ ,  $R \geq 1$  and  $d > n/r$ . If  $\phi \in \mathcal{S}(\mathbb{R}^n)$  and  $f$  is such that  $\text{supp}(\widehat{f}) \subset \{\xi \in \mathbb{R}^n : |\xi| \leq AR\}$ , it holds that

$$|\phi * f(x)| \lesssim R^{n(\frac{1}{r}-1)} A^{-n} \|(1 + |A \cdot|)^d \phi\|_{L^\infty} \mathcal{M}_r f(x) \quad \forall x \in \mathbb{R}^n,$$

where the implicit constant is independent of  $A$ ,  $R$ ,  $\phi$ , and  $f$ .

*Remark 2.3.1.* <sup>26</sup> Corollary 2.11 incorrectly states  $A^{-n/r}$  instead of  $A^{-n}$ . Also, it states  $A \geq 1$ , but the result is true for  $A > 0$  as stated in Lemma 2.3.2.



**Lemma 2.3.3.** *Suppose  $w \in A_\infty$ ,  $0 < p \leq \infty$ ,  $A > 0$ ,  $R \geq 1$ , and  $d > b > n/\min(1, p/\tau_w)$ . If  $\phi \in \mathcal{S}(\mathbb{R}^n)$  and  $f$  is such that  $\text{supp}(\widehat{f}) \subset \{\xi \in \mathbb{R}^n : |\xi| \leq AR\}$ , it holds that*

$$\|\phi * f\|_{L^p(w)} \lesssim R^{b-n} A^{-n} \|(1 + |A \cdot|^d) \phi\|_{L^\infty} \|f\|_{L^p(w)},$$

where the implicit constant is independent of  $A$ ,  $R$ ,  $\phi$  and  $f$ .

*Proof.* Set  $r := n/b < \min(1, p/\tau_w)$ . The hypothesis  $d > b$  means  $d > n/r$  and Lemma 2.3.2 yields

$$|\phi * f(x)| \lesssim R^{n(\frac{1}{r}-1)} A^{-n} \|(1 + |A \cdot|^d) \phi\|_{L^\infty} \mathcal{M}_r f(x) \quad \forall x \in \mathbb{R}^n.$$

Since  $r < p/\tau_w$ , we have  $\|\mathcal{M}_r f\|_{L^{p,t}(w)} \lesssim \|f\|_{L^{p,t}(w)}$  and therefore

$$\|\phi * f\|_{L^{p,t}(w)} \lesssim R^{n(\frac{1}{r}-1)} A^{-n} \|(1 + |A \cdot|^d) \phi\|_{L^\infty} \|f\|_{L^{p,t}(w)};$$

observing that  $1/r - 1 = (b - n)/n$ , the desired estimate follows.  $\square$

The following lemma is a modified version of<sup>26</sup> Lemma 3.8.

**Lemma 2.3.4.** *Let  $\tau < 0$ ,  $\lambda \in \mathbb{R}$ ,  $0 < q \leq \infty$ , and  $k_0 \in \mathbb{Z}$ . Then, for any sequence  $\{d_j\}_{j \in \mathbb{Z}} \subset [0, \infty)$  it holds that*

$$\left\| \left\{ \sum_{k=k_0}^{\infty} 2^{\tau k} 2^{\lambda(j+k)} d_{j+k} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} \lesssim \left\| \{2^{j\lambda} d_j\}_{j \in \mathbb{Z}} \right\|_{\ell^q},$$

where the implicit constant depends only on  $k_0, \tau, \lambda$  and  $q$ .

*Proof.* Suppose first that  $0 < q \leq 1$ . Then,

$$\begin{aligned}
\left\| \left\{ \sum_{k=k_0}^{\infty} 2^{\tau k} 2^{\lambda(j+k)} d_{j+k} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} &= \left[ \sum_{j \in \mathbb{Z}} \left( \sum_{k=k_0}^{\infty} 2^{\tau k} 2^{\lambda(j+k)} d_{j+k} \right)^q \right]^{\frac{1}{q}} \\
&\leq \left[ \sum_{j \in \mathbb{Z}} \sum_{k=k_0}^{\infty} 2^{\tau q k} 2^{\lambda q(j+k)} d_{j+k}^q \right]^{\frac{1}{q}} = \left[ \sum_{k=k_0}^{\infty} 2^{\tau q k} \sum_{j \in \mathbb{Z}} 2^{\lambda q(j+k)} d_{j+k}^q \right]^{\frac{1}{q}} \\
&= \left( \sum_{k=k_0}^{\infty} 2^{\tau q k} \right)^{\frac{1}{q}} \left\| \{2^{j\lambda} d_j\}_{j \in \mathbb{Z}} \right\|_{\ell^q} = C_{k_0, \tau, q} \left\| \{2^{j\lambda} d_j\}_{j \in \mathbb{Z}} \right\|_{\ell^q},
\end{aligned}$$

where in the last equality we have used that  $\tau < 0$ . If  $1 < q < \infty$  we use Hölder's inequality with  $q$  and  $q'$  to write

$$\begin{aligned}
\left\| \left\{ \sum_{k=k_0}^{\infty} 2^{\tau k} 2^{\lambda(j+k)} d_{j+k} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q} &\leq \left[ \sum_{j \in \mathbb{Z}} \left( \sum_{k=k_0}^{\infty} 2^{\tau k q/2} 2^{\lambda q(j+k)} d_{j+k}^q \right) \left( \sum_{k=k_0}^{\infty} 2^{\tau k q'/2} \right)^{q/q'} \right]^{\frac{1}{q}} \\
&= C_{k_0, \tau, q} \left\| \{2^{j\lambda} d_j\}_{j \in \mathbb{Z}} \right\|_{\ell^q}.
\end{aligned}$$

The case  $q = \infty$  is straightforward. □

We now show the proof of Theorem 2.3.1.

*Proof.* We first establish the theorem for finite families of functions and then extend this result to families that are not necessarily finite. We will do this in the homogeneous settings, with the proof in the inhomogeneous settings being similar. Suppose  $\{u_j\}_{j \in \mathbb{Z}}$  is such that  $u_j = 0$  for all  $j$  except those belonging to some finite subset of  $\mathbb{Z}$ ; this assumption allows us to avoid convergence issues since all the sums considered will be finite.

For Part (2.3.6), let  $D, w, p, q$  and  $s$  be as in the hypotheses. Fix  $0 < r < \min(1, p/\tau_w, q)$  such that  $s > n(1/r - 1)$ ; note that the latter is possible since  $s > \tau_{p,q}(w)$ .

Let  $k_0 \in \mathbb{Z}$  be such that  $2^{k_0-1} < D \leq 2^{k_0}$ , then

$$\text{supp}(\widehat{u}_\ell) \subset B(0, 2^\ell D) \subset B(0, 2^{\ell+k_0}) \quad \forall \ell \in \mathbb{Z}.$$

Define  $u = \sum_{\ell \in \mathbb{Z}} u_\ell$  and let  $\psi$  be as in the definition of  $\dot{F}_{(p,t),q}^s(w)$  in Section ?? . We have

$$\Delta_j^\psi u = \sum_{\ell \in \mathbb{Z}} \Delta_j^\psi u_\ell = \sum_{\ell=j-k_0}^{\infty} \Delta_j^\psi u_\ell = \sum_{k=-k_0}^{\infty} \Delta_j^\psi u_{j+k}. \quad (2.3.26)$$

We will use Lemma 2.3.2 with  $\phi(x) = 2^{jn}\psi(2^j x)$ ,  $f = u_{j+k}$ ,  $A = 2^j > 0$ , and  $R = 2^{k+k_0}$ . (Notice that  $\text{supp}(\widehat{u_{j+k}}) \subset B(0, 2^j 2^{k+k_0})$  and, since  $k \geq -k_0$ , we get  $R \geq 1$ .) Fixing  $d > n/r$  and applying Lemma 2.3.2, we get

$$\begin{aligned} |\Delta_j^\psi u_{j+k}(x)| &\lesssim 2^{k_0 n(\frac{1}{r}-1)} 2^{kn(\frac{1}{r}-1)} 2^{-jn} \left\| (1 + |2^j \cdot|)^d 2^{jn} \psi(2^j \cdot) \right\|_{L^\infty} \mathcal{M}_r(u_{j+k})(x) \\ &\sim 2^{kn(\frac{1}{r}-1)} \left( \sup_{y \in \mathbb{R}^n} (1 + |2^j y|)^d |\psi(2^j y)| \right) \mathcal{M}_r(u_{j+k})(x). \end{aligned}$$

Hence,

$$2^{js} |\Delta_j^\psi u_{j+k}(x)| \lesssim 2^{kn(\frac{1}{r}-1-\frac{s}{n})} 2^{s(j+k)} \mathcal{M}_r(u_{j+k})(x),$$

and then, recalling (2.3.32),

$$2^{js} |\Delta_j^\psi u(x)| \lesssim \sum_{k=-k_0}^{\infty} 2^{kn(\frac{1}{r}-1-\frac{s}{n})} 2^{s(j+k)} \mathcal{M}_r(u_{j+k})(x).$$

Since  $1/r - 1 - s/n < 0$ , Lemma 2.3.4 yields

$$\left\| \{2^{js} |\Delta_j^\psi u|\}_{j \in \mathbb{Z}} \right\|_{L^{p,t}(w)(\ell^q)} \lesssim \left\| \{2^{js} \mathcal{M}_r u_j\}_{j \in \mathbb{Z}} \right\|_{L^{p,t}(w)(\ell^q)}$$

with an implicit constant independent of  $\{u_j\}_{j \in \mathbb{Z}}$ . Applying the weighted Fefferman-Stein inequality to the right-hand side of the last inequality leads to the desired estimate

$$\|u\|_{\dot{F}_{(p,t),q}^s(w)} \lesssim \left\| \{2^{js} u_j\}_{j \in \mathbb{Z}} \right\|_{L^{p,t}(\ell^q)}.$$

Assume that the theorem is true for finite families. Let  $\{u_j\} \subset S'$  be as in the hypothesis.

Then using the theorem for finite families we have that

$$\|U_N - U_M\|_{F_{(p,t),q}^s} \lesssim \|\{2^{js}u_j\}_{M+1 \leq j \leq N}\|_{L^{p,t}(w)(\ell^q)}. \quad (2.3.27)$$

Now we apply the following version of the dominated convergence theorem.

*Suppose  $f_n \rightarrow f$  in measure and  $|f_n(x)| \leq |g(x)|$  for some  $g \in L^{p,t}(w)$ . Then*

$$\lim_{n \rightarrow \infty} \|f_n - f\|_{L^{p,t}(w)} = 0.$$

We need to check that the functions  $U_{N,M} = \left(\sum_{j=M+1}^N (2^{js}u_j)^q\right)^{\frac{1}{q}}$ ,  $U = 0$ , and  $g = \left(\sum_{j=0}^{\infty} (2^{js}u_j)^q\right)^{\frac{1}{q}}$  satisfy the hypotheses of the theorem. Now we need to check that  $f_{N,M} \rightarrow f$  in measure. Let  $A_N = \{x : \sum_{j=N}^{\infty} |2^{js}u_j|^q > \tau^q\}$ . Because  $\{u_j\} \subset L^{(p,t)}(w)(\ell^q)$  we have that  $|A_1| < \infty$ . Then since  $|A_{N+1}| \leq |A_N|$  it follows that  $\lim_{N \rightarrow \infty} |A_N| = |\cap A_N|$ .  $\{u_j\} \subset L^{(p,t)}(w)(\ell^q)$  so  $u_j \rightarrow 0$  as  $j \rightarrow \infty$ . Therefore  $f_N \rightarrow 0$  in measure. This implies that

$$\|U_N - U_M\|_{F_{(p,t),q}^s} \lesssim \|\{2^{js}u_j\}_{M+1 \leq j \leq N}\|_{L^{p,t}(w)(\ell^q)} \rightarrow 0.$$

Then  $\sum_{j=0}^{\infty} u_j$  converges in  $F_{(p,t),q}^s$ .

Now we consider the case where  $q = \infty$ . Then  $\{2^{j(s-\epsilon)}\}_{j \geq 0}$  belongs to  $\ell^1(L^{p,t}(w))$  for any  $\epsilon > 0$ . Then by the case for  $q < \infty$   $\sum_{j=0}^{\infty} u_j$  converges in  $B_{(p,t),1}^s$  and so it converges in  $S'$ . Then by using the case for finite families applied to  $\{u_j\}_{0 \leq j \leq N}$  (this case holds because the lemmas used only depend on the indexes considered and maximal function inequalities in Lorentz spaces) we have that

$$\|U_N\|_{F_{(p,t),q}^s} \lesssim \|\{2^{js}u_j\}_{0 \leq j \leq N}\|_{F_{(p,t),\infty}^s} \leq \|\{2^{js}u_j\}\|_{F_{(p,t),\infty}^s}.$$

Then using the Fatou property finishes the proof. □

With these four properties theorems analogous those earlier in this chapter hold in the setting of weighted Lorentz-based Triebel-Lizorkin and Besov spaces and as an example the analogue to Theorem 2.2.7 in the context of the spaces  $F_{(p,t),q}^s(w)$  is below.

**Theorem 2.3.5.** *For  $m \in \mathbb{R}$ , let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be an inhomogeneous Coifman-Meyer multiplier of order  $m$ . If  $w \in A_\infty$ ,  $0 < p, p_1, p_2 < \infty$  and  $0 < t, t_1, t_2 \leq \infty$  are such that  $1/p = 1/p_1 + 1/p_2$  and  $1/t = 1/t_1 + 1/t_2$ ,  $0 < q \leq \infty$  and  $s > \tau_{p,t,q}(w)$ , it holds that*

$$\|T_\sigma(f, g)\|_{F_{(p,t),q}^s(w)} \lesssim \|f\|_{F_{(p_1,t_1),q}^{s+m}(w)} \|g\|_{h^{p_2,t_2}(w)} + \|f\|_{h^{p_1,t_1}(w)} \|g\|_{F_{(p_2,t_2),q}^{s+m}(w)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n).$$

*Different pairs of  $p_1, p_2$  and  $t_1, t_2$  can be used on the right-hand side of the inequality above.*

*Moreover, if  $w \in A_\infty$ ,  $0 < p < \infty$ ,  $0 < t, q \leq \infty$  and  $s > \tau_{p,t,q}(w)$ , it holds that*

$$\|T_\sigma(f, g)\|_{F_{(p,t),q}^s(w)} \lesssim \|f\|_{F_{(p,t),q}^{s+m}(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{F_{(p,t),q}^{s+m}(w)} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n).$$

By the Fefferman–Stein inequality (2.3.25) the lifting property  $\|f\|_{F_{(p,t),q}^s} \simeq \|J^s f\|_{F_{(p,t),q}^0}$  holds true for  $s \in \mathbb{R}$ ,  $0 < p < \infty$  and  $0 < t, q \leq \infty$ . Then, under the assumptions of Theorem 2.3.5 we obtain, in particular,

$$\|J^s(fg)\|_{F_{(p,t),q}^0(w)} \lesssim \|J^s f\|_{F_{(p_1,t_1),q}^0(w)} \|g\|_{h^{p_2,t_2}(w)} + \|f\|_{h^{p_1,t_1}(w)} \|J^s g\|_{F_{(p_2,t_2),q}^0(w)};$$

$$\|J^s(fg)\|_{F_{(p,t),q}^0(w)} \lesssim \|J^s f\|_{F_{(p,t),q}^0(w)} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|J^s g\|_{F_{(p,t),q}^0(w)}.$$

These last two estimates supplement the results in <sup>18</sup> Theorem 6.1, where related Leibniz-type rules in Lorentz spaces were obtained.

### 2.3.2 Morrey spaces

Given  $0 < p \leq t < \infty$  and  $w \in A_\infty$ , we denote by  $M_p^t(w)$  the weighted Morrey space consisting of functions  $f \in L_{\text{loc}}^p(\mathbb{R}^n)$  such that

$$\|f\|_{M_p^t(w)} = \sup_{B \subset \mathbb{R}^n} w(B)^{\frac{1}{t} - \frac{1}{p}} \left( \int_B |f(x)|^p w(x) dx \right)^{\frac{1}{p}} < \infty,$$

where the supremum is taken over all Euclidean balls  $B$  contained in  $\mathbb{R}^n$ . We note that  $M_p^p(w) = L^p(w)$ . For more details on Morrey spaces see the work Rosenthal–Schmeisser<sup>29</sup> and the references contained therein. The corresponding weighted inhomogeneous Triebel–Lizorkin spaces and inhomogeneous Besov spaces are denoted by  $F_{[p,t],q}^s(w)$  and  $B_{[p,t],q}^s(w)$ , respectively. These Morrey-based Triebel–Lizorkin and Besov spaces are independent of the choice of  $\varphi$  and  $\psi$  given in Section ?? and are quasi-Banach spaces that contain  $\mathcal{S}(\mathbb{R}^n)$  (see the works Kozono–Yamazaki<sup>30</sup>, Mazzucato<sup>31</sup>, Izuki et al.<sup>32</sup> and the references they contain). The corresponding local Hardy spaces are denoted by  $h_p^t(w)$ .

We now show the analogues to properties 2.3–2.3 for weighted Morrey spaces.

$$\|f + g\|_{M_p^t(w)}^r \lesssim \|f\|_{M_p^t(w)}^r + \|g\|_{M_p^t(w)}^r$$

for  $r = \min(1, p)$ . It follows that for  $r := \min(1, p, q)$

$$\|f + g\|_{F_{[p,t],q}^s(w)}^r \lesssim \|f\|_{F_{[p,t],q}^s(w)}^r + \|g\|_{F_{[p,t],q}^s(w)}^r$$

with similar inequalities for inhomogeneous weighted Morry-based Besov spaces and homogeneous weighted Morry-based Triebel–Lizorkin and Besov spaces.

A version of property 2.3 follows from Hölder’s inequality for weighted Lebesgue spaces. For  $0 < p \leq t < \infty$ ,  $0 < p_1 \leq t_1 < \infty$  and  $0 < p_2 \leq t_2 < \infty$  are such that  $1/p = 1/p_1 + 1/p_2$

and  $1/t = 1/t_1 + 1/t_2$ , then

$$\|fg\|_{M_p^t(w)} \leq \|f\|_{M_{p_1}^{t_1}(w)} \|g\|_{M_{p_2}^{t_2}(w)};$$

also, if  $0 < p \leq t < \infty$ ,  $0 < p_1, p_2 < \infty$  are such that  $1/p = 1/p_1 + 1/p_2$  and  $w = w_1^{p/p_1} w_2^{p/p_2}$  for weights  $w_1$  and  $w_2$ , then

$$\|fg\|_{M_p^t(w)} \leq \|f\|_{M_{p_1}^{\frac{p_1 t}{p}}(w_1)} \|g\|_{M_{p_2}^{\frac{p_2 t}{p}}(w_2)}.$$

The analogue to property 2.3 is the following Fefferman-Stein inequality: Let  $0 < p \leq t < \infty$ ,  $0 < q \leq \infty$  and  $0 < r < \min(p/\tau_w, q)$ , then

$$\left\| \left( \sum_{j \in \mathbb{Z}} |\mathcal{M}_r(f_j)|^q \right)^{\frac{1}{q}} \right\|_{M_p^t(w)} \lesssim \left\| \left( \sum_{j \in \mathbb{N}_0} |f_j|^q \right)^{\frac{1}{q}} \right\|_{M_p^t(w)} \quad \forall \{f_j\}_{j \in \mathbb{N}_0} \in M_p^t(w)(\ell^q). \quad (2.3.28)$$

If  $0 < p \leq t < \infty$  and  $0 < r < p/\tau_w$ , then

$$\|\mathcal{M}_r(f)\|_{M_p^t(w)} \lesssim \|f\|_{M_p^t(w)} \quad \forall f \in M_p^t(w).$$

The case for  $r = 1$ ,  $1 < p \leq t < \infty$  and  $1 < q \leq \infty$  are shown using extrapolation and the Fefferman-Stein inequality in weighted Lebesgue spaces. For the extrapolation theorem see<sup>29</sup> Theorem 5.3. The remaining cases are shown using that for  $0 < s < \infty$   $\| |f|^s \|_{M_p^t(w)} = \|f\|_{M_{sp}^{st}(w)}^s$  and the previous case.

The Nikol'skij representation for weighted Morrey-based Triebel-Lizorkin and Besov spaces is as follows.

**Theorem 2.3.6.** *For  $D > 0$ , let  $\{u_j\}_{j \in \mathbb{N}_0} \subset \mathcal{S}'(\mathbb{R}^n)$  be a sequence of tempered distributions such that*

$$\text{supp}(\widehat{u}_j) \subset B(0, D 2^j) \quad \forall j \in \mathbb{N}_0.$$

If  $w \in A_\infty$ , then the following holds:

(i) Let  $0 < p \leq t < \infty$ ,  $0 < q \leq \infty$  and  $s > \left(\frac{1}{\min(p/\tau_w, t, q, 1)} - 1\right)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{M_p^t(w)(\ell^q)} < \infty$ , then the series  $\sum_{j \in \mathbb{N}_0} u_j$  converges in  $\mathcal{S}'(\mathbb{R}^n)$  and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{F_{[p, t], q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{M_p^t(w)(\ell^q)},$$

where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p$  and  $q$ .

(ii) Let  $0 < p, q \leq \infty$  and  $s > \tau_{p, t}(w)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(M_p^t(w))} < \infty$ , then the series  $\sum_{j \in \mathbb{N}_0} u_j$  converges in  $B_{[p, t], q}^s(w)$  (in  $\mathcal{S}'(\mathbb{R}^n)$  if  $q = \infty$ ) and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{B_{[p, t], q}^s(w)} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(M_p^t(w))},$$

where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p$  and  $q$ .

The proof of Theorem (2.3.6) uses Lemma 2.3.2, Lemma 2.3.4, and a modified version of Lemma (2.3.3).

**Lemma 2.3.7.** Let  $0 < p \leq t < \infty$ ,  $A > 0$ ,  $R \geq 1$  and  $d > b > \frac{n}{\min(p/\tau_w, 1)}$ . If  $\phi \in \mathcal{S}(\mathbb{R}^n)$  and  $\text{supp}(\widehat{f}) \subset \{\xi : |\xi| \leq AR\}$ . Then

$$\|\phi * f\|_{M_p^t(w)} \lesssim R^{b-n} A^{-n} \|(1 + |A \cdot|^d)\phi\|_{L^\infty} \|f\|_{M_p^t(w)} \quad (2.3.29)$$

with the implicit constant independent of  $R$ ,  $A$ ,  $\phi$ , and  $f$ .

The proof follows along the same lines as Lemma 2.3.2.

*Proof.* Proof of part (ii): Using Lemma 4.3 we have that

$$\left\| \Delta_j^\psi u_{j+k} \right\|_{M_p^t(w)} \lesssim 2^{k(b-n)} \|u_{j+k}\|_{M_p^t(w)}$$



where we have used boundedness of the Hardy-Littlewood maximal function on Morrey spaces when  $0 < r < p/\tau_w$ . Then by setting  $r^* = \min(1, p, q)$  and using property (1) for Morrey spaces

$$2^{jsr^*} \left\| \Delta_j^\psi u \right\|_{M_p^t(w)}^{r^*} \lesssim 2^{jsr^*} \sum_{k=-k_0}^{\infty} \left\| \Delta_j^\psi u_{j+k} \right\|_{M_p^t(w)}^{r^*} = \sum_{k=-k_0}^{\infty} 2^{k(b-n-s)r^*} 2^{sr^*(j+k)} \|u_{j+k}\|_{M_p^t(w)}^{r^*}.$$

Now by applying Lemma A.3 we have

$$\|u\|_{\dot{B}_{[p,t],q}^s(w)} \lesssim \left\| \left\{ \sum_{k=-k_0}^{\infty} 2^{k(b-n-s)r^*} 2^{sr^*(j+k)} \|u_{j+k}\|_{M_p^t(w)}^{p^*} \right\}_{j \in \mathbb{N}_0} \right\|_{\ell^q(w/r^*)}^{\frac{1}{r^*}} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(M_p^t(w))}$$

and part (ii) is shown for the finite family case. For a family that is not necessarily finite we apply the finite family case to  $U_N - U_M := \sum_{j=M+1}^N u_j$ . For finite  $q$  this gives us

$$\|U_N - U_M\|_{\dot{B}_{[p,t],q}^s(w)} \lesssim \|\{2^{js}u_j\}_{M+1 \leq j \leq N}\|_{\ell^q(M_p^t(w))} \leq \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{\ell^q(M_p^t(w))} < \infty$$

and by the dominated convergence theorem the left side converges to 0 as  $M, N \rightarrow \infty$ . So  $U_N$  converges in  $B_{[p,t],q}^s(w)$ . If  $q = \infty$  then  $\{2^{j(s-\epsilon)}\} \in \ell^1(M_p^t(w))$ . By the case for finite  $q$  we have that  $\sum_{j=0}^N u_j$  converges in  $B_{[p,t],q}^{s-\epsilon}(w)$ . Therefore it converges in  $\mathcal{S}'$ .

Proof of part (i): First we will assume that  $u_j = 0$  for all but finitely many  $j$ . From Lemma 4.2 we have that

$$|\Delta_j^\psi u_{j+k}(x)| \lesssim 2^{kn(\frac{1}{r}-1-\frac{s}{n})} 2^{j(s+k)} \mathcal{M}_r(u_{j+k})(x)$$

where  $\Delta_j^\psi u = \sum_{\ell \in \mathbb{N}_0} \Delta_j^\psi u_\ell$ . By Lemma 3.4 and the Fefferman-Stein inequality for Morrey spaces we get

$$\left\| \{2^{js} |\Delta_j^\psi u|\}_{j \in \mathbb{N}_0} \right\|_{M_t^p(w)(\ell^q)} \lesssim \left\| \{2^{js} \mathcal{M}_r u_j\}_{j \in \mathbb{N}_0} \right\|_{M_t^p(w)(\ell^q)} \lesssim \left\| \{2^{js} u_j\}_{j \in \mathbb{N}_0} \right\|_{M_t^p(w)(\ell^q)}$$

for  $0 < r < \min(\frac{p}{\tau_w}, q)$ . Now we prove the theorem for families with are not necessarily

finite. Let  $U_N := \sum_{j=0}^N u_j$ . Then by the finite family case we have that

$$\|u_N\|_{F_{[p,t],q}^s(w)} \lesssim \|\{2^{js}u_j\}_{0 \leq j \leq N}\|_{M_t^p(w)(\ell^q)} \lesssim \|\{2^{js}u_j\}\|_{M_t^p(w)(\ell^q)} < \infty.$$

Assume that  $1 < q < \infty$ . Because  $\|\{2^{js}u_j\}\|_{M_t^p(w)(\ell^q)} < \infty$  we have that  $\sup_j \|\{2^{js}u_j\}\|_{M_t^p(w)} < \infty$  which falls in the  $q = \infty$  case for part (ii). Then  $\sum_{j=0}^\infty u_j$  converges in  $S'$  and from the Fatou property

$$\liminf_{N \rightarrow \infty} \|U_N\|_{F_{[p,t],q}^s(w)} \leq \|U\|_{F_{[p,t],q}^s(w)} v \lesssim \|\{2^{js}u_j\}\|_{M_t^p(w)(\ell^q)}.$$

If  $q = \infty$  then  $\{2^{j(s-\epsilon)}u_j\}_{j \in \mathbb{N}_0}$  is in  $\ell^1(M_p^t(w))$  for any  $\epsilon > 0$ . Then the case for finite  $q$  shows that  $\{2^{j(s-\epsilon)}u_j\}_{j \in \mathbb{N}_0}$  converges in  $B_{[p,t],1}^{s-\epsilon}(w)$  and we have convergence in  $S'$ . Then using the finite family case we have

$$\|U_N\|_{F_{[p,t],\infty}^s(w)} \lesssim \|\{2^{js}u_j\}_{0 \leq j \leq N}\|_{\ell^1(M_p^t(w))} \leq \|\{2^{js}u_j\}\|_{\ell^1(M_p^t(w))} < \infty$$

Then after using the Fatou property of  $F_{[p,t],q}^s(w)$  we are finished.  $\square$

### 2.3.3 Variable Lebesgue spaces

Let  $\mathcal{P}_0$  be the collection of measurable functions  $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$  such that

$$p_- := \operatorname{ess\,inf}_{x \in \mathbb{R}^n} p(x) > 0 \quad \text{and} \quad p_+ := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x) < \infty.$$

For  $p(\cdot) \in \mathcal{P}_0$ , the variable-exponent Lebesgue space  $L^{p(\cdot)}$  consists of all measurable functions  $f$  such that

$$\|f\|_{L^{p(\cdot)}} := \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \left| \frac{f(x)}{\lambda} \right|^{p(x)} dx \leq 1 \right\} < \infty;$$

such quasi-norm turns  $L^{p(\cdot)}$  into a quasi-Banach space (Banach space if  $p_- \geq 1$ ). We note that if  $p(\cdot) = p$  is constant then  $L^{p(\cdot)} \simeq L^p$  with equality of norms and that

$$\| |f|^t \|_{L^{p(\cdot)}} = \|f\|_{L^{tp(\cdot)}}^t \quad \forall t > 0. \quad (2.3.30)$$

Let  $\mathcal{B}$  be the family of all  $p(\cdot) \in \mathcal{P}_0$  such that  $\mathcal{M}$ , the Hardy–Littlewood maximal operator, is bounded from  $L^{p(\cdot)}$  to  $L^{p(\cdot)}$ . Such exponents satisfy  $p_- > 1$  and the following log-Hölder continuity properties

- there exists a constant  $C_0$  such that for all  $x, y \in \mathbb{R}^n$ ,  $|x - y| < 1/2$

$$|p(x) - p(y)| \leq \frac{C_0}{-\log(|x - y|)},$$

- there exist constants  $C_\infty$  and  $p_\infty$  such that for all  $x \in \mathbb{R}^n$

$$|p(x) - p_\infty| \leq \frac{C_\infty}{\log(e + |x|)}.$$

Furthermore if  $\tau_0 > 0$  is such that  $p(\cdot)/\tau_0 \in \mathcal{B}$  then  $p(\cdot)/\tau \in \mathcal{B}$  for  $0 < \tau < \tau_0$ . Indeed, by Jensen's inequality it holds that  $\mathcal{M}(f)^{\tau_0/\tau}(x) \leq \mathcal{M}(|f|^{\tau_0/\tau})(x)$  so by 2.3.30 we have that

$$\begin{aligned} \|\mathcal{M}f\|_{L^{\frac{p(\cdot)}{\tau}}} &\leq \|\mathcal{M}(|f|^{\tau_0/\tau})^{\tau/\tau_0}\|_{L^{\frac{p(\cdot)}{\tau}}} \\ &= \|\mathcal{M}(|f|^{\tau_0/\tau})\|_{L^{\frac{p(\cdot)}{\tau_0}}}^{\tau/\tau_0} \\ &\leq \left\| |f|^{\frac{\tau_0}{\tau}} \right\|_{L^{\frac{p(\cdot)}{\tau}}}^{\frac{\tau}{\tau_0}} \\ &= \|f\|_{L^{\frac{p(\cdot)}{\tau}}}. \end{aligned}$$

We then define

$$\tau_{p(\cdot)} = \sup\{\tau > 0 : \frac{p(\cdot)}{\tau} \in \mathcal{B}\}, \quad p(\cdot) \in \mathcal{P}_0^*,$$

where  $\mathcal{P}_0^*$  denotes the class of variable exponents in  $\mathcal{P}_0$  such that  $p(\cdot)/\tau_0 \in \mathcal{B}$  for some  $\tau_0 > 0$ .

Note that  $\tau_{p(\cdot)} \leq p_-$ .

Given  $s \in \mathbb{R}$ ,  $0 < q \leq \infty$  and  $p(\cdot) \in \mathcal{P}_0$ , the corresponding inhomogeneous Triebel–Lizorkin and Besov spaces are denoted by  $F_{p(\cdot),q}^s$  and  $B_{p(\cdot),q}^s$ , respectively. If  $p(\cdot) \in \mathcal{P}_0^*$ , these spaces are independent of the functions  $\psi$  and  $\varphi$  given in Section 2.1.2 (see Xu<sup>33</sup>), contain  $\mathcal{S}(\mathbb{R}^n)$  and are quasi-Banach spaces. If  $p(\cdot) \in \mathcal{B}$  and  $s > 0$ ,  $F_{p(\cdot),2}^s$  coincides with the variable-exponent Sobolev space  $W^{s,p(\cdot)}$  (see Gurka et al.<sup>34</sup> and Xu<sup>35</sup>). More general versions of variable-exponent Triebel–Lizorkin and Besov spaces, where  $s$  and  $q$  are also allowed to be functions, were introduced in Diening et al.<sup>36</sup> and Almeida–Hästö<sup>37</sup>, respectively. The local Hardy space with variable exponent  $p(\cdot) \in \mathcal{P}_0$ , denoted  $h^{p(\cdot)}$ , is defined analogously to  $h^p(w)$  with the quasi-norm in  $L^p(w)$  replaced by the quasi-norm in  $L^{p(\cdot)}$ .

We now consider the analogues of properties 2.3–2.3 in the setting of variable Lebesgue based spaces.

For 2.3 we apply 2.3.30 to get for  $r = \min(p_-, q, 1)$

$$\|f + g\|_{F_{p(\cdot),q}^s}^r \leq \|f\|_{F_{p(\cdot),q}^s}^r + \|g\|_{F_{p(\cdot),q}^s}^r,$$

$$\|f + g\|_{B_{p(\cdot),q}^s}^r \leq \|f\|_{B_{p(\cdot),q}^s}^r + \|g\|_{B_{p(\cdot),q}^s}^r.$$

To prove 2.3 we use<sup>38</sup> Corollary 2.28 and 2.3.30 to get the following version of Hölder’s inequality: If  $p_1(\cdot)$ ,  $p_2(\cdot)$ ,  $p(\cdot) \in \mathcal{P}_0$  are such that  $1/p(\cdot) = 1/p_1(\cdot) + 1/p_2(\cdot)$  then

$$\|fg\|_{L^{p(\cdot)}} \lesssim \|f\|_{L^{p_1(\cdot)}} \|g\|_{L^{p_2(\cdot)}} \quad \forall f \in L^{p_1(\cdot)}, g \in L^{p_2(\cdot)}.$$

The case when  $p(\cdot) \in \mathcal{P}_0$  has  $p_- \geq 1$  is shown in<sup>38</sup> Corollary 2.28. If  $0 < p_- < 1$  then we use 2.3.30 to get

$$\|fg\|_{L^{p(\cdot)}} = \| |fg|^{p_-} \|_{L^{\frac{p(\cdot)}{p_-}}}^{\frac{1}{p_-}}$$

and then use the first case since  $\frac{p(\cdot)}{p_-} > 1$ .

A Fefferman-Stein inequality in variable exponent Lebesgue spaces follows from the discussion in<sup>38</sup> Section 5.6.8 and (2.3.30). For property 2.3 we have the following: If  $p(\cdot) \in \mathcal{P}_0^*$ ,  $0 < q \leq \infty$  and  $0 < r < \min(\tau_{p(\cdot)}, q)$  then

$$\left\| \left( \sum_{j \in \mathbb{Z}} |\mathcal{M}_r(f_j)|^q \right)^{\frac{1}{q}} \right\|_{L^{p(\cdot)}} \lesssim \left\| \left( \sum_{j \in \mathbb{N}_0} |f_j|^q \right)^{\frac{1}{q}} \right\|_{L^{p(\cdot)}} \quad \forall \{f_j\}_{j \in \mathbb{N}_0} \in L^{p(\cdot)}(\ell^q);$$

in particular, if  $0 < r < \tau_{p(\cdot)}$  it holds that

$$\|\mathcal{M}_r(f)\|_{L^{p(\cdot)}} \lesssim \|f\|_{L^{p(\cdot)}} \quad \forall f \in L^{p(\cdot)}.$$

Property 2.3, the Nikol'skij representation for  $F_{p(\cdot),q}^s$  and  $B_{p(\cdot),q}^s$ , for variable exponent Lebesgue spaces is stated below.

**Theorem 2.3.8.** *For  $D > 0$ , let  $\{u_j\}_{j \in \mathbb{Z}} \subset \mathcal{S}'(\mathbb{R}^n)$  be a sequence of tempered distributions such that  $\text{supp}(\widehat{u}_j) \subset B(0, D2^j)$  for all  $j \in \mathbb{Z}$ . Let  $p(\cdot) \in \mathcal{P}_0^*$ ,  $0 < q \leq \infty$  and  $s > n(1/\min(\tau_{p(\cdot)}, q, 1) - 1)$ . If  $\|\{2^{js}u_j\}_{j \in \mathbb{Z}}\|_{L^{p(\cdot)}(\ell^q)} < \infty$ , then the series  $\sum_{j \in \mathbb{Z}} u_j$  converges in  $F_{p(\cdot),q}^s$  (in  $\mathcal{S}'(\mathbb{R}^n)$  if  $q = \infty$ ) and*

$$\left\| \sum_{j \in \mathbb{N}_0} u_j \right\|_{F_{p(\cdot),q}^s} \lesssim \|\{2^{js}u_j\}_{j \in \mathbb{N}_0}\|_{L^{p(\cdot)}(\ell^q)},$$

where the implicit constant depends only on  $n$ ,  $D$ ,  $s$ ,  $p(\cdot)$  and  $q$ . An analogous statement holds true for  $B_{p(\cdot),q}^s$  with  $s > n(1/\min(\tau_{p(\cdot)}, 1) - 1)$ .

**Lemma 2.3.9.** *Let  $p(\cdot) \in \mathcal{P}_0$ ,  $A > 0$ ,  $R \geq 1$  and  $d > b > \frac{n}{\min(\tau_{p(\cdot)}, 1)}$ . If  $\phi \in \mathcal{S}(\mathbb{R}^n)$  and  $\text{supp}(\widehat{f}) \subset \{\xi : |\xi| \leq AR\}$ . Then*

$$\|\phi * f\|_{L^{p(\cdot)}} \lesssim R^{b-n} A^{-n} \|(1 + |A \cdot|^d) \phi\|_{L^\infty} \|f\|_{L^{p(\cdot)}}, \quad (2.3.31)$$

with the implicit constant independent of  $R$ ,  $A$ ,  $\phi$ , and  $f$ .

*Proof.* We first prove the theorem for finite families. Assume that  $\{u_j\}_{j \in \mathbb{Z}} \subset \mathcal{S}'(\mathbb{R}^n)$  is such that  $u_j \equiv 0$  for all but finitely many  $j$ . Let  $D$ ,  $p(\cdot)$ ,  $q$ , and  $s$  be as in the hypotheses. Fix  $0 < r < \min(1, p_-, q)$  such that  $s > n(1/r - 1)$ ; note that this is possible since  $s > n(1/\min(\tau_{p(\cdot)}, q, 1) - 1)$ . Let  $k_0 \in \mathbb{Z}$  be such that  $2^{k_0-1} < D \leq 2^{k_0}$ , then

$$\text{supp}(\widehat{u}_\ell) \subset B(0, 2^\ell D) \subset B(0, 2^{\ell+k_0}) \quad \forall \ell \in \mathbb{Z}.$$

Define  $u = \sum_{\ell \in \mathbb{Z}} u_\ell$  and let  $\psi$  be as in the definition of  $\dot{F}_{p(\cdot), q}^s(w)$ . We have

$$\Delta_j^\psi u = \sum_{\ell \in \mathbb{Z}} \Delta_j^\psi u_\ell = \sum_{\ell=j-k_0}^{\infty} \Delta_j^\psi u_\ell = \sum_{k=-k_0}^{\infty} \Delta_j^\psi u_{j+k}. \quad (2.3.32)$$

We will use Lemma 2.3.2 with  $\phi(x) = 2^{jn}\psi(2^j x)$ ,  $f = u_{j+k}$ ,  $A = 2^j > 0$ , and  $R = 2^{k+k_0}$ . (Notice that  $\text{supp}(\widehat{u_{j+k}}) \subset B(0, 2^j 2^{k+k_0})$  and, since  $k \geq -k_0$ , we get  $R \geq 1$ .) Fixing  $d > n/r$  and applying Lemma 2.3.2, we get

$$\begin{aligned} |\Delta_j^\psi u_{j+k}(x)| &\lesssim 2^{k_0 n(\frac{1}{r}-1)} 2^{kn(\frac{1}{r}-1)} 2^{-jn} \left\| (1 + |2^j \cdot|)^d 2^{jn} \psi(2^j \cdot) \right\|_{L^\infty} \mathcal{M}_r(u_{j+k})(x) \\ &\sim 2^{kn(\frac{1}{r}-1)} \left( \sup_{y \in \mathbb{R}^n} (1 + |2^j y|)^d |\psi(2^j y)| \right) \mathcal{M}_r(u_{j+k})(x). \end{aligned}$$

Hence,

$$2^{js} |\Delta_j^\psi u_{j+k}(x)| \lesssim 2^{kn(\frac{1}{r}-1-\frac{s}{n})} 2^{s(j+k)} \mathcal{M}_r(u_{j+k})(x),$$

and then, recalling (2.3.32),

$$2^{js} |\Delta_j^\psi u(x)| \lesssim \sum_{k=-k_0}^{\infty} 2^{kn(\frac{1}{r}-1-\frac{s}{n})} 2^{s(j+k)} \mathcal{M}_r(u_{j+k})(x).$$

Since  $1/r - 1 - s/n < 0$ , Lemma 2.3.4 yields

$$\left\| \{2^{js} |\Delta_j^\psi u|\}_{j \in \mathbb{Z}} \right\|_{L^p(w)(\ell^q)} \lesssim \left\| \{2^{js} \mathcal{M}_r u_j\}_{j \in \mathbb{Z}} \right\|_{L^p(w)(\ell^q)}$$

with an implicit constant independent of  $\{u_j\}_{j \in \mathbb{Z}}$ . Applying the weighted Fefferman-Stein inequality to the right-hand side of the last inequality leads to the desired estimate

$$\|u\|_{\dot{F}_{p,q}^s(w)} \lesssim \left\| \{2^{js} u_j\}_{j \in \mathbb{Z}} \right\|_{L^p(\ell^q)}.$$

For the space  $\dot{B}_{p(\cdot),q}^s()$ , let  $D$ ,  $w$ ,  $p$ ,  $q$  and  $s$  be as in the hypotheses and  $k_0$  be as above. Consider  $\Delta_j^\psi u_{j+k}$  in (2.3.32) and apply Lemma 2.3.3 with  $\phi(x) = 2^{jn} \psi(2^{-j}x)$ ,  $f = u_{j+k}$ ,  $A = 2^j$ ,  $R = 2^{k+k_0}$ ,  $d > b$  and  $n/\min(1, p/\tau_w) < b < n + s$ ; note that such  $b$  exists since  $s > \tau_p(w)$ . We get

$$\left\| \Delta_j^\psi u_{j+k} \right\|_{L^{p(\cdot)}} \lesssim 2^{(k+k_0)(b-n)} 2^{-jn} \left\| (1 + |2^j \cdot|)^d 2^{jn} \psi(2^{-j} \cdot) \right\|_{L^\infty} \|u_{j+k}\|_{L^p(w)} \sim 2^{k(b-n)} \|u_{j+k}\|_{L^{p(\cdot)}},$$

and setting  $p^* := \min(p_-, 1)$  we obtain

$$2^{jsp^*} \left\| \Delta_j^\psi u \right\|_{L^{p(\cdot)}}^{p^*} \lesssim 2^{jsp^*} \sum_{k=-k_0}^{\infty} \left\| \Delta_j^\psi u_{j+k} \right\|_{L^{p(\cdot)}}^{p^*} = \sum_{k=-k_0}^{\infty} 2^{k(b-n-s)p^*} 2^{sp^*(j+k)} \|u_{j+k}\|_{L^{p(\cdot)}}^{p^*}.$$

Hence, applying Lemma 2.3.4, it follows that

$$\|u\|_{\dot{B}_{p(\cdot),q}^s} \lesssim \left\| \left\{ \sum_{k=-k_0}^{\infty} 2^{k(b-n-s)p^*} 2^{sp^*(j+k)} \|u_{j+k}\|_{L^{p(\cdot)}}^{p^*} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^q/p^* \frac{1}{p^*}} \lesssim \left\| \{2^{js} u_j\}_{j \in \mathbb{Z}} \right\|_{\ell^q(L^{p(\cdot)})},$$

as desired. Now for families that are not necessarily finite let  $U_N := \sum_{j=0}^N u_j$ . First we assume that  $0 < q < \infty$ . Then for  $M \leq N$  we have

$$\|U_N - U_M\|_{\dot{F}_{p(\cdot),q}^s} \lesssim \left\| \{2^{js} u_j\}_{M+1 \leq j \leq N} \right\|_{L^{p(\cdot)}(\ell^q)}. \quad (2.3.33)$$

Now we use the following dominated convergence theorem in variable exponent Lebesgue spaces: Suppose  $f_n \rightarrow f$  pointwise a.e. and  $|f_n(x)| \leq |g(x)|$  for some  $g \in L^{p(\cdot)}$ . Then  $f_n \rightarrow f$  in  $L^{p(\cdot)}$ .

Applying this theorem with  $f_{N,M} = \sum_{j=M+1}^N 2^{js} U_j$ ,  $f = 0$ , and  $g = \sum_{j=0}^{\infty} u_j$  to get

$$\|U_N - U_M\|_{\dot{F}_{p(\cdot),q}^s} \lesssim \|\{2^{js} u_j\}_{M+1 \leq j \leq N}\|_{L^{p(\cdot)}(\ell^q)} \rightarrow 0 \text{ as } M, N \rightarrow \infty. \quad (2.3.34)$$

Because  $\dot{F}_{p(\cdot),q}^s$  is a quasi-Banach space it is complete so  $U_N$  converges in  $\dot{F}_{p(\cdot),q}^s$  and

$$\left\| \sum_{j=0}^{\infty} \right\|_{\dot{F}_{p(\cdot),q}^s} \lesssim \|\{2^{js} u_j\}\|_{L^{p(\cdot)}(\ell^q)}. \quad (2.3.35)$$

If  $q = \infty$ , we use that  $\{2^{(s-\varepsilon)j} u_j\}_{j \geq 0}$  and  $\{2^{(s+\varepsilon)j} u_j\}_{j < 0}$  belong to  $\ell^1(L^{p(\cdot)})$  for any  $\varepsilon > 0$  and apply Theorem 2.3.6 under the case of finite  $q$  to conclude that  $\sum_{j=0}^N u_j$  and  $\sum_{j=-N}^{-1} u_j$  converge in  $\dot{B}_{p(\cdot),1}^{s-\varepsilon}$  and  $\dot{B}_{p(\cdot),1}^{s+\varepsilon}$ , respectively (choosing  $\varepsilon > 0$  so that  $s - \varepsilon > n(1/\min(\tau_{p(\cdot)}, 1) - 1)$ ). Therefore,  $U_N$  convergence in  $\mathcal{S}'(\mathbb{R}^n)$ . Moreover, by Theorem 2.3.6 applied to the finite sequence  $\{u_j\}_{-N \leq j \leq N}$ , we have that  $U_N \in \dot{F}_{p(\cdot),\infty}^s$  and

$$\|U_N\|_{\dot{F}_{p(\cdot),\infty}^s} \lesssim \|\{2^{js} u_j\}_{-N \leq j \leq N}\|_{L^{p(\cdot)}(\ell^\infty)} \leq \|\{2^{js} u_j\}_{j \in \mathbb{Z}}\|_{L^{p(\cdot)}(\ell^\infty)},$$

with the implicit constant independent of  $N$  and  $\{u_j\}_{j \in \mathbb{Z}}$ . Since  $\dot{F}_{p(\cdot),\infty}^s$  has the Fatou property, we conclude that  $\lim_{N \rightarrow \infty} U_N = \sum_{j \in \mathbb{Z}} u_j$  belongs to  $\dot{F}_{p(\cdot),\infty}^s$  and

$$\left\| \sum_{j \in \mathbb{Z}} u_j \right\|_{\dot{F}_{p(\cdot),\infty}^s} \lesssim \|\{2^{js} u_j\}_{j \in \mathbb{Z}}\|_{L^{p(\cdot)}(\ell^\infty)}.$$

□

As a model result the we state the Leibniz type rule for variable exponent Triebel-Lizorkin spaces.



**Theorem 2.3.10.** *For  $m \in \mathbb{R}$ , let  $\sigma(\xi, \eta)$ ,  $\xi, \eta \in \mathbb{R}^n$ , be an inhomogeneous Coifman-Meyer multiplier of order  $m$ . If  $p(\cdot), p_1(\cdot), p_2(\cdot) \in \mathcal{P}_0^*$  are such that  $1/p(\cdot) = 1/p_1(\cdot) + 1/p_2(\cdot)$ ,  $0 < q \leq \infty$  and  $s > n(1/\min(\tau_{p(\cdot)}, q, 1) - 1)$ , it holds that*

$$\|T_\sigma(f, g)\|_{F_{p(\cdot), q}^s} \lesssim \|f\|_{F_{p_1(\cdot), q}^{s+m}} \|g\|_{h^{p_2(\cdot)}} + \|f\|_{h^{p_1(\cdot)}} \|g\|_{F_{p_2(\cdot), q}^{s+m}} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n).$$

*Moreover, if  $p(\cdot) \in \mathcal{P}_0^*$ ,  $0 < q \leq \infty$  and  $s > n(1/\min(\tau_{p(\cdot)}, q, 1) - 1)$ , it holds that*

$$\|T_\sigma(f, g)\|_{F_{p(\cdot), q}^s} \lesssim \|f\|_{F_{p(\cdot), q}^{s+m}} \|g\|_{L^\infty} + \|f\|_{L^\infty} \|g\|_{F_{p(\cdot), q}^{s+m}} \quad \forall f, g \in \mathcal{S}(\mathbb{R}^n).$$

## 2.4 Applications to scattering properties of PDEs

# Bibliography

- [1] T. Kato and G. Ponce. Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.*, 41(7):891–907, 1988. ISSN 0010-3640. doi: 10.1002/cpa.3160410704. URL <http://dx.doi.org.er.lib.k-state.edu/10.1002/cpa.3160410704>.
- [2] A. Gulisashvili and M. Kon. Exact smoothing properties of Schrödinger semi-groups. *Amer. J. Math.*, 118(6):1215–1248, 1996. ISSN 0002-9327. URL [http://muse.jhu.edu.er.lib.k-state.edu/journals/american\\_journal\\_of\\_mathematics/v118/118.6gulisashvili.pdf](http://muse.jhu.edu.er.lib.k-state.edu/journals/american_journal_of_mathematics/v118/118.6gulisashvili.pdf).
- [3] L. Grafakos and S. Oh. The Kato-Ponce inequality. *Comm. Partial Differential Equations*, 39(6):1128–1157, 2014. ISSN 0360-5302. doi: 10.1080/03605302.2013.822885. URL <http://dx.doi.org.er.lib.k-state.edu/10.1080/03605302.2013.822885>.
- [4] C. Muscalu and W. Schlag. *Classical and multilinear harmonic analysis. Vol. II*, volume 138 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2013. ISBN 978-1-107-03182-1.
- [5] J. Bourgain and D. Li. On an endpoint Kato-Ponce inequality. *Differential Integral Equations*, 27(11-12):1037–1072, 2014. ISSN 0893-4983. URL <http://projecteuclid.org/euclid.die/1408366784>.
- [6] L. Grafakos, D. Maldonado, and V. Naibo. A remark on an endpoint Kato-Ponce inequality. *Differential Integral Equations*, 27(5-6):415–424, 2014. ISSN 0893-4983. URL <http://projecteuclid.org.er.lib.k-state.edu/euclid.die/1396558089>.

- [7] M. Christ and M. Weinstein. Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation. *J. Funct. Anal.*, 100(1):87–109, 1991. ISSN 0022-1236. doi: 10.1016/0022-1236(91)90103-C. URL [http://dx.doi.org.er.lib.k-state.edu/10.1016/0022-1236\(91\)90103-C](http://dx.doi.org.er.lib.k-state.edu/10.1016/0022-1236(91)90103-C).
- [8] C. Kenig, G. Ponce, and L. Vega. Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle. *Comm. Pure Appl. Math.*, 46(4):527–620, 1993. ISSN 0010-3640. doi: 10.1002/cpa.3160460405. URL <http://dx.doi.org.er.lib.k-state.edu/10.1002/cpa.3160460405>.
- [9] J. Brummer and V. Naibo. Bilinear operators with homogeneous symbols, smooth molecules, and Kato-Ponce inequalities. *Proc. Amer. Math. Soc.*, 146(3):1217–1230, 2018. ISSN 0002-9939. URL <https://doi-org.er.lib.k-state.edu/10.1090/proc/13841>.
- [10] M. Frazier and B. Jawerth. Decomposition of Besov spaces. *Indiana Univ. Math. J.*, 34(4):777–799, 1985. ISSN 0022-2518. doi: 10.1512/iumj.1985.34.34041. URL <http://dx.doi.org.er.lib.k-state.edu/10.1512/iumj.1985.34.34041>.
- [11] M. Frazier and B. Jawerth. A discrete transform and decompositions of distribution spaces. *J. Funct. Anal.*, 93(1):34–170, 1990. ISSN 0022-1236. doi: 10.1016/0022-1236(90)90137-A. URL [http://dx.doi.org.er.lib.k-state.edu/10.1016/0022-1236\(90\)90137-A](http://dx.doi.org.er.lib.k-state.edu/10.1016/0022-1236(90)90137-A).
- [12] J. Hart, R.H. Torres, and X. Wu. Smoothing properties of bilinear operators and Leibniz-type rules in Lebesgue and mixed Lebesgue spaces. *Trans. Amer. Math. Soc.*, DOI: <https://doi.org/10.1090/tran/7312>.
- [13] Á. Bényi and R. H. Torres. Symbolic calculus and the transposes of bilinear pseudodifferential operators. *Comm. Partial Differential Equations*, 28(5-6):1161–1181, 2003.

ISSN 0360-5302. doi: 10.1081/PDE-120021190. URL <http://dx.doi.org/10.1081/PDE-120021190>.

- [14] Á. Bényi, A. R. Nahmod, and R. H. Torres. Sobolev space estimates and symbolic calculus for bilinear pseudodifferential operators. *J. Geom. Anal.*, 16(3):431–453, 2006. ISSN 1050-6926. doi: 10.1007/BF02922061. URL <http://dx.doi.org.er.lib.k-state.edu/10.1007/BF02922061>.
- [15] Á. Bényi. Bilinear pseudodifferential operators with forbidden symbols on Lipschitz and Besov spaces. *J. Math. Anal. Appl.*, 284(1):97–103, 2003. ISSN 0022-247X. doi: 10.1016/S0022-247X(03)00245-2. URL [http://dx.doi.org.er.lib.k-state.edu/10.1016/S0022-247X\(03\)00245-2](http://dx.doi.org.er.lib.k-state.edu/10.1016/S0022-247X(03)00245-2).
- [16] V. Naibo. On the bilinear Hörmander classes in the scales of Triebel-Lizorkin and Besov spaces. *J. Fourier Anal. Appl.*, 21(5):1077–1104, 2015. ISSN 1069-5869. doi: 10.1007/s00041-015-9398-x. URL <http://dx.doi.org.er.lib.k-state.edu/10.1007/s00041-015-9398-x>.
- [17] K. Koezuka and N. Tomita. Bilinear Pseudo-differential Operators with Symbols in  $BS_{1,1}^m$  on Triebel-Lizorkin Spaces. *J. Fourier Anal. Appl.*, 24(1):309–319, 2018. ISSN 1069-5869. URL <https://doi-org.er.lib.k-state.edu/10.1007/s00041-016-9518-2>.
- [18] D. Cruz-Uribe and V. Naibo. Kato-Ponce inequalities on weighted and variable Lebesgue spaces. *Differential Integral Equations*, 29(9-10):801–836, 2016. ISSN 0893-4983. URL <http://projecteuclid.org/euclid.die/1465912605>.
- [19] J. Brummer and V. Naibo. Weighted fractional Leibniz-type rules for bilinear multiplier operators. *Preprint*, 2017.
- [20] R. Coifman and Y. Meyer. *Au delà des opérateurs pseudo-différentiels*, volume 57 of *Astérisque*. Société Mathématique de France, Paris, 1978.

- [21] Loukas Grafakos and Rodolfo H. Torres. Multilinear Calderón-Zygmund theory. *Adv. Math.*, 165(1):124–164, 2002. ISSN 0001-8708. doi: 10.1006/aima.2001.2028. URL <https://doi-org.er.lib.k-state.edu/10.1006/aima.2001.2028>.
- [22] Loukas Grafakos and Rodolfo H. Torres. Maximal operator and weighted norm inequalities for multilinear singular integrals. *Indiana Univ. Math. J.*, 51(5):1261–1276, 2002. ISSN 0022-2518. doi: 10.1512/iumj.2002.51.2114. URL <https://doi-org.er.lib.k-state.edu/10.1512/iumj.2002.51.2114>.
- [23] Andrei K. Lerner, Sheldy Ombrosi, Carlos Pérez, Rodolfo H. Torres, and Rodrigo Trujillo-González. New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory. *Adv. Math.*, 220(4):1222–1264, 2009. ISSN 0001-8708. doi: 10.1016/j.aim.2008.10.014. URL <https://doi-org.er.lib.k-state.edu/10.1016/j.aim.2008.10.014>.
- [24] L. Grafakos. *Modern Fourier analysis*, volume 250 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014. ISBN 978-1-4939-1229-2; 978-1-4939-1230-8. doi: 10.1007/978-1-4939-1230-8. URL <http://dx.doi.org.er.lib.k-state.edu/10.1007/978-1-4939-1230-8>.
- [25] H. Triebel. *Theory of function spaces*. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 2010. ISBN 978-3-0346-0415-4; 978-3-0346-0416-1. Reprint of 1983 edition. Also published in 1983 by Birkhäuser Verlag.
- [26] M. Yamazaki. A quasihomogeneous version of paradifferential operators. I. Boundedness on spaces of Besov type. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 33(1):131–174, 1986. ISSN 0040-8980.
- [27] R. Hunt. On  $L(p, q)$  spaces. *Enseignement Math. (2)*, 12:249–276, 1966. ISSN 0013-8584.

- [28] D. Cruz-Uribe, J. M. Martell, and C. Pérez. *Weights, extrapolation and the theory of Rubio de Francia*, volume 215 of *Operator Theory: Advances and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011. ISBN 978-3-0348-0071-6. doi: 10.1007/978-3-0348-0072-3. URL <http://dx.doi.org/10.1007/978-3-0348-0072-3>.
- [29] M. Rosenthal and H.-J. Schmeisser. The boundedness of operators in Muckenhoupt weighted Morrey spaces via extrapolation techniques and duality. *Rev. Mat. Complut.*, 29(3):623–657, 2016. ISSN 1139-1138. doi: 10.1007/s13163-016-0208-z. URL <https://doi-org.er.lib.k-state.edu/10.1007/s13163-016-0208-z>.
- [30] H. Kozono and M. Yamazaki. Semilinear heat equations and the Navier-Stokes equation with distributions in new function spaces as initial data. *Comm. Partial Differential Equations*, 19(5-6):959–1014, 1994. ISSN 0360-5302. doi: 10.1080/03605309408821042. URL <https://doi-org.er.lib.k-state.edu/10.1080/03605309408821042>.
- [31] A. L. Mazzucato. Besov-Morrey spaces: function space theory and applications to non-linear PDE. *Trans. Amer. Math. Soc.*, 355(4):1297–1364, 2003. ISSN 0002-9947. doi: 10.1090/S0002-9947-02-03214-2. URL <https://doi-org.er.lib.k-state.edu/10.1090/S0002-9947-02-03214-2>.
- [32] M. Izuki, Y. Sawano, and H. Tanaka. Weighted Besov-Morrey spaces and Triebel-Lizorkin spaces. In *Harmonic analysis and nonlinear partial differential equations*, RIMS Kôkyûroku Bessatsu, B22, pages 21–60. Res. Inst. Math. Sci. (RIMS), Kyoto, 2010.
- [33] J. Xu. Variable Besov and Triebel-Lizorkin spaces. *Ann. Acad. Sci. Fenn. Math.*, 33(2):511–522, 2008. ISSN 1239-629X.
- [34] P. Gurka, P. Harjulehto, and A. Nekvinda. Bessel potential spaces with variable exponent. *Math. Inequal. Appl.*, 10(3):661–676, 2007. ISSN 1331-4343. doi: 10.7153/mia-10-61. URL <http://dx.doi.org.er.lib.k-state.edu/10.7153/mia-10-61>.

- [35] J. Xu. The relation between variable Bessel potential spaces and Triebel-Lizorkin spaces. *Integral Transforms Spec. Funct.*, 19(7-8):599–605, 2008. ISSN 1065-2469. doi: 10.1080/10652460802030631. URL <http://dx.doi.org.er.lib.k-state.edu/10.1080/10652460802030631>.
- [36] L. Diening, P. Hästö, and S. Roudenko. Function spaces of variable smoothness and integrability. *J. Funct. Anal.*, 256(6):1731–1768, 2009. ISSN 0022-1236. doi: 10.1016/j.jfa.2009.01.017. URL <http://dx.doi.org.er.lib.k-state.edu/10.1016/j.jfa.2009.01.017>.
- [37] A. Almeida and P. Hästö. Besov spaces with variable smoothness and integrability. *J. Funct. Anal.*, 258(5):1628–1655, 2010. ISSN 0022-1236. doi: 10.1016/j.jfa.2009.09.012. URL <http://dx.doi.org.er.lib.k-state.edu/10.1016/j.jfa.2009.09.012>.
- [38] D. Cruz-Uribe and A. Fiorenza. *Variable Lebesgue spaces*. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, Heidelberg, 2013. ISBN 978-3-0348-0547-6; 978-3-0348-0548-3. doi: 10.1007/978-3-0348-0548-3. URL <http://dx.doi.org.er.lib.k-state.edu/10.1007/978-3-0348-0548-3>. Foundations and harmonic analysis.
- [39] Á. Bényi, D. Maldonado, V. Naibo, and R. H. Torres. On the Hörmander classes of bilinear pseudodifferential operators. *Integral Equations Operator Theory*, 67(3):341–364, 2010. ISSN 0378-620X. doi: 10.1007/s00020-010-1782-y. URL <http://dx.doi.org/10.1007/s00020-010-1782-y>.
- [40] F. Bernicot and P. Germain. Bilinear oscillatory integrals and boundedness for new bilinear multipliers. *Adv. Math.*, 225(4):1739–1785, 2010. ISSN 0001-8708. doi: 10.1016/j.aim.2010.03.032. URL <http://dx.doi.org.er.lib.k-state.edu/10.1016/j.aim.2010.03.032>.

# Appendix A

## Title for This Appendix

Enter the content for Appendix A in the appendixA.tex file. If you do not have an Appendix A, see comments in the etdrtemplate.tex file for instructions on how to remove this page.