
#### relativistische Betrachtung von Bewegungen


Alexander Helbok

19. Januar 2024

## Radiobeobachtungen




(Matthews, Morgan und Schmidt 1964)



(Matthews, Morgan und Schmidt 1964)

#### Leuchtkraft von 3C 273



Entfernung über Rotverschiebung


$$(z = 0.158)$$


$$L = 4\pi d^2 F$$


Flussmessung im Radiobereich


 $\Rightarrow$  extrem Hell


## Helligkeitsschwankungen in 3C 273














































## Helligkeitsschwankungen in 3C 273








## sphärische Expansion $v \ll c$



## sphärische Expansion $v \sim c$








"Because the observed intensity of a source, for a given surface brightness, is proportional to the apparent size" [4]


$$F = B < \propto 2 \frac{\gamma vt}{R}$$



"Because the observed intensity of a source, for a given surface brightness, is proportional to the apparent size" [4]


$$F = B < \propto 2 \frac{\gamma vt}{R} \quad \frac{\mathrm{d}F}{\mathrm{d}t} = 2 \frac{\gamma v}{R} \propto \gamma$$

"it is already clear that an expanding source could exhibit a rate of increase of flux density high enough to explain the observations." [4]




"If, however, a source at the distance of 3C 273 were to start to explode with a velocity corresponding to  $\gamma=5$ , and if H (measured in a frame sharing the mean particle motion)  $\sim 10^{-2}$  gauss, the flux density would have risen to  $\sim 15$  F.U. in 3 years" [4]

#### Konsequenz





$$\begin{split} \beta &= \frac{v}{c} \in [0;1] \\ \gamma &= \frac{1}{\sqrt{1-\beta^2}} \in [1,\infty) \quad \Rightarrow \quad \gamma v > c \text{ m\"{o}glich!} \end{split}$$


#### **Kosmische Jets**





3C 273 © ESA/HUBBLE & NASA














Fragen?

$$\beta = 0.835, \ \gamma = 1.82$$





$$\beta = 0.835, \ \gamma = 1.82$$


$$(t = 0.25)$$






$$\beta = 0.835, \ \gamma = 1.82$$

$$t = 0.59$$








$$\beta = 0.835, \ \gamma = 1.82$$






$$\beta = 0.835, \ \gamma = 1.82$$

$$t = 2.5$$


to O





$$\beta = 0.835, \ \gamma = 1.82$$

$$t = 4$$

to O







#### Sources i

#### Literatur

- [1] T. A. Matthews, W. W. Morgan und M. Schmidt. "A Discussion of Galaxies Indentified with Radio Sources.". In: Astrophysical Journal, vol. 140, p. 35 140 (1964), S. 35.
- [2] P. Maltby und A. T. Moffett. "Spectrum of the Intensity Variations in 3C 273B". In: Science 150.3692 (1965), S. 63–64. ISSN: 00368075, 10959203. URL: http://www.jstor.org/stable/1717963 (besucht am 28.12.2023).

#### Sources ii

- [3] S. Fernandes u. a. "Multiwavelength analysis of the variability of the blazar 3C 273". In: Monthly Notices of the Royal Astronomical Society 497.2 (Juli 2020), S. 2066–2077. ISSN: 1365-2966. URL: http://dx.doi.org/10.1093/mnras/staa2013.
- [4] M. Rees. "Appearance of relativistically expanding radio sources". In: *Nature* 211.5048 (1966), S. 468–470.