177. Stabpendel mit Kugeln

$$L=0.3 \text{ m}; \qquad r_1=0.02 \text{ m}; \qquad r_2=0.03 \text{ m} \ m_0=0.1 \text{ kg}; \qquad m_1=0.1 \text{ kg}; \qquad m_2=0.3 \text{ kg}$$

a)
$$\begin{split} I_{ges} &= I_{Stab} + I_{Kugel_1} + I_{Kugel_2} \\ &I_{Stab} = \tfrac{1}{12} m_0 L^2 = 7.5 * 10^{-4} \text{ kg m}^2 \\ &I_{Kugel_1} = \tfrac{2}{5} m_1 r_1^2 + m_1 (\tfrac{L}{2} + r_1)^2 = 2.9 * 10^{-3} \text{ kg m}^2 \\ &I_{Kugel_2} = \tfrac{2}{5} m_2 r_2^2 + m_2 (\tfrac{L}{2} + r_2)^2 = 9.8 * 10^{-3} \text{ kg m}^2 \end{split}$$

$$\Rightarrow I_{ges} = \underbrace{1.3 * 10^{-2} \text{ kg m}^2}$$

b)

$$\begin{split} x_{Stab_{sp}} &= 0 \text{ m} \\ x_{Kugeln_{sp}} &= \frac{m_1 \left(-\frac{2r_1 + L}{2} \right) + m_2 \left(\frac{2r_2 + L}{2} \right)}{m_1 + m_2} = 0.0925 \text{ m} \\ x_{sp} &= \frac{x_{Stab_{sp}} * m_0 + x_{Kugeln_{sp}} * (m_1 + m_2)}{m_0 + m_1 + m_2} = \underline{0.074} \text{ m} \end{split}$$

c)
$$T = 2\pi \sqrt{\frac{I}{Mgl}}$$
$$T = 2\pi \sqrt{\frac{I}{(m_0 + m_1 + m_2) g x_{sp}}}$$
$$= \underline{1.2 \text{ s}}$$

184. Resonanzverhalten einer Stimmgabel

$$f_0 = 440 \text{ Hz}; \quad \omega_0 = 2\pi f_0; \quad x(\tau) = x_m e^{-\beta \tau/2}; \quad x(5) = \frac{1}{10}x(0)$$

$$\frac{x_m}{10} = x_m e^{-\beta 5/2}$$

$$\beta = -\frac{2}{5} \ln(0.1)$$

$$\tau = \frac{1}{\beta} = \underline{1.09 \text{ s}}$$

b)
$$\omega' = \sqrt{\omega_0^2 - \frac{\beta^2}{4}}$$

$$\omega' = 2764.6 \text{ Hz} \qquad (\omega_0 \approx \omega')$$

$$f' = \frac{\omega'}{2\pi} = 440 \text{ Hz} \qquad (f_0 \approx f')$$

$$f' * \tau = \underline{477.7}$$

c)
$$Q = \frac{\omega_0}{\beta}$$

$$Q = \underline{3002}$$

d)
$$\delta f = \frac{f'}{Q}$$

$$\delta f = 0.15 \text{ Hz}$$

e)
$$x_m(f) = \frac{2\pi f_0}{\sqrt{16\pi^2 (f - f_0)^2 + \beta^2}};$$
 $E(f) = \frac{(\pi f_0)^2}{4\pi^2 (f - f_0)^2 + (\beta/2)^2};$ $f_1 = 439.5 \text{ Hz}$

$$x_m(f_0) = \frac{2\pi f_0}{\sqrt{16\pi^2 (f_0 - f_0)^2 + \beta^2}} = 3001.63$$

$$x_m(f_1) = \frac{2\pi f_0}{\sqrt{16\pi^2 (f_1 - f_0)^2 + \beta^2}} = 435.348$$

$$x_{rel} = \frac{x_m(f_1)}{x_m(f_0)} = \underline{0.15}$$

$$E(f_0) = \frac{(\pi f_0)^2}{4\pi^2 (f_0 - f_0)^2 + (\beta/2)^2} = 9.0 * 10^6$$

$$E(f_1) = \frac{(\pi f_0)^2}{4\pi^2 (f_1 - f_0)^2 + (\beta/2)^2} = 1.9 * 10^5$$

$$E_{rel} = \frac{E(f_1)}{E(f_0)} = \underline{0.021}$$

185. Resonante Anregung eines Federpendels

$$m = 0.1 \text{ kg}; \quad k = 40 \text{ N/m}$$

a)
$$\omega_0 = \sqrt{\frac{k}{m}} = 20 \text{ rad/s}$$

$$\frac{x_m}{2} = x_m * e^{-\beta t/2}$$

$$\beta = -\frac{\ln(0.5)2}{t} = \underline{0.14 \text{ s}^{-1}}$$

$$\omega' = \sqrt{\omega_0^2 - \frac{\beta^2}{4}} = \underline{19.99988 \text{ rad/s}}$$

$$Q = \frac{\omega_0}{\beta} = \underline{144.3}$$

c)
$$\hat{y}_{res} = \frac{\omega_0}{\beta} y_0$$
; $\hat{y}_{res} = 0.012 \text{ m}$
 $y_0 = \hat{y}_{res} \frac{\beta}{\omega_0} = \underline{8.32 * 10^{-5} \text{ m}}$

d)
$$y_2 = y_0 \frac{\omega_0}{\sqrt{4(\omega_1 - \omega_0)^2 + \beta^2}}; \quad \omega_1 = \omega_0 \pm 0.2\pi$$

 $y_2 = \underline{0.0013 \text{ m}}$