





VMM2 with MMFE8: TDO distributions

NSW Electronics Meeting
April 1, 2016

M. Bledsoe, N. Felt, M. Franklin, B. Garber, P. Giromini, J.
Grotto, J. Huth, C. Rogan, A. Tuna, A. Wang

Overview




Working to put together a functioning system

Timing studies

- Our goal was to examine timing information for data taken with the VMM2 on the MMFE8 board
- Studied TDO distributions with varying peak times and DAC pulse heights
- Compared to data from VMM1 studies


VMM2 timing study methods

- In order to do the TDO calibration, we have synchronized CKTP to CKBC
- Then we “delay” it using ticks of the faster mother clock (200 MHz) : 5 delays
- Rising edge of CKTP can be held fixed with respect to falling edge of CKBC



VMM7, Ch 14, TDO vs. Test Pulse DAC

For every TP DAC, pulsed 100 times




- CKTP pulses with every fixed test pulse DAC TDO distribution are fully synchronized with respect to CKBC


VMM 7, Ch 14

TDO distributions

Test Pulse DAC = 120



Test Pulse DAC = 140




- CKTP in each plot is fully synchronized with respect to CKBC


VMM 7, Ch 14

TDO distributions

Test Pulse DAC = 160




Test Pulse DAC = 180



- Large decrease in RMS

Studying the TDO as a function of the peak time




Smaller the peak time -> smaller TDO RMS

Investigating origin of the dependence


- Since TDO is the difference between the next falling edge of CKBC and the peak time, we don't know if the large RMS is because of the bunch crossing or the peak detection
- In order to see if the large dependence of TDO on PDO / peaktime was caused by some phenomenon unique to each channel, we pulsed two channels simultaneously and examined the differences in TDO for each data point

Difference in TDO values for two simultaneously pulsed channels



- The RMS of the difference is larger than the RMS of just one channel -> problem is with peak detection
- We are surprised that the RMS is so large

Going back in time to VMM1



ATL-UPGRADE-PUB-2014-001

Between 0.3 to 0.9 V,
timing resolution < 3 ns
for a 200 ns peak time

Going back in time to VMM1: logbook edition


June 2013

Measure how TDO changes with PDAC

PDAC ← 250 300 350 400

E2E	250	300	350	400
0	TDO = 262.8 RMS = 6.50	T: 276.6 R: 7.39	T: 277.2 R: 1.75	T: 274.4 R: 1.57
10	TDO = 414.3 RMS = 6.77	T: 348.4 R: 7.91	T: 427.8 R: 2.33	T: 422.93 R: 2.11
20	TDO: 572.4 RMS: 7.29	T: 585.4 R: 8.33	T: 583.5 R: 2.95	T: 577.53 R: 2.80
30	T: 733.1 R: 7.58	T: 746.0 R: 8.53	T: 743.0 R: 3.58	T: 736.2 R: 3.43
40	T: 892.8 R: 7.72	T: 903.8 R: 8.64	T: 901.0 R: 4.26	T: 894.1 R: 4.11

1 count ~ 1 ns



Dependence of test pulse amplitude on peak time


- After we noticed that the RMS was improved with shorter peak time and larger test pulse DAC, we wanted to see if a change in peak time would change the voltage of the test pulse amplitude
- Also collected approximate values for test pulse amplitudes on VMM7, Ch 14

Changing peak time changes shaped pulse amplitude

TP DAC	200 ns Peak Time	25 ns Peak Time
200	~920 mV	~650 mV
120	~330 mV	~230 mV
80	~290 mV	~220 mV

- Pulse height is affected by the peak time
- Shorter peak time ~ shorter pulse amplitude

Fall time of DAC step pulse




- The fall/rise time of the DAC step pulse is long, $> 100\text{ ns}$
- Accounts for the dependence of test pulse pulse amplitude on peak time
- For TP DAC = 0, we still have a pulse amplitude of $\sim 100\text{-}200\text{ mV}$
- Not probing low enough voltages

Summary


- TDO distributions at lower test pulse DACs / long peak time are very wide and span the full bunch cross
- At even lower voltages, using just the BCID may be better than using the time of flight (with just the BCID, $25 \text{ ns} / \sqrt{12} \sim 7.2 \text{ ns RMS}$)
- May signal a problem in the peak detection of the signal
- Test pulse amplitude dependent on peak time of test pulse, most likely because DAC step pulse has a long rise time

Backup

VMM 7, Ch 14, overlaid distributions





VMM 7, Ch 14, overlaid distributions



7-8 bins ~ 5 ns

Distributions have become much narrower

VMM 7, Ch 14, TDO distributions




- All of data is from one synchronization point with respect to CKBC

VMM 7, Ch 14



TDO distributions

Test Pulse DAC = 200





With some reduced noise
(disconnected logic
probes)



VMM 7, Ch 14, TDO distributions




VMM 7, Ch 14, TDO distributions




VMM 7, Ch 14, TDO distributions



VMM 7, Ch 14, TDO distributions



Rough calibration plot



Talk by C. Rogan: <https://indico.cern.ch/event/512405/>