
Contents

62	Acknowledgements	iii
63	Abstract	iv
64	Contents	v
65	Preface	ix
66	1 Introduction	1
67	2 The Standard Model	2
68	3 The LHC and the ATLAS detector	3
69	3.1 The LHC	3
70	3.2 The ATLAS detector	3
71	3.2.1 Tracking	3
72	3.2.2 Calorimetry	3
73	3.2.3 Muon spectrometry	3
74	3.3 Particle identification	3
75	3.3.1 Muons	3
76	3.3.2 Electrons and photons	3
77	3.3.3 Hadrons	3
78	3.3.4 Neutrinos	3
79	3.4 Triggering	3
80	3.4.1 L1	3
81	3.4.2 HLT	3

82	4 Tau leptons	11
83	4.1 Tau leptons	11
84	4.2 Leptonic tau decays, τ_ℓ	11
85	4.3 Hadronic tau decays, τ_{had}	13
86	4.3.1 Reconstruction	13
87	4.3.1.1 Calorimeter seeding	13
88	4.3.1.2 Track and vertex association	13
89	4.3.1.3 Calibration	14
90	4.4 Identification	15
91	4.5 Leptons mis-identified as τ_{had}	20
92	4.5.1 Electrons	20
93	4.5.2 Muons	20
94	5 $H \rightarrow \tau_\ell \tau_{\text{had}}$ strategy	23
95	5.1 Introduction	23
96	5.1.1 ATLAS Higgs program	23
97	5.1.2 $H \rightarrow \tau\tau$	23
98	5.2 Physics objects	23
99	5.2.1 Electrons, muons, and τ_{had}	23
100	5.2.2 Jets and $E_{\text{T}}^{\text{miss}}$	23
101	5.3 Categorization	23
102	5.3.1 Pre-selection	23
103	5.3.2 VBF category	29
104	5.3.3 Boosted category	29
105	5.4 $\tau\tau$ mass reconstruction	30
106	5.4.1 $\tau\tau$ systems	30
107	5.4.2 $m_{\tau\tau}^{\text{MMC}}$ algorithm	30
108	5.4.3 Performance	30
109	5.5 MVA discrimination	30
110	5.5.1 Inputs	30
111	5.5.2 Correlations	32
112	5.5.3 MVAs in other VBF analyses	40
113	6 Signal and background predictions	42

114	6.1	$Z \rightarrow \tau\tau$	42
115	6.1.1	Mis-modeling of $Z(\rightarrow \ell\ell)+\text{jets}$ in simulation	42
116	6.1.2	Embedding	42
117	6.1.3	Validation	42
118	6.1.4	Uncertainties	45
119	6.2	$j \rightarrow \tau_{\text{had}}$ mis-identification	46
120	6.2.1	Mis-modeling of $j \rightarrow \tau_{\text{had}}$ in simulation	46
121	6.2.2	Fake factor method	46
122	6.2.3	Composition of $j \rightarrow \tau_{\text{had}}$ in the SR	50
123	6.2.4	Validation	55
124	6.2.5	Uncertainties	60
125	6.3	top, $Z \rightarrow \ell\ell$, diboson	61
126	6.4	$H \rightarrow \tau\tau$	61
127	6.4.1	Generators	61
128	6.4.2	Uncertainties	61
129	7	Results	63
130	7.1	Predictions in the signal region	63
131	7.2	$H \rightarrow \tau_{\text{had}}\tau_{\text{had}}$ and $H \rightarrow \tau_{\ell}\tau_{\ell}$	63
132	7.3	Fit results	63
133	8	Prospects for $H \rightarrow \tau\tau$	69
134	8.1	Run-II	69
135	8.1.1	Run-I triggers for $H \rightarrow \tau\tau$	70
136	8.1.2	Run-II triggers	72
137	8.1.2.1	Object thresholds	72
138	8.1.2.2	Topological requirements	73
139	8.1.2.3	Gains with $\ell+\tau_{\text{had}}$ triggers	76
140	8.1.3	Updates to the L1 τ_{had} object	79
141	8.1.3.1	Size	79
142	8.1.3.2	Isolation	81
143	8.1.4	Conclusions and contingencies	81
144	8.2	HL-LHC	84
145	8.2.1	Selection	84

146	8.2.2	Emulation of High-Luminosity LHC conditions	86
147	8.2.2.1	Performance assumptions	86
148	8.2.2.2	Pileup emulation	86
149	8.2.2.3	Impact on observables	87
150	8.2.3	Analysis	87
151	8.2.3.1	Boosted decision tree training	87
152	8.2.3.2	Kinematic distributions	89
153	8.2.4	Results	89
154	8.2.4.1	Uncertainties assumptions	89
155	8.2.5	Conclusions	93
156	A	Control regions for fakes	95
157	A.1	$W \rightarrow \ell \nu_\ell$ CR	95
158	A.2	QCD CR	98
159	A.3	$Z \rightarrow \ell \ell$ CR	100
160	A.4	top CR	102
161	B	Inputs to the τ_{had} BDT identifier	104
162	C	Performance of $m_{\tau\tau}$ algorithms	107
163		Bibliography	110