
List of Figures

192	3.1	Aerial view of Geneva with an overlaid drawing of the LHC and associated experiments .	4
193	3.2	The LHC accelerator complex. Before reaching the LHC, protons are accelerated at Linac	
194		2, the Proton Synchrotron Booster (PSB), the Proton Synchrotron (PS), and the Super	
195		Proton Synchrotron (SPS)	5
196	3.3	Cross sections for pp and $p\bar{p}$ processes in the center-of-mass energy regime relevant to the	
197		Tevatron and LHC, courtesy of W.J. Stirling	6
198	3.4	Scale rendering of the ATLAS detector with the various sub-detectors highlighted	7
199	3.5	Transverse schematic view of a wedge of the ATLAS detector. Charged particles leave	
200		tracks in the tracker, electrons and photons typically stop in the electromagnetic calorime-	
201		ter, hadrons like charged pions typically stop in the hadronic calorimeter, and muons are	
202		tagged by the muon system as they exit. Neutrinos escape undetected.	8
203	3.6	The peak ATLAS online luminosity as measured in different data-taking periods . The	
204		peak Run-I luminosity is $0.8 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$	8
205	3.7	Distributions of the recorded luminosity in bins of $\langle\mu\rangle$ (left) and the total integrated	
206		luminosity as a function of time (right) . In 2011 (2012), the average $\langle\mu\rangle$ is 9.1 (20.7) and	
207		the total integrated luminosity for physics analysis is 4.6 fb^{-1} . (20.3 fb^{-1}).	9
208	3.8	Summary of cross sections measured at ATLAS in 7 and 8 TeV data-taking	9
209	3.9	Schematic view of the calorimeter granularity available at the L1 trigger	10
210	4.1	Pie chart of tau lepton decay branching fractions, grouped by hadronic decays (65%) and	
211		leptonic decays (35%).	12
212	4.2	True p_T and reconstructed d_0 for muons from simulated W , Z , and tau lepton decays.	
213		Muons from tau lepton decays are shown for $Z \rightarrow \tau\tau$, $H^{\text{ggF}} \rightarrow \tau\tau$, and $H^{\text{VBF}} \rightarrow \tau\tau$	
214		processes.	12
215	4.3	Track selection efficiency for τ_{had} candidates with the default vertex selection (highest	
216		Σp_T^2) versus the dedicated TJVA algorithm, for true 1-track (left) and 3-track (right) τ_{had} ,	
217		as a function of $\langle\mu\rangle$	14
218	4.4	τ_{had} energy response curves measured in simulation, for 1-track (left) and 2,3-track (right)	
219		τ_{had} , as a function of the reconstructed energy	15
220	4.5	τ_{had} energy resolution measured, for 1-track (left) and 2,3-track (right) τ_{had} , as a function	
221		of the true visible energy	15

222	4.6	Event display of a $tt \rightarrow (b\mu\nu_\mu)(b\tau_{\text{had}}\nu_\tau)$ candidate during 2011 data-taking . The τ_{had} candidate has 3 tracks, the b -jet candidates each have more than 10 tracks, and the muon is in red. The estimated purity of the selection is greater than 75%.	17
223			
224			
225	4.7	Fit of the predicted τ_{had} track multiplicity to data in a $Z \rightarrow \tau_\mu \tau_{\text{had}}$ event selection before applying tau identification algorithms . The τ_{had} candidates have much lower track multiplicity relative to the large jet background.	18
226			
227			
228	4.8	Signal and background distributions for two of the discriminating variables in the 1-track τ_{had} jet discrimination algorithm: f_{cent} (left) and $N_{\text{track}}^{\text{iso}}$ (right) . The remaining distributions are shown in Appendix B.	19
229			
230			
231	4.9	Signal and background distributions for two of the discriminating variables in the 3-track τ_{had} jet discrimination algorithm: R_{track} (left) and $m_{\pi^0+\text{track}}$ (right) . The remaining distributions are shown in Appendix B.	20
232			
233			
234	4.10	Signal efficiency versus inverse background efficiency for 1-track and 3-track τ_{had} jet discrimination algorithms in a lower- p_T regime (left) and higher- p_T regime (right) . The loose, medium, and tight operating points are highlighted with red markers.	20
235			
236			
237	4.11	Signal and background distribution for one of the discriminating variables in the τ_{had} electron discrimination algorithm: the TRT high threshold fraction (left), and signal efficiency versus inverse background efficiency for the algorithm (right)	20
238			
239			
240	4.12	The visible mass $m_{e\tau_{\text{had}}}$ in a $Z \rightarrow ee$ selection in data after requiring the τ_{had} candidate pass the medium jet discrimination algorithm and not overlap spatially with a tight identified electron (left) and after additionally requiring the τ_{had} pass the loose τ_{had} electron discrimination algorithm (right).	21
241			
242			
243			
244	4.13	True $m_{\mu\mu}$ (left) and $\eta(\mu)$ (right) in $Z \rightarrow \mu\mu$ events where a muon is mis-identified as a τ_{had} . The muons are split into combined muons (black), muons which pass tracking requirements but fail combined requirements (green), are reconstructed but fail tracking requirements (blue), and are not reconstructed (red). A large fraction of non-reconstructed muons have $ \eta \approx 0$, which is a poorly covered region of the muon system.	21
245			
246			
247			
248			
249	4.14	True $m_{\mu\mu\gamma}$ (left) and $\Delta R(\mu, \gamma)$ (right) in $Z \rightarrow \mu\mu$ events where a muon is mis-identified as a τ_{had} and a true ISR photon is associated to the muon. The muons are split into combined muons (black), muons which pass tracking requirements but fail combined requirements (green), are reconstructed but fail tracking requirements (blue), and are not reconstructed (red).	22
250			
251			
252			
253			
254	5.1	Pie chart of di-tau lepton decay branching fractions.	24
255	5.2	Measurement of the efficiency of the muon reconstructions algorithms in data and in simulation	24
256			
257	5.3	Validation of the muon energy scale corrections in J/Ψ events (left), Υ events (center), and Z events (right)	25
258			
259	5.4	Data and predictions of m_{ee} before the electron identification algorithm is applied (top, left) and after (top, right), and the measured efficiency of the algorithm as a function of E_T (bottom, left) and η (bottom, right)	25
260			
261			
262	5.5	Resolution of various E_T^{miss} reconstruction algorithms as a function of the number of reconstructed primary vertices in $Z \rightarrow \mu\mu$ events in data (left) and $W \rightarrow e\nu$ events in simulation (right)	26
263			
264			
265	5.6	Bias of various E_T^{miss} reconstruction algorithms as a function of p_T^Z measured in data events inclusively (left) and with no additional jets (right)	26
266			
267	5.7	Efficiency of b -jet identification algorithms measured in simulation as a function of light jet rejection (left) and b -jet p_T (right)	27
268			
269	5.8	Cartoon depiction of the relevant categories in the $H \rightarrow \tau_\ell \tau_{\text{had}}$ analysis: pre-selection, boosted, and VBF.	28
270			

271	5.9	Kinematic distributions in the pre-selection category of the 8 TeV $H \rightarrow \tau_\ell \tau_{\text{had}}$ analysis. .	29
272	5.10	Kinematic distributions in the pre-selection category of the 8 TeV $H \rightarrow \tau_\ell \tau_{\text{had}}$ analysis. .	30
273	5.11	Cartoon of the $m_{\tau\tau}^{\text{MMC}}$ reconstruction algorithm. Black, filled lines indicate items measured directly (ℓ , τ_{had}). Red, dotted lines indicate items which cannot be measured (neutrinos). The black, dashed line indicates the E_T^{miss} , which is measured indirectly. Blue indicates items which the $m_{\tau\tau}^{\text{MMC}}$ scans to find an optimal solution ($\Delta\phi$, E_T^{miss}).	31
274			
275	5.12	Input assumptions of the three-dimensional angle between the visible and invisible tau lepton decay products, for leptonic decays (left), 1-track hadronic decays (center), and 3-track hadronic decays (right)	31
276			
277	5.13	Efficiency for $H \rightarrow \tau_\ell \tau_{\text{had}}$ versus the efficiency for $Z \rightarrow \tau_\ell \tau_{\text{had}}$ for various $m_{\tau\tau}$ reconstruction algorithms in the boosted category (left) and VBF category (right).	32
278			
279	5.14	Cartoons of lepton η -centrality (left) and E_T^{miss} ϕ -centrality (right), courtesy of Tae Min Hong.	33
280			
281	5.15	Predicted signal and background distributions in the boosted category normalized to unit area and overlaid.	35
282			
283	5.16	Predicted signal and background distributions in the boosted category normalized to unit area and overlaid.	36
284			
285	5.17	Predicted signal and background distributions in the VBF category normalized to unit area and overlaid.	37
286			
287	5.18	Predicted signal and background distributions in the VBF category normalized to unit area and overlaid.	38
288			
289	5.19	Predicted signal and background distributions in the VBF category normalized to unit area and overlaid.	39
290			
291	5.20	Contours of kinematic correlations in the VBF category for VBF $H \rightarrow \tau\tau$ (left), $Z \rightarrow \tau\tau$ (center), and fakes (right).	40
292			
293	5.21	Contours of kinematic correlations in the VBF category for VBF $H \rightarrow \tau\tau$ (left), $Z \rightarrow \tau\tau$ (center), and fakes (right).	41
294			
295	5.22	Contours of kinematic correlations in the VBF category for VBF $H \rightarrow \tau\tau$ (left), $Z \rightarrow \tau\tau$ (center), and fakes (right).	42
296			
297	5.23	Contours of kinematic correlations in the VBF category for VBF $H \rightarrow \tau\tau$ (left), $Z \rightarrow \tau\tau$ (center), and fakes (right).	43
298			
299	5.24	Overlaid shapes of BDT outputs for signal and background processes in the VBF $H \rightarrow \gamma\gamma$, VBF $H \rightarrow ZZ^*$, and VBF $H \rightarrow WW^*$ analyses.	44
300			
301	6.1	Comparison of data and various predictions in $Z \rightarrow \ell\ell$ events of the charged particle scalar momentum density (left) and multiplicity density (right) as a function of p_T^Z in 2011 data-taking . Mis-modeling is observed for all predictions.	46
302			
303	6.2	Comparison of data and various predictions in $Z \rightarrow \ell\ell$ events of $\Delta Y(jj)$ (left) and m_{jj} (right) in 2011 data-taking . Mis-modeling is observed for all predictions.	46
304			
305	6.3	Comparison of data and various predictions of p_T^Z for $Z \rightarrow ee$ (left) and $Z \rightarrow \mu\mu$ (right) in 2011 data-taking . Mis-modeling is observed for all predictions.	47
306			
307	6.4	Validation of the embedding technique for simulated tau lepton decays in simulated $Z \rightarrow \mu\mu$ events (left) and simulated muons in data $Z \rightarrow \mu\mu$ events (right) . Good agreement is observed in both.	47
308			
309	6.5	The fractional uncertainty on the embedded $Z \rightarrow \tau_\ell \tau_{\text{had}}$ prediction in each bin of the VBF category for uncertainties pertaining to the embedding procedure and τ_{had} performance. .	48
310			
311	6.6	Data events in the VBF category which fail τ_{had} identification but fulfill all other requirements. The contamination of $Z \rightarrow \tau_\ell \tau_{\text{had}}$ and other processes without $j \rightarrow \tau_{\text{had}}$ is less than 10%.	50
312			
313			
314			
315			
316			
317			
318			

319	6.7	Data events in the VBF category which fail τ_{had} identification but fulfill all other requirements. The contamination of $Z \rightarrow \tau_\ell \tau_{\text{had}}$ and other processes without $j \rightarrow \tau_{\text{had}}$ is less than 10%.	51
320			
321			
322	6.8	Correlations between the τ_{had} BDT identification score and event kinematics in data events in the VBF same-sign region which fail τ_{had} identification but fulfill all other requirements. No strong correlations are observed.	52
323			
324			
325	6.9	Requirements on the τ_{had} jet discriminant, which are defined to have constant signal efficiency as a function of $p_T(\tau_{\text{had}})$, of various operating points for 1-track τ_{had} (left) and 3-track τ_{had} (right).	53
326			
327			
328	6.10	Fake factors in the VBF category measured in the various control regions in data for 1-track τ_{had} (left) and 3-track τ_{had} (right).	53
329			
330	6.11	A pie chart of the composition of $j \rightarrow \tau_{\text{had}}$ processes in the anti-identified CR as predicted by simulation and data (left) and the systematic variations on the composition (right). . .	54
331			
332	6.12	The composition of $j \rightarrow \tau_{\text{had}}$ processes in the anti-identified CR as predicted by simulation and data as a function of event kinematics.	55
333			
334	6.13	The composition of $j \rightarrow \tau_{\text{had}}$ processes in the anti-identified CR as predicted by simulation and data as a function of event kinematics.	56
335			
336	6.14	Fake factors in the VBF category mixed from the various control regions in data for 1-track τ_{had} (left) and 3-track τ_{had} (right). Statistical and systematic uncertainties are shown. . .	57
337			
338	6.15	Cartoon of the signal, control, and validation regions used which are used in the $j \rightarrow \tau_{\text{had}}$ estimate.	58
339			
340	6.16	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the same-sign validation region for various event kinematics. The purity of $j \rightarrow \tau_{\text{had}}$ is $\approx 97\%$. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	59
341			
342			
343	6.17	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the same-sign validation region for various event kinematics. The purity of $j \rightarrow \tau_{\text{had}}$ is $\approx 97\%$. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	60
344			
345			
346	6.18	Comparison of the prediction of identified taus and the $j \rightarrow \tau_{\text{had}}$ prediction, both in simulation, in the signal region for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	61
347			
348			
349	6.19	Comparison of the prediction of identified taus and the $j \rightarrow \tau_{\text{had}}$ prediction, both in simulation, in the signal region for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	62
350			
351			
352	6.20	The fractional uncertainty on the $j \rightarrow \tau_{\text{had}}$ prediction in each bin of the VBF category. . .	63
353	6.21	The fractional uncertainty on the VBF $H \rightarrow \tau_\ell \tau_{\text{had}}$ prediction in each bin of the VBF category for uncertainties pertaining to the jet energy scale.	64
354			
355	6.22	The fractional uncertainty on the VBF $H \rightarrow \tau_\ell \tau_{\text{had}}$ prediction in each bin of the VBF category for uncertainties pertaining to τ_{had} performance, theory, and the luminosity. . .	65
356			
357	7.1	Data and prediction for input variables to the BDT in the $H \rightarrow \tau_\ell \tau_{\text{had}}$ VBF signal region .	67
358	7.2	Display of one of the most signal-like events in the $H \rightarrow \tau_\ell \tau_{\text{had}}$ VBF category in data . The blue track matched to the green cluster indicates an electron, the green track matched to the yellow cluster indicates a τ_{had} , the pink dotted line indicates the E_T^{miss} in the transverse plane, and the turquoise cones indicates the VBF jets. The reconstructed $m_{\tau\tau}^{\text{MMC}} = 127$ GeV and $m_{jj} = 1.53$ TeV.	68
359			
360			
361			
362			
363	7.3	Distributions of the 8 TeV BDT discriminants in all six analysis categories after the global fit	69
364			
365	7.4	Plots of data and prediction which emphasize the most sensitive regions . The individual BDT bins from all six categories are ordered by S/B and plotted on a shared axis (left) and entries in the $m_{\tau\tau}^{\text{MMC}}$ distribution are weighted by $\log(1 + S/B)$ (right).	70
366			
367			

368	7.5	Two-dimensional contours of the fitted signal strength μ comparing VBF and ggF production mechanisms (left) and the local p_0 for the MVA and cuts-based analyses as a function of the Higgs mass hypothesis (right)	70
369			
370			
371	7.6	Comparison of the impact of the statistical and systematic uncertainties on the absolute uncertainty on μ	71
372			
373	7.7	Comparison of the impact and pull of the dominant individual systematic uncertainties on the absolute uncertainty on μ	72
374			
375	7.8	The fitted signal strength μ split by category, final state, and data-taking period	73
376	8.1	Tau trigger rates in 2012 data-taking as a function of instantaneous luminosity for L1 (left) and HLT (right)	75
377			
378	8.2	Significance (p_0) of the $H \rightarrow \tau\tau$ VBF category in 2012 for the $H \rightarrow \tau_{\text{had}}\tau_{\text{had}}$ (left) and $H \rightarrow \tau_{\ell}\tau_{\text{had}}$ (right) analyses as a function of offline or L1 threshold for various objects. . .	77
379			
380	8.3	Topological distributions at L1 for $H \rightarrow \tau_e\tau_{\text{had}}$ MC versus high-pileup ($\langle\mu\rangle = 81$) minimum bias MC.	79
381			
382	8.4	L1 angular resolution for τ_{had} in simulation and data	79
383	8.5	Momentum resolution for τ_{had} in simulation and data at L1 (left) and HLT (right). The resolution is significantly improved at HLT	80
384			
385	8.6	Kinematic distributions in the $\ell + \tau_{\text{had}}$ category of the 8 TeV VBF $H \rightarrow \tau_{\ell}\tau_{\text{had}}$ analysis. .	82
386	8.7	Kinematic distributions in the $\ell + \tau_{\text{had}}$ category of the 8 TeV VBF $H \rightarrow \tau_{\ell}\tau_{\text{had}}$ analysis. .	83
387	8.8	Efficiency for firing the 20 GeV L1 τ_{had} trigger as a function of offline $p_T(\tau_{\text{had}})$ for no isolation requirement (left) and the 2012 isolation requirement (right) for various definitions of the L1 τ_{had} item. The current definition (2×1 EM, 2×2 had.) has the slowest efficiency turn-on. Fits are performed with a Fermi-Dirac distribution.	84
388			
389			
390			
391	8.9	L1 rate for the di- τ_{had} trigger in 14 TeV minimum bias MC for various p_T^{L1} -dependent isolation definitions relative to the 2012 definition: $p_T^{\text{L1,iso}} \leq 4$ GeV. Many options give the same rate (white color). The rate is calculated irrespective of the lowest unscaled single τ_{had} trigger (left) and with a logical OR of it (right).	86
392			
393			
394			
395	8.10	Efficiency for firing the L1 τ_{had} trigger for various p_T^{L1} -dependent isolation definitions which have similar rates, as derived from Fig. 8.9. The 2012 definition is the black line. The efficiency is calculated irrespective of the lowest unscaled single τ_{had} trigger (left) and with a logical OR of it (right).	87
396			
397			
398			
399	8.11	Degradation of E_T^{miss} -related observables at HL-LHC conditions for VBF $H \rightarrow \tau_{\ell}\tau_{\text{had}}$: the E_T^{miss} resolution (left) and reconstructed $m_{\tau\tau}^{\text{MMC}}$ (right) . The underflow of the $m_{\tau\tau}^{\text{MMC}}$ shows the fraction of events which fail the mass reconstruction.	93
400			
401			
402	8.12	Signal efficiency versus background efficiency for scenarios of generic forward tracker coverage and rejection power (left) and zoomed in to lower signal efficiency (right) . A BDT is trained in the VBF category for each scenario.	94
403			
404			
405	8.13	Signal and background HL-LHC predictions of (a) leading jet p_T , (b) sub-leading jet p_T , (c) leading jet η , (d) sub-leading jet η , (e) $\Delta\eta_{jj}$, (f) m_{jj} , (g) $\eta_{\text{leadjet}} \times \eta_{\text{sub-leadjet}}$ and (h) E_T^{miss} . The last bin contains the overflow events.	95
406			
407			
408	8.14	Signal and background HL-LHC predictions of (a) $p_T(\tau_{\text{had}})$, (b) $p_T(\text{lepton})$, (c) $\eta(\tau_{\text{had}})$, (d) $\eta(\text{lepton})$, (e) $\Delta R(\tau_{\text{had}}, \text{lepton})$, (f) MMC (g) $m_{\tau\tau}^{\text{vis}}$ and (h) $m_T(\ell, E_T^{\text{miss}})$. The last bin contains the overflow events.	96
409			
410			
411	8.15	Signal and background HL-LHC predictions of (a) E_T^{miss} ϕ -centrality, (b) lepton η -centrality and (c) p_T^{Total} . The last bin contains the overflow events.	97
412			
413	8.16	Signal and background HL-LHC predictions of the BDT spectrum in the (a) full range and (b) highest bins range . Signal and background are overlaid in (a) and stacked in (b). . .	97
414			

415	A.1	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the $W \rightarrow \ell \nu_\ell$ CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	101
416			
417	A.2	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the $W \rightarrow \ell \nu_\ell$ CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	102
418			
419	A.3	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the QCD CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	103
420			
421	A.4	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the QCD CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	104
422			
423	A.5	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the $Z \rightarrow \ell \ell$ CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	105
424			
425	A.6	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the $Z \rightarrow \ell \ell$ CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	106
426			
427	A.7	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the top CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	107
428			
429	A.8	Comparison of data and $j \rightarrow \tau_{\text{had}}$ prediction in the top CR for various event kinematics. Only statistical uncertainties are shown, and no sign of systematic bias is observed.	108
430			
431	B.1	Signal and background distributions for the full set of the discriminating variables in the 1-track τ_{had} jet discrimination algorithm.	110
432			
433	B.2	Signal and background distributions for the full set of the discriminating variables in the 3-track τ_{had} jet discrimination algorithm.	111
434			
435	C.1	Simulated predictions of $m_{Z \rightarrow \tau \ell \tau_{\text{had}}}$ and $m_{H \rightarrow \tau \ell \tau_{\text{had}}}$ in the boosted category for various $m_{\tau \tau}$ reconstruction algorithms.	113
436			
437	C.2	Simulated predictions of $m_{Z \rightarrow \tau \ell \tau_{\text{had}}}$ and $m_{H \rightarrow \tau \ell \tau_{\text{had}}}$ in the VBF category for various $m_{\tau \tau}$ reconstruction algorithms.	114
438			