

# CSCI 1550: PROBABILISTIC METHODS IN CS

## NOTES

ALEXANDER W. LEE

### 1. EVENTS AND PROBABILITY

**Definition.** A probability space has three components:

- (1) a sample space  $\Omega$ , which is the set of all possible outcomes of the random process modeled by the probability space;
- (2) a family of sets  $\mathcal{F}$  representing the allowable events, where each set in  $\mathcal{F}$  is a subset of the sample space  $\Omega$ ; and
- (3) a probability function  $\Pr : \mathcal{F} \rightarrow \mathbb{R}$  satisfying the following definition.

**Definition.** A probability function is any function  $\Pr : \mathcal{F} \rightarrow \mathbb{R}$  that satisfies the following conditions:

- (1) for any event  $E$ ,  $0 \leq \Pr(E) \leq 1$ ;
- (2)  $\Pr(\Omega) = 1$ ; and
- (3) for any finite or countably infinite sequence of pairwise mutually disjoint events  $E_1, E_2, E_3, \dots$ ,

$$\Pr \left( \bigcup_{i \geq 1} E_i \right) = \sum_{i \geq 1} \Pr(E_i).$$

**Lemma.** For any two events  $E_1$  and  $E_2$ ,

$$\Pr(E_1 \cup E_2) = \Pr(E_1) + \Pr(E_2) - \Pr(E_1 \cap E_2).$$

**Lemma (Union Bound).** For any finite or countably infinite sequence of events  $E_1, E_2, \dots$ ,

$$\Pr \left( \bigcup_{i \geq 1} E_i \right) \leq \sum_{i \geq 1} \Pr(E_i).$$

**Lemma (1.3).** Let  $E_1, \dots, E_n$  be any  $n$  events. Then

$$\begin{aligned} \Pr \left( \bigcup_{i=1}^n E_i \right) &= \sum_{i=1}^n \Pr(E_i) - \sum_{i < j} \Pr(E_i \cap E_j) + \sum_{i < j < k} \Pr(E_i \cap E_j \cap E_k) \\ &\quad - \dots + (-1)^{l+1} \sum_{i_1 < i_2 < \dots < i_l} \Pr \left( \bigcap_{r=1}^l E_{i_r} \right) + \dots \end{aligned}$$

**Definition.** Two events  $E$  and  $F$  are independent if and only if

$$\Pr(E \cap F) = \Pr(E) \cdot \Pr(F).$$

More generally, events  $E_1, E_2, \dots, E_k$  are mutually independent if and only if, for any subset  $I \subseteq [1, k]$ ,

$$\Pr\left(\bigcap_{i \in I} E_i\right) = \prod_{i \in I} \Pr(E_i).$$

**Definition.** The conditional probability that event  $E$  occurs given that event  $F$  occurs is

$$\Pr(E \mid F) = \frac{\Pr(E \cap F)}{\Pr(F)}.$$

The conditional probability is well-defined only if  $\Pr(F) > 0$ .

**Theorem** (Law of Total Probability). Let  $E_1, E_2, \dots, E_n$  be mutually disjoint events in the sample space  $\Omega$ , and let  $\bigcup_{i=1}^n E_i = \Omega$ . Then

$$\Pr(B) = \sum_{i=1}^n \Pr(B \cap E_i) = \sum_{i=1}^n \Pr(B \mid E_i) \Pr(E_i).$$

**Theorem** (Bayes' Law). Assume that  $E_1, E_2, \dots, E_n$  are mutually disjoint events in the sample space  $\Omega$  such that  $\bigcup_{i=1}^n E_i = \Omega$ . Then

$$\Pr(E_j \mid B) = \frac{\Pr(E_j \cap B)}{\Pr(B)} = \frac{\Pr(B \mid E_j) \Pr(E_j)}{\sum_{i=1}^n \Pr(B \mid E_i) \Pr(E_i)}.$$

## 2. DISCRETE RANDOM VARIABLES AND EXPECTATION

**Definition.** A random variable  $X$  on a sample space  $\Omega$  is a real-valued (measurable) function on  $\Omega$ ; that is,  $X : \Omega \rightarrow \mathbb{R}$ . A discrete random variable is a random variable that takes on only a finite or countably infinite numbers of values.

**Definition.** Two random variable  $X$  and  $Y$  are independent if and only if

$$\Pr((X = x) \cap (Y = y)) = \Pr(X = x) \cdot \Pr(Y = y)$$

for all values  $x$  and  $y$ . Similarly, random variables  $X_1, X_2, \dots, X_k$  are mutually independent if and only if, for any subset  $I \subseteq [1, k]$  and any values  $x_i, i \in I$ ,

$$\Pr\left(\bigcap_{i \in I} (X_i = x_i)\right) = \prod_{i \in I} \Pr(X_i = x_i).$$

**Definition.** The expectation of a discrete random variable  $X$ , denoted by  $\mathbf{E}[X]$ , is given by

$$\mathbf{E}[X] = \sum_i i \Pr(X = i),$$

where the summation is over all values in the range of  $X$ . The expectation is finite if  $\sum_i |i| \Pr(X = i)$  converges; otherwise, the expectation is unbounded.

**Theorem** (Linearity of Expectations). For any finite collection of discrete random variables  $X_1, X_2, \dots, X_n$  with finite expectations,

$$\mathbf{E}\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n \mathbf{E}[X_i].$$

**Lemma.** For any constant  $c$  and discrete random variable  $X$ ,

$$\mathbf{E}[cX] = c\mathbf{E}[X].$$

**Definition.** Consider a sequence of  $n$  independent experiments, each of which succeeds with probability  $p$ . If we let  $X$  represent the number of successes in the  $n$  experiments, then  $X$  has a binomial distribution. A binomial random variable  $X$  with parameters  $n$  and  $p$ , denoted by  $B(n, p)$ , is defined by the following probability distribution on  $j = 0, 1, 2, \dots, n$ :

$$\Pr(X = j) = \binom{n}{j} p^j (1 - p)^{n-j}.$$

That is, the binomial random variable  $X$  equals  $j$  when there are exactly  $j$  successes and  $n - j$  failures in  $n$  independent experiments, each of which is successful with probability  $p$ .

**Lemma.** For a binomial random variable  $X$  with parameters  $n$  and  $p$ ,

$$\mathbf{E}[X] = np.$$

**Definition.**

$$\mathbf{E}[Y \mid Z = z] = \sum_y y \Pr(Y = y \mid Z = z),$$

where the summation is over all  $y$  in the range of  $Y$ .

**Lemma.** For any random variables  $X$  and  $Y$ ,

$$\mathbf{E}[X] = \sum_y \Pr(Y = y) \mathbf{E}[X \mid Y = y],$$

where the sum is over all values in the range of  $Y$  and all of the expectations exist.

**Lemma.** For any finite collection of discrete random variables  $X_1, X_2, \dots, X_n$  with finite expectations and for any random variable  $Y$ ,

$$\mathbf{E} \left[ \sum_{i=1}^n X_i \mid Y = y \right] = \sum_{i=1}^n \mathbf{E}[X_i \mid Y = y].$$

**Definition.** The expression  $\mathbf{E}[X \mid Y]$  is a random variable  $f(Y)$  that takes on the value  $\mathbf{E}[X \mid Y = z]$  when  $Y = z$ .

**Theorem.**

$$\mathbf{E}[Y] = \mathbf{E}[\mathbf{E}[Y \mid Z]].$$

**Definition.** We perform a sequence of independent trials until the first success, where each trial succeeds with probability  $p$ . A geometric random variable  $X$  with parameter  $p$  is given by the following probability distribution on  $n = 1, 2, \dots$ :

$$\Pr(X = n) = (1 - p)^{n-1} p.$$

That is, for the geometric random variable  $X$  to equal  $n$ , there must be  $n-1$  failures, followed by a success.

**Lemma.** For a geometric random variable  $X$  with parameter  $p$  and for  $n > 0$ ,

$$\Pr(X = n + k \mid X > k) = \Pr(X = n).$$

**Lemma.** Let  $X$  be a discrete random variable that takes on only non-negative integer values. Then

$$\mathbf{E}[X] = \sum_{i=1}^{\infty} \Pr(X \geq i).$$

**Lemma.** For a geometric random variable  $X$  with parameter  $p$ ,

$$\mathbf{E}[X] = \frac{1}{p}.$$

**Lemma.** The harmonic number  $H(n) = \sum_{i=1}^n 1/i$  satisfies  $H(n) = \ln n + \Theta(1)$ .

### 3. MOMENTS AND DEVIATIONS

**Theorem** (Markov's Inequality). Let  $X$  be a random variable that assumes only non-negative values. Then, for all  $a > 0$ ,

$$\Pr(X \geq a) \leq \frac{\mathbf{E}[X]}{a}.$$

**Definition.** The  $k$ th moment of a random variable  $X$  is  $\mathbf{E}[X^k]$ .

**Definition.** The variance of a random variable  $X$  is defined as

$$\mathbf{Var}[X] = \mathbf{E}[(X - \mathbf{E}[X])^2] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2.$$

The standard deviation of a random variable  $X$  is

$$\sigma[X] = \sqrt{\mathbf{Var}[X]}.$$

**Definition.** The covariance of two random variables  $X$  and  $Y$  is

$$\mathbf{Cov}(X, Y) = \mathbf{E}[(X - \mathbf{E}[X])(Y - \mathbf{E}[Y])].$$

**Theorem.** For any two random variables  $X$  and  $Y$ ,

$$\mathbf{Var}[X + Y] = \mathbf{Var}[X] + \mathbf{Var}[Y] + 2\mathbf{Cov}(X, Y).$$

**Theorem.** If  $X$  and  $Y$  are two independent random variables, then

$$\mathbf{E}[X \cdot Y] = \mathbf{E}[X] \cdot \mathbf{E}[Y].$$

**Corollary.** If  $X$  and  $Y$  are independent random variables, then

$$\mathbf{Cov}(X, Y) = 0$$

and

$$\mathbf{Var}[X + Y] = \mathbf{Var}[X] + \mathbf{Var}[Y].$$

**Theorem.** Let  $X_1, X_2, \dots, X_n$  be mutually independent random variables. Then

$$\mathbf{Var}\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n \mathbf{Var}[X_i].$$

**Lemma.** For a binomial random variable  $X$  with parameters  $n$  and  $p$ ,

$$\mathbf{Var}[X] = np(1 - p).$$

**Theorem** (Chebyshev's Inequality). For any  $a > 0$ ,

$$\Pr(|X - \mathbf{E}[X]| \geq a) \leq \frac{\mathbf{Var}[X]}{a^2}.$$

**Corollary.** For any  $t > 1$ ,

$$\Pr(|X - \mathbf{E}[X]| \geq t \cdot \sigma[X]) \leq \frac{1}{t^2} \text{ and}$$

$$\Pr(|X - \mathbf{E}[X]| \geq t \cdot \mathbf{E}[X]) \leq \frac{\mathbf{Var}[X]}{t^2(\mathbf{E}[X])^2}.$$

**Lemma.** *For a geometric random variable  $X$  with parameter  $p$ ,*

$$\mathbf{Var}[X] = (1 - p)/p^2.$$