

Mathematical Programs with Complementarity Constraints and Related Problems

Prof. Dr. Alexandra Schwartz
schwartz@gsc.tu-darmstadt.de



TECHNISCHE
UNIVERSITÄT
DARMSTADT



You can find

- ▶ my slides
- ▶ and some more extensive lecture notes

at

github.com/alexandrabschwartz/Winterschool2018

If you have any questions, please come to me during the week or contact me at

schwartz@gsc.tu-darmstadt.de



At the end of the week you should

- ▶ know the definition and applications of MPCCs,
- ▶ be able to explain why MPCCs cannot be solved directly using standard theory,
- ▶ understand the origins and relations of different optimality conditions for MPCCs,
- ▶ know what to look for in a relaxation algorithm for MPCCs,
- ▶ be able to transfer the MPCC ideas to some related problem classes.




- ▶ Luo, Pang, Ralph: *Mathematical Programs with Equilibrium Constraints*, Cambridge University Press
- ▶ Outrata, Kocvara, Zowe: *Nonsmooth Approach to Optimization Problems with Equilibrium Constraints*, Kluwer Academic Publishers
- ▶ Mordukhovich: *Variational Analysis and Generalized Differentiation I and II*, Springer
- ▶ Rockafellar, Wets: *Variational Analysis*, Springer
- ▶ Numerous articles by Jane Ye, Jiri Outrata, Christian Kanzow, and many others

Contents of the Course




TECHNISCHE
UNIVERSITÄT
DARMSTADT



What is the plan for today?



TECHNISCHE
UNIVERSITÄT
DARMSTADT





Problem Formulation and Examples


Mathematical Program with Complementarity Constraints (MPCC)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A mathematical program with complementarity constraints (MPCC) is of the form

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0, \\ 0 \leq G(x) \perp H(x) \geq 0.$$



We assume that

- ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$, and $G, H : \mathbb{R}^n \rightarrow \mathbb{R}^q$ are continuously differentiable,
- ▶ the feasible set

$$X := \{x \in \mathbb{R}^n \mid g(x) \leq 0, \quad h(x) = 0, \quad 0 \leq G(x) \perp H(x) \geq 0\}$$

is nonempty.

Mathematical Program with Equilibrium Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A mathematical program with equilibrium constraints (MPEC) is of the form

$$\min_{x,y} f(x, y) \quad \text{s.t.} \quad x \in X, y \in S(x),$$

where $x \Rightarrow S(x)$ is the solution set of the equilibrium constraint, e.g. of

- ▶ a lower-level optimization problem,
- ▶ a lower-level Nash game,
- ▶ or a physical equilibrium.

Mathematical Program with Equilibrium Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A **mathematical program with equilibrium constraints (MPEC)** is of the form

$$\min_{x,y} f(x,y) \quad \text{s.t.} \quad x \in X, y \in S(x),$$

where $x \mapsto S(x)$ is the solution set of the **equilibrium constraint**, e.g. of

- ▶ a lower-level optimization problem,
- ▶ a lower-level Nash game,
- ▶ or a physical equilibrium.

The solution set $S(x)$ can often be described using

- ▶ complementarity constraints $0 \leq G(x,y) \perp H(x,y) \geq 0$,
- ▶ variational inequalities $F(y^*)^T(y - y^*) \geq 0 \quad \forall y \in Y$,
- ▶ generalized equations $0 \in F(y^*) + N_Y(y^*)$.

Example: Bilevel Optimization



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A **bilevel optimization problem** (in optimistic formulation) is of the form

$$\min_{x,y} f(x,y) \quad \text{s.t.} \quad x \in X, y \in \underset{y}{\operatorname{argmin}}\{F(x,y) \mid y \in Y(x)\},$$

where

$$Y(x) = \{y \mid G(x,y) \leq 0, H(x,y) = 0\}.$$

Example: Bilevel Optimization



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A **bilevel optimization problem** (in optimistic formulation) is of the form

$$\min_{x,y} f(x,y) \quad \text{s.t.} \quad x \in X, y \in \underset{y}{\operatorname{argmin}}\{F(x,y) \mid y \in Y(x)\},$$

where

$$Y(x) = \{y \mid G(x,y) \leq 0, H(x,y) = 0\}.$$

Exercise

How can you rewrite the lower level $y \in \underset{y}{\operatorname{argmin}}\{F(x,y) \mid y \in Y(x)\}$ using

- ▶ complementarity constraints,
- ▶ a variational inequality,
- ▶ a generalized equation?

Example: Reformulations of a Bilevel Problem



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the lower level $y \in \operatorname{argmin}_y \{F(x, y) \mid y \in Y(x)\}$.

- Under a suitable constraint qualification, we can replace the lower level by the KKT conditions

$$\nabla_y F(x, y) + \nabla_y G(x, y)\lambda + \nabla_y H(x, y)\mu = 0,$$

$$0 \leq \lambda \perp G(x, y) \leq 0,$$

$$H(x, y) = 0.$$

Example: Reformulations of a Bilevel Problem



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the lower level $y \in \operatorname{argmin}_y \{F(x, y) \mid y \in Y(x)\}$.

- Under a suitable constraint qualification, we can replace the lower level by the **KKT conditions**

$$\nabla_y F(x, y) + \nabla_y G(x, y)\lambda + \nabla_y H(x, y)\mu = 0,$$

$$0 \leq \lambda \perp G(x, y) \leq 0,$$

$$H(x, y) = 0.$$

- If $Y(x)$ is convex, we can use the **variational inequality**

$$\nabla_y F(x, y)^T (z - y) \geq 0 \quad \forall z \in Y(x).$$

Example: Reformulations of a Bilevel Problem



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the lower level $y \in \operatorname{argmin}_y \{F(x, y) \mid y \in Y(x)\}$.

- Under a suitable constraint qualification, we can replace the lower level by the **KKT conditions**

$$\nabla_y F(x, y) + \nabla_y G(x, y)\lambda + \nabla_y H(x, y)\mu = 0,$$

$$0 \leq \lambda \perp G(x, y) \leq 0,$$

$$H(x, y) = 0.$$

- If $Y(x)$ is convex, we can use the **variational inequality**

$$\nabla_y F(x, y)^T (z - y) \geq 0 \quad \forall z \in Y(x).$$

- Or we rewrite **B-stationarity** as a generalized equation

$$\nabla_y F(x, y)^T d \geq 0 \quad \forall d \in T_{Y(x)}(y) \quad \Longleftrightarrow \quad 0 \in \nabla_y F(x, y) + N_{Y(x)}(y).$$



Challenges

Nonlinear Program (NLP)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A **nonlinear program (NLP)** is of the form

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0.$$

Denote the feasible set by

$$Z := \{x \in \mathbb{R}^n \mid g(x) \leq 0, \quad h(x) = 0\}$$

and for a point $x^* \in Z$ define the set of **active inequalities**

$$I_g(x^*) = \{i \mid g_i(x^*) = 0\}.$$

Optimality Conditions for NLPs: B-Stationarity





TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let x^* be a local minimum of NLP. Then x^* is **B-stationary**, i.e.

$$\nabla f(x^*)^T d \geq 0 \quad \forall d \in T_Z(x^*),$$

where the (**Bouligand**) **tangent cone** is defined as

$$T_X(x^*) := \{d \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow_X x^*, (t_k)_k \geq 0 : t_k(x^k - x^*) \rightarrow d\}.$$



Constraint Qualifications for NLPs



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Under a **constraint qualification** in x^* such as

- **LICQ**: the gradients

$$\nabla g_i(x^*) \ (i \in I_g(x^*)), \quad \nabla h_i(x^*) \ (i = 1, \dots, p)$$

are linearly independent;

- **MFCQ**: the gradients

$$\nabla g_i(x^*) \ (i \in I_g(x^*)) \quad \text{and} \quad \nabla h_i(x^*) \ (i = 1, \dots, p)$$

are **positively** linearly independent;

- **Abadie CQ**: $T_Z(x^*) = L_Z(x^*)$;
- **Guignard CQ**: $T_Z(x^*)^\circ = L_Z(x^*)^\circ$

we can replace the tangent cone by the simpler **linearized tangent cone**

$$\begin{aligned} L_Z(x^*) = \{d \in \mathbb{R}^n \mid & \nabla g_i(x^*)^T d \leq 0 \quad \forall i \in I_g(x^*), \\ & \nabla h_i(x^*)^T d = 0 \quad \forall i = 1, \dots, p\} \end{aligned}$$

Optimality Conditions for NLPs:

KKT Conditions



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let x^* be a local minimum of NLP, where a CQ holds. Then x^* is a **KKT point**, i.e.

$$\nabla f(x^*)^T d \geq 0 \quad \forall d \in L_Z(x^*),$$

which is equivalent to the existence of **(Lagrange) multipliers** $\lambda \in \mathbb{R}^n$ and $\mu \in \mathbb{R}^m$ such that

$$\nabla f(x^*) + \nabla g(x^*)\lambda + \nabla h(x^*)\mu = 0,$$

$$0 \leq \lambda \perp g(x^*) \leq 0,$$

$$h(x^*) = 0.$$



Exercise

Show that the following conditions are equivalent for $a, b \in \mathbb{R}^n$:

- ▶ $0 \leq a \perp b \geq 0$
- ▶ $a \geq 0, b \geq 0, a \circ b = (a_i b_i)_i = 0$
- ▶ $a \geq 0, b \geq 0, a \circ b \leq 0$
- ▶ $a \geq 0, b \geq 0, a^T b \leq 0$



Exercise

Show that the following conditions are equivalent for $a, b \in \mathbb{R}^n$:

- ▶ $0 \leq a \perp b \geq 0$
- ▶ $a \geq 0, b \geq 0, a \circ b = (a_i b_i)_i = 0$
- ▶ $a \geq 0, b \geq 0, a \circ b \leq 0$
- ▶ $a \geq 0, b \geq 0, a^T b \leq 0$

Thus, we can reformulate MPCC as the equivalent NLP

$$\begin{aligned} \min_x f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & G(x) \geq 0, \quad H(x) \geq 0, \quad G(x) \circ H(x) = 0. \end{aligned}$$



Exercise

Show that the following conditions are equivalent for $a, b \in \mathbb{R}^n$:

- ▶ $0 \leq a \perp b \geq 0$
- ▶ $a \geq 0, b \geq 0, a \circ b = (a_i b_i)_i = 0$
- ▶ $a \geq 0, b \geq 0, a \circ b \leq 0$
- ▶ $a \geq 0, b \geq 0, a^T b \leq 0$

Thus, we can reformulate MPCC as the equivalent NLP

$$\begin{aligned} \min_x f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & G(x) \geq 0, \quad H(x) \geq 0, \quad G(x) \circ H(x) = 0. \end{aligned}$$

Why are we not done here?

NLP Reformulation versus Constraint Qualifications




TECHNISCHE
UNIVERSITÄT
DARMSTADT

Exercise

Consider the simple MPCC

$$\min_{x \in \mathbb{R}^2} f(x) \quad \text{s.t.} \quad 0 \leq x_1 \perp x_2 \geq 0.$$

In which feasible points is LICQ / MFCQ / Abadie CQ / Guignard CQ for the NLP reformulation satisfied?






Exercise

Consider the simple MPCC

$$\min_{x \in \mathbb{R}^2} f(x) \quad \text{s.t.} \quad 0 \leq x_1 \perp x_2 \geq 0.$$

In which feasible points is LICQ / MFCQ / Abadie CQ / Guignard CQ for the NLP reformulation satisfied?



For general MPCCs one knows that

- ▶ LICQ and MFCQ are violated in all feasible points,
- ▶ Abadie CQ is usually violated,
- ▶ only Guignard CQ has a chance to hold.

A “simple” MPCC by S. Scholtes



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the linear MPCC

$$\begin{aligned} \min_{x \in \mathbb{R}^3} \quad & x_1 + x_2 - x_3 \quad \text{s.t.} \quad -4x_1 + x_3 \leq 0, \\ & -4x_2 + x_3 \leq 0, \\ & 0 \leq x_1 \perp x_2 \geq 0. \end{aligned}$$

A “simple” MPCC by S. Scholtes



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the linear MPCC

$$\begin{aligned} \min_{x \in \mathbb{R}^3} \quad & x_1 + x_2 - x_3 \quad \text{s.t.} \quad -4x_1 + x_3 \leq 0, \\ & -4x_2 + x_3 \leq 0, \\ & 0 \leq x_1 \perp x_2 \geq 0. \end{aligned}$$

Its global minimum $x^* = (0, 0, 0)$ is not a KKT point:

$$\begin{aligned} \lambda_3 &= 1 - 4\lambda_1 \geq 0, \\ \lambda_4 &= 1 - 4\lambda_2 \geq 0, \\ \lambda_1 + \lambda_2 &= 1, \text{ where } \lambda_1 \geq 0, \lambda_2 \geq 0, \end{aligned}$$

Consequently Guignard CQ cannot be satisfied.

An even “simpler” MPCC by A. Schiela




TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the linear MPCC

$$\min_{x \in \mathbb{R}^2} -x_2 \quad \text{s.t.} \quad x_2 - x_1 \leq 0, \quad 0 \leq x_1 \perp x_2 \geq 0.$$

Then $X = [0, \infty) \times \{0\}$ and $x^* = (0, 0)$ is a global minimum with

$$T_X(x^*) = [0, \infty) \times \{0\} \quad \text{and} \quad L_X(x^*) = \{d \in \mathbb{R}^2 \mid d_1 \geq 0, d_2 \geq 0, d_2 - d_1 \leq 0\},$$
$$T_X(x^*)^\circ = (-\infty, 0] \times \mathbb{R} \quad \text{and} \quad L_X(x^*)^\circ = \{w \in \mathbb{R}^2 \mid w_1 \leq 0, w_1 + w_2 \leq 0\}.$$



Challenges of the NLP Reformulation of MPCC



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- Most standard CQs for the NLP reformulation of MPCC are violated.

Challenges of the NLP Reformulation of MPCC



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ Most standard CQs for the NLP reformulation of MPCC are violated.
- ▶ Without CQ the KKT conditions are not necessary optimality conditions.

Challenges of the NLP Reformulation of MPCC



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ Most standard CQs for the NLP reformulation of MPCC are violated.
- ▶ Without CQ the KKT conditions are not necessary optimality conditions.
- ▶ The **Fritz-John conditions**, i.e. the existence of $\alpha \geq 0$, $\lambda \in \mathbb{R}^n$ and $\mu \in \mathbb{R}^m$ such that

$$\alpha \nabla f(x^*) + \nabla g(x^*)\lambda + \nabla h(x^*)\mu = 0,$$

$$0 \leq \lambda \perp g(x^*) \leq 0,$$

$$h(x^*) = 0,$$

$$(\alpha, \lambda, \mu) \neq 0$$

are necessary optimality conditions without a CQ, but they are satisfied at every feasible point of the NLP reformulation.



Background: Normal Cones and Subdifferentials

Polar Cone and Fréchet Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $X \subseteq \mathbb{R}^n$ be nonempty and $x^* \in X$.

- The (Bouligand) tangent cone is defined as

$$T_X(x^*) := \{d \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow_X x^*, (t_k)_k \geq 0 : t_k(x^k - x^*) \rightarrow d\}.$$

Polar Cone and Fréchet Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $X \subseteq \mathbb{R}^n$ be nonempty and $x^* \in X$.

- ▶ The (Bouligand) tangent cone is defined as

$$T_X(x^*) := \{d \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow_X x^*, (t_k)_k \geq 0 : t_k(x^k - x^*) \rightarrow d\}.$$

- ▶ The polar cone to a set $K \subseteq \mathbb{R}^n$ is defined as

$$K^\circ := \{w \in \mathbb{R}^n \mid w^T d \leq 0 \quad \forall d \in K\}.$$

Polar Cone and Fréchet Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $X \subseteq \mathbb{R}^n$ be nonempty and $x^* \in X$.

- ▶ The (Bouligand) tangent cone is defined as

$$T_X(x^*) := \{d \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow_X x^*, (t_k)_k \geq 0 : t_k(x^k - x^*) \rightarrow d\}.$$

- ▶ The polar cone to a set $K \subseteq \mathbb{R}^n$ is defined as

$$K^\circ := \{w \in \mathbb{R}^n \mid w^T d \leq 0 \quad \forall d \in K\}.$$

- ▶ For a polyhedral cone $P = \{d \mid A^T d \leq 0, B^T d = 0\}$ one knows

$$P^\circ = \{w = A\lambda + B\mu \mid \lambda \geq 0\}.$$

Polar Cone and Fréchet Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $X \subseteq \mathbb{R}^n$ be nonempty and $x^* \in X$.

- ▶ The (Bouligand) tangent cone is defined as

$$T_X(x^*) := \{d \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow_X x^*, (t_k)_k \geq 0 : t_k(x^k - x^*) \rightarrow d\}.$$

- ▶ The polar cone to a set $K \subseteq \mathbb{R}^n$ is defined as

$$K^\circ := \{w \in \mathbb{R}^n \mid w^T d \leq 0 \quad \forall d \in K\}.$$

- ▶ For a polyhedral cone $P = \{d \mid A^T d \leq 0, B^T d = 0\}$ one knows

$$P^\circ = \{w = A\lambda + B\mu \mid \lambda \geq 0\}.$$

- ▶ The Fréchet normal cone of X at x^* is defined as

$$N_X^F(x^*) := T_X(x^*)^\circ = \{w \in \mathbb{R}^n \mid \limsup_{x \rightarrow x^*, x \in X \setminus \{x^*\}} w^T \frac{x - x^*}{\|x - x^*\|} \leq 0\}.$$

For $x^* \notin X$ one defines $N_X^F(x^*) := \emptyset$.

Examples for the Fréchet Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Exercise

Consider the feasible sets

- ▶ $X = \{0\} \subset \mathbb{R}$,
- ▶ $X = (-\infty, 0] \subset \mathbb{R}$,
- ▶ $X = \{x \in \mathbb{R}^2 \mid 0 \leq x_1 \perp x_2 \geq 0\} \subset \mathbb{R}^2$

and compute the Fréchet normal cone at all feasible points.



Limiting Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

The **limiting or Mordukhovich normal cone** of X at x^* is defined as

$$N_X^M(x^*) := \{w \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow x^*, w^k \in N_X^F(x^k) : w^k \rightarrow w\}.$$

For $x^* \notin X$ one defines $N_X^M(x^*) := \emptyset$.



The **limiting or Mordukhovich normal cone** of X at x^* is defined as

$$N_X^M(x^*) := \{w \in \mathbb{R}^n \mid \exists (x^k)_k \rightarrow x^*, w^k \in N_X^F(x^k) : w^k \rightarrow w\}.$$

For $x^* \notin X$ one defines $N_X^M(x^*) := \emptyset$.

Exercise

Consider the feasible set $X = \{x \in \mathbb{R}^2 \mid 0 \leq x_1 \perp x_2 \geq 0\}$ and compute the limiting normal cone at $x^ = (0, 0)^T$.*

A Calculus Rule for the Limiting Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

$$N_{A \cap B}^M(x^*) = ???$$

A Calculus Rule for the Limiting Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

$$N_{A \cap B}^M(x^*) = ???$$

- Let $A, B \subseteq \mathbb{R}^n$ be nonempty and closed, $x^* \in A \cap B$ and assume that the set-valued function

$$M(y) := \{x \in A \mid x + y \in B\}$$

is **calm** at $(0, x^*)$. Then

$$N_{A \cap B}^M(x^*) \subseteq N_A^M(x^*) + N_B^M(x^*).$$

A Calculus Rule for the Limiting Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

$$N_{A \cap B}^M(x^*) = ???$$

- Let $A, B \subseteq \mathbb{R}^n$ be nonempty and closed, $x^* \in A \cap B$ and assume that the set-valued function

$$M(y) := \{x \in A \mid x + y \in B\}$$

is **calm** at $(0, x^*)$. Then

$$N_{A \cap B}^M(x^*) \subseteq N_A^M(x^*) + N_B^M(x^*).$$

- A set-valued map $M : Y \subseteq \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ is **calm** at $(y^*, x^*) \in \text{graph}(M)$ if there exist $\delta > 0, \varepsilon > 0, L > 0$ such that

$$M(y) \cap B_\varepsilon(x^*) \subseteq M(y^*) + L\|y - y^*\|B_1(0) \quad \forall y \in B_\delta(y^*).$$

A Calculus Rule for the Limiting Normal Cone



TECHNISCHE
UNIVERSITÄT
DARMSTADT

$$N_{A \cap B}^M(x^*) = ???$$

- ▶ Let $A, B \subseteq \mathbb{R}^n$ be nonempty and closed, $x^* \in A \cap B$ and assume that the set-valued function

$$M(y) := \{x \in A \mid x + y \in B\}$$

is **calm** at $(0, x^*)$. Then

$$N_{A \cap B}^M(x^*) \subseteq N_A^M(x^*) + N_B^M(x^*).$$

- ▶ A set-valued map $M : Y \subseteq \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ is **calm** at $(y^*, x^*) \in \text{graph}(M)$ if there exist $\delta > 0, \varepsilon > 0, L > 0$ such that

$$M(y) \cap B_\varepsilon(x^*) \subseteq M(y^*) + L\|y - y^*\|B_1(0) \quad \forall y \in B_\delta(y^*).$$

- ▶ **Polyhedral multifunctions** M , i.e. set-valued maps whose graph can be written as the union of finitely many polyhedra, are calm at all $(y^*, x^*) \in \text{graph}(M)$.

The Bouligand and Clarke Subdifferential



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider a locally Lipschitz continuous map $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

- **Rademacher's Theorem:** F is differentiable almost everywhere, i.e. the complement of

$$D_F := \{x \in \mathbb{R}^n \mid F \text{ is differentiable in } x\}.$$

has Lebesgue measure zero.

The Bouligand and Clarke Subdifferential



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider a locally Lipschitz continuous map $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

- **Rademacher's Theorem:** F is differentiable almost everywhere, i.e. the complement of

$$D_F := \{x \in \mathbb{R}^n \mid F \text{ is differentiable in } x\}.$$

has Lebesgue measure zero.

- The **Bouligand subdifferential** of F at $x^* \in \mathbb{R}^n$ is defined as

$$\partial^B F(x^*) := \{M \in \mathbb{R}^{m \times n} \mid \exists (x^k)_k \rightarrow_{D_F} x^* : F'(x^k) \rightarrow M\}.$$

The Bouligand and Clarke Subdifferential



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider a locally Lipschitz continuous map $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

- **Rademacher's Theorem:** F is differentiable almost everywhere, i.e. the complement of

$$D_F := \{x \in \mathbb{R}^n \mid F \text{ is differentiable in } x\}.$$

has Lebesgue measure zero.

- The **Bouligand subdifferential** of F at $x^* \in \mathbb{R}^n$ is defined as

$$\partial^B F(x^*) := \{M \in \mathbb{R}^{m \times n} \mid \exists (x^k)_k \rightarrow_{D_F} x^* : F'(x^k) \rightarrow M\}.$$

- The **Clarke subdifferential** of F at $x^* \in \mathbb{R}^n$ is defined as

$$\partial^C F(x^*) := \text{conv } \partial^B F(x^*).$$

Examples for the Bouligand and Clarke Subdifferential






TECHNISCHE
UNIVERSITÄT
DARMSTADT

Exercise

Consider the functions

- ▶ $h(x) = |x|$,
- ▶ $g(x) = \max\{0, x\}$,
- ▶ $c(x, y) = \min\{x, y\}$

and compute the Bouligand and Clarke subdifferentials.



Properties of the Clarke Subdifferential



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be locally Lipschitz continuous and $x^* \in \mathbb{R}^n$.

- ▶ $\partial F(x^*)$ is nonempty, compact, and convex.
- ▶ If F is continuously differentiable around x^* , then $\partial^C F(x^*) = \{F'(x^*)\}$.

Properties of the Clarke Subdifferential



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be locally Lipschitz continuous and $x^* \in \mathbb{R}^n$.

- ▶ $\partial F(x^*)$ is nonempty, compact, and convex.
- ▶ If F is continuously differentiable around x^* , then $\partial^C F(x^*) = \{F'(x^*)\}$.
- ▶ Let $f, f_1, f_2 : \mathbb{R}^n \rightarrow \mathbb{R}$, $G : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be locally Lipschitz continuous and $x \in \mathbb{R}^n, y \in \mathbb{R}^m$. Then the following holds:
 - ▶ $\partial^C(cf)(x) = c\partial^C f(x)$ for all $c \in \mathbb{R}$.
 - ▶ $\partial^C(f_1 + f_2)(x) \subseteq \partial^C f_1(x) + \partial^C f_2(x)$.
 - ▶ $\partial^C(f_1 \cdot f_2)(x) \subseteq f_2(x)\partial^C f_1(x) + f_1(x)\partial^C f_2(x)$.
 - ▶ $\partial(f \circ G)(y) \subseteq \text{conv}(\partial^C f(G(y)) \cdot \partial^C G(y))$ with equality if f is continuously differentiable or f convex and G continuously differentiable.



Let f, g, h be locally Lipschitz continuous and x^* be a local minimum of

$$\min_x f(x) \text{ s.t. } x \in Z := \{x \in \mathbb{R}^n \mid g(x) \leq 0, h(x) = 0\}.$$

Then there exist multipliers $\alpha \in \mathbb{R}, \lambda \in \mathbb{R}^m, \mu \in \mathbb{R}^p$ such that

- ▶ $(\alpha, \lambda, \mu) \neq (0, 0, 0)$,
- ▶ $\alpha \geq 0$,
- ▶ $\lambda \geq 0$ and $\lambda_i = 0$ for all $i \notin I_g(x^*)$
- ▶ and

$$0 \in \alpha \partial^C f(x^*) + \sum_{i=1}^m \lambda_i \partial^C g_i(x^*) + \sum_{i=1}^p \mu_i \partial^C h_i(x^*)$$

What should you remember?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ The definition of MPCC.
- ▶ Why MPCC cannot be solved using standard NLP theory.
- ▶ The definition and calculus rule for the Fréchet and limiting normal cone.
- ▶ The definition and basic properties of the Clarke subdifferential.

What is the plan for tomorrow?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

