

Mathematical Programs with Complementarity Constraints and Related Problems

Prof. Dr. Alexandra Schwartz
schwartz@gsc.tu-darmstadt.de



TECHNISCHE
UNIVERSITÄT
DARMSTADT



You can find

- ▶ my slides
- ▶ and some more extensive lecture notes

at

github.com/alexandrab Schwartz/Winterschool2018


If you have any questions, please come to me during the week or contact me at

schwartz@gsc.tu-darmstadt.de

Contents of the Course




TECHNISCHE
UNIVERSITÄT
DARMSTADT



What did we do yesterday?







TECHNISCHE
UNIVERSITÄT
DARMSTADT



Relaxation Methods: Exact KKT Points







TECHNISCHE
UNIVERSITÄT
DARMSTADT

relaxation	global	local	nonsmooth	kinked
authors	Scholtes	Steff./Ulbr.	Kadrani et al.	Kanzow, S.
geometry				
stationarity under CQ	C-stat. MPEC-MFCQ	C-stat. MPEC-CPLD	M-stat. MPEC-CPLD	M-stat. MPEC-CPLD

Relaxation Methods: Inexact KKT Points


Let $t_k \downarrow 0$ and $\varepsilon_k = o(t_k)$. Let x^k be ε_k -KKT points of $\text{NLP}(t_k)$ and assume $x^k \rightarrow x^*$. Then the limit x^* satisfies:

relaxation	global	local	nonsmooth	kinked
authors	Scholtes	Steff./Ulbr.	Kadrani et al.	Kanzow, S.
geometry				
stationarity under CQ	C-stat. MPEC-MFCQ	W-stat. MPEC-MFCQ	W-stat. MPEC-MFCQ	W-stat. MPEC-MFCQ

What is the plan for today?



TECHNISCHE
UNIVERSITÄT
DARMSTADT





Vanishing Constraints


Mathematical Program with Vanishing Constraints (MPVC)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A mathematical program with vanishing constraints (MPVC) is of the form

$$\begin{aligned} \min_x f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & H(x) \geq 0, \quad G(x) \circ H(x) \geq 0. \end{aligned}$$




Mathematical Program with Vanishing Constraints (MPVC)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

A mathematical program with vanishing constraints (MPVC) is of the form

$$\begin{aligned} \min_x f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & H(x) \geq 0, \quad G(x) \circ H(x) \geq 0. \end{aligned}$$




Two cases are possible:

- ▶ $H_i(x^*) > 0$ and thus $G_i(x^*) \leq 0$
- ▶ $H_i(x^*) = 0$ and then $G_i(x^*) \in \mathbb{R}$ arbitrarily.

Example: Truss Design



TECHNISCHE
UNIVERSITÄT
DARMSTADT



Given some anchor points and a load point, how should you construct a load bearing truss?

Example: Truss Design (continued)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

$$\min_{a,u} \sum_i l_i a_i$$

$$\text{s.t. } 0 \leq a \leq a_{\max}$$

$$K(a)u = f$$

$$f^T u \leq c$$

$$(\sigma_i(u)^2 - \sigma_{\max}^2) a_i \leq 0$$

minimize volume of the truss

bounds on cross sections

force equilibrium

bound on the deformation

bound on the stress on bar i

Variables

a_i cross sectional area of bar i

u_i nodal displacements at the hinges

Constants

f applied force

l_i length of bar i

$K(a)$ stiffness matrix of the truss

$\sigma_i(u)$ stress on bar i

Relation between Vanishing and Complementarity Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

One can reformulate an MPVC as an MPCC using slack variables as follows:

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, h(x) = 0, \\ & G(x) - y \leq 0, \\ & 0 \leq H(x) \perp y \geq 0. \end{aligned}$$

Relation between Vanishing and Complementarity Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

One can reformulate an MPVC as an MPCC using slack variables as follows:

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, h(x) = 0, \\ & G(x) - y \leq 0, \\ & 0 \leq H(x) \perp y \geq 0. \end{aligned}$$

Exercise

How is the relation between the MPCC reformulation and the original MPVC?

Relation between Vanishing and Complementarity Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

One can reformulate an MPVC as an MPCC using slack variables as follows:

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, h(x) = 0, \\ & G(x) - y \leq 0, \\ & 0 \leq H(x) \perp y \geq 0. \end{aligned}$$

Then x^* is a solution of the MPVC if and only if (x^*, y^*) with

$$y_i^* \begin{cases} = 0 & \text{if } H_i(x^*) > 0, \\ \geq \max\{0, G_i(x^*)\} & \text{if } H_i(x^*) = 0 \end{cases}$$

if a solution of the corresponding MPCC. But y^* is not uniquely determined.




Vanishing Constraints:

$$H(x)_i \geq 0, \quad G(x)_i H(x)_i \geq 0 \quad \forall i = 1, \dots, q$$

Let x^* be feasible for MPVC.

- ▶ If $H_i(x^*) = 0$ for at least one index $i \in \{1, \dots, q\}$, then LICQ is violated in x^* .
- ▶ If there exists an index i such that $H_i(x^*) = 0$ and $G_i(x^*) \geq 0$, then MFCQ is violated in x^* .
- ▶ The vanishing constraints i with $H_i(x^*) > 0$ are “well-behaved”.



How to handle MPVCs?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ One can try to apply NLP theory.

How to handle MPVCs?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ One can try to apply NLP theory.
- ▶ One can use the MPCC reformulation.

How to handle MPVCs?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ One can try to apply NLP theory.
- ▶ One can use the MPCC reformulation.
- ▶ One can define $V := \{(a, b) \in \mathbb{R}^2 \mid b \geq 0, ab \leq 0\}$ and use the condition

$$(\gamma_i, \nu_i) \in N_V^F(G_i(x^*), H_i(x^*)) \subseteq N_V^M(G_i(x^*), H_i(x^*))$$

to obtain possible optimality conditions.

How to handle MPVCs?



- ▶ One can try to apply NLP theory.
- ▶ One can use the MPCC reformulation.
- ▶ One can define $V := \{(a, b) \in \mathbb{R}^2 \mid b \geq 0, ab \leq 0\}$ and use the condition

$$(\gamma_i, \nu_i) \in N_V^F(G_i(x^*), H_i(x^*)) \subseteq N_V^M(G_i(x^*), H_i(x^*))$$

to obtain possible optimality conditions.

- ▶ One can locally decompose the MPVC into TNLP(x^*, I) for all $I \subseteq \{i \mid G_i(x^*) = H_i(x^*) = 0\}$

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0,$$

$$G_i(x) \leq 0, \quad H_i(x) \geq 0 \quad \forall i \in \{i \mid G_i(x^*) < 0\} \cup I,$$

$$H_i(x) = 0 \quad \forall i \in \{i \mid G_i(x^*) > 0\} \cup I^c$$

How to handle MPVCs?



- ▶ One can try to apply NLP theory.
- ▶ One can use the MPCC reformulation.
- ▶ One can define $V := \{(a, b) \in \mathbb{R}^2 \mid b \geq 0, ab \leq 0\}$ and use the condition

$$(\gamma_i, \nu_i) \in N_V^F(G_i(x^*), H_i(x^*)) \subseteq N_V^M(G_i(x^*), H_i(x^*))$$

to obtain possible optimality conditions.

- ▶ One can locally decompose the MPVC into TNLP(x^*, I) for all $I \subseteq \{i \mid G_i(x^*) = H_i(x^*) = 0\}$

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0,$$

$$G_i(x) \leq 0, \quad H_i(x) \geq 0 \quad \forall i \in \{i \mid G_i(x^*) < 0\} \cup I,$$

$$H_i(x) = 0 \quad \forall i \in \{i \mid G_i(x^*) > 0\} \cup I^c$$

- ▶ One can use a nonsmooth reformulation

$$\varphi(a, b) = \max\{-b, \min\{a, b\}\} \leq 0.$$



Cardinality Constraints

Cardinality Constrained Optimization Problems



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ A **cardinality constrained problem** is of the form

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0, \quad \|x\|_0 \leq \kappa$$

with the “sparsity”-term

$$\|x\|_0 := \text{number of nonzero components of } x.$$

- ▶ These problems appear e.g. in
 - ▶ portfolio optimization,
 - ▶ compressed sensing,
 - ▶ feature selection,
 - ▶ communications engineering,
 - ▶ truss design.

Example: Portfolio Optimization



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider a basic portfolio selection problem

$$\begin{aligned} \min_x \quad & x^T Q x \quad \text{s.t.} \quad \mu^T x \geq \rho, \\ & e^T x = 1, \\ & 0 \leq x_i \leq u_i \quad \forall i = 1, \dots, n, \\ & \|x\|_0 \leq \kappa, \end{aligned}$$

where

- ▶ Q is the covariance matrix of the n possible assets,
- ▶ μ is the expected revenue,
- ▶ $e = (1, \dots, 1)^T$.

Relation to Sparse Optimization Problems



TECHNISCHE
UNIVERSITÄT
DARMSTADT

For some weight $\rho > 0$ consider

$$\min_x f(x) + \rho \|x\|_0 \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0.$$

Relation to Sparse Optimization Problems



TECHNISCHE
UNIVERSITÄT
DARMSTADT

For some weight $\rho > 0$ consider

$$\min_x f(x) + \rho \|x\|_0 \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0.$$

- **Sparse \rightarrow Cardinality:** A solution x^* of the sparse problem is always a solution of the cardinality constrained problem with $\kappa := \|x^*\|_0$.

Relation to Sparse Optimization Problems



TECHNISCHE
UNIVERSITÄT
DARMSTADT

For some weight $\rho > 0$ consider

$$\min_x f(x) + \rho \|x\|_0 \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0.$$

- **Sparse \rightarrow Cardinality**: A solution x^* of the sparse problem is always a solution of the cardinality constrained problem with $\kappa := \|x^*\|_0$.
- **Cardinality \nrightarrow Sparse**: Consider the following example by O. Burdakov

$$f(x) = \|Ax - b\|_2^2, \quad \text{where} \quad A = \begin{pmatrix} 0 & 3 & -3 \\ 3 & 2 & 2 \\ 3 & 3 & 3 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 0 \\ 0 \\ 6 \end{pmatrix}.$$

There is no $\rho \geq 0$ for which one obtains the same solution as for $\kappa = 2$.


Continuous Reformulation using Complementarity Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- **Idea:** Introduce **binary variables** y counting the zeros of x :

$$x \circ y = 0, \quad \|x\|_0 \leq n - e^T y \leq \kappa.$$



Continuous Reformulation using Complementarity Constraints




TECHNISCHE
UNIVERSITÄT
DARMSTADT

- **Idea:** Introduce **binary variables** y counting the zeros of x :

$$x \circ y = 0, \quad \|x\|_0 \leq n - e^T y \leq \kappa.$$

Relax the binary variables to $y \in [0, e]$.



Continuous Reformulation using Complementarity Constraints




TECHNISCHE
UNIVERSITÄT
DARMSTADT

- **Idea:** Introduce **binary variables y** counting the zeros of x :

$$x \circ y = 0, \quad \|x\|_0 \leq n - e^T y \leq \kappa.$$

Relax the binary variables to $y \in [0, e]$.

- **Advantages:** No bounds on x needed; relaxation possible; solves the “right” problem.
- **Disadvantages:** Introduces a complementarity-type condition.



Relation between the Original Problem and the Continuous Reformulation



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Exercise

What is the relation between *continuous reformulation*

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & 0 \leq y \leq e, \quad e^T y \geq n - \kappa, \\ & x \circ y = 0. \end{aligned}$$

and the original problem

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0, \quad \|x\|_0 \leq \kappa?$$

Relation between the Original Problem and the Continuous Reformulation (continued)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Global Solutions:

- ▶ x^* is a global solution of the original problem if and only if there exists y^* such that (x^*, y^*) is a global solution of the continuous reformulation.

Relation between the Original Problem and the Continuous Reformulation (continued)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Global Solutions:

- ▶ x^* is a global solution of the original problem if and only if there exists y^* such that (x^*, y^*) is a global solution of the continuous reformulation.

Local Solutions:

- ▶ If x^* is a local solution of the original problem, then for every y such that (x^*, y) is feasible, it is a local solution of the continuous reformulation.

Relation between the Original Problem and the Continuous Reformulation (continued)



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Global Solutions:

- ▶ x^* is a global solution of the original problem if and only if there exists y^* such that (x^*, y^*) is a global solution of the continuous reformulation.

Local Solutions:

- ▶ If x^* is a local solution of the original problem, then for every y such that (x^*, y) is feasible, it is a local solution of the continuous reformulation.
- ▶ If (x^*, y^*) is a local solution of the continuous reformulation and $\|x^*\|_0 = \kappa$, then x^* is a local solution of the original problem.


Example for Spurious Local Solutions



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the 3-dimensional cardinality constrained problem

$$\min_x \|x - (2, 1, 0)^T\|_2^2 \quad \text{s.t.} \quad \|x\|_0 \leq 1.$$



Example for Spurious Local Solutions




TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the 3-dimensional cardinality constrained problem

$$\min_x \|x - (2, 1, 0)^T\|_2^2 \quad \text{s.t.} \quad \|x\|_0 \leq 1.$$

The global minimizer is $x^* = (2, 0, 0)^T$ and the local minimizer $\hat{x} = (0, 1, 0)^T$.



Example for Spurious Local Solutions




TECHNISCHE
UNIVERSITÄT
DARMSTADT

Consider the 3-dimensional cardinality constrained problem

$$\min_x \|x - (2, 1, 0)^T\|_2^2 \quad \text{s.t.} \quad \|x\|_0 \leq 1.$$

The global minimizer is $x^* = (2, 0, 0)^T$ and the local minimizer $\hat{x} = (0, 1, 0)^T$.



However, the continuous relaxation has additional local minimizer such as (\tilde{x}, \tilde{y}) with $\tilde{x} = (0, 0, 0)^T$ and $\tilde{y} = (1, 1, 0)$.


Half and Full Complementarity Formulation



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- The **continuous reformulation** results in

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & 0 \leq y \leq e, \quad e^T y \geq n - \kappa, \\ & x \circ y = 0. \end{aligned}$$




Half and Full Complementarity Formulation



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- The **continuous reformulation** results in

$$\begin{aligned} \min_{x,y} f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & 0 \leq y \leq e, \quad e^T y \geq n - \kappa, \\ & x \circ y = 0. \end{aligned}$$



- The **full complementarity formulation** is given by

$$\begin{aligned} \min_{x^+ - x^-, y} f(x^+ - x^-) \quad \text{s.t.} \quad & g(x^+ - x^-) \leq 0, \quad h(x^+ - x^-) = 0, \\ & y \leq e, \quad e^T y \geq n - \kappa, \\ & x^+, x^- \geq 0, \quad y \geq 0, \quad (x^+ + x^-) \circ y = 0. \end{aligned}$$

Why should you not treat the continuous reformulation as an MPCC?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ In all points of interest MPCC-LICQ and MPCC-MFCQ are violated.

Why should you not treat the continuous reformulation as an MPCC?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ In all points of interest MPCC-LICQ and MPCC-MFCQ are violated.
- ▶ MPCC theory generates conditions on the gradients with respect to y which are either violated or meaningless.

Why should you not treat the continuous reformulation as an MPCC?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ In all points of interest MPCC-LICQ and MPCC-MFCQ are violated.
- ▶ MPCC theory generates conditions on the gradients with respect to y which are either violated or meaningless.
- ▶ The analogues of M-, C- and W-stationarity coincide.

Why should you not treat the continuous reformulation as an MPCC?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ In all points of interest MPCC-LICQ and MPCC-MFCQ are violated.
- ▶ MPCC theory generates conditions on the gradients with respect to y which are either violated or meaningless.
- ▶ The analogues of M-, C- and W-stationarity coincide.
- ▶ All local minima of the continuous reformulation are S-stationary (= KKT points) under a Guignard-type CQ.

Why should you not treat the continuous reformulation as an MPCC?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ In all points of interest MPCC-LICQ and MPCC-MFCQ are violated.
- ▶ MPCC theory generates conditions on the gradients with respect to y which are either violated or meaningless.
- ▶ The analogues of M-, C- and W-stationarity coincide.
- ▶ All local minima of the continuous reformulation are S-stationary (= KKT points) under a Guignard-type CQ.
- ▶ The Scholtes relaxation converges to S-stationary points under an MFCQ-type CQ.



Chance Constraints

Optimization Problems with Chance Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

An **optimization problems with chance constraints** is of the form

$$\begin{aligned} \min_x f(x) \quad \text{s.t.} \quad & g(x) \leq 0, \quad h(x) = 0, \\ & P(G(x, \xi) \leq 0) \geq \eta, \end{aligned}$$

where

- ▶ $\eta \in [0, 1]$ is the level of confidence, e.g. $\eta = 0.9, 0.95, 0.99$,
- ▶ ξ is a random variable with finitely many possible realization ξ_1, \dots, ξ_N and known probabilities p_1, \dots, p_N .

Example: Portfolio Optimization



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Assume that

- ▶ we can invest in n assets,
- ▶ N possible scenarios i can occur with probabilities p_i and prognosed returns $\xi_i \in \mathbb{R}^n$,
- ▶ we want to invest a budget of 1 to maximize the expected return
- ▶ but limit the probability of having losses to $1 - \eta \in (0, 1)$.

We have to solve

$$\max_x \sum_{i=1}^N p_i \xi_i^T x \quad \text{s.t.} \quad x \geq 0, \quad \sum_{i=1}^n x_i = 1, \\ P(\xi^T x \geq 0) \geq \eta$$

Reformulation of Chance Constraints



TECHNISCHE
UNIVERSITÄT
DARMSTADT

The chance constrained problem

$$\min_x f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0, \quad P(G(x, \xi) \leq 0) \geq \eta,$$


can equivalently be rewritten as

$$\min_{x, y} f(x) \quad \text{s.t.} \quad g(x) \leq 0, \quad h(x) = 0,$$

$$y \in \{0, 1\}^N,$$

$$p^T y \geq \eta,$$

$$y_i G(x, \xi_i) \leq 0.$$



Similarly to cardinality constraints, we can relax the binary variables to $y \in [0, 1]^N$.

What should you remember?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- ▶ There are many classes of optimization problems with similar structures/problems as MPCCs.
- ▶ Reformulating them as an MPCC is not always the best approach.
- ▶ But one can use ideas from MPCCs to handle these classes such as
 - ▶ decomposition of the feasible set into easier pieces,
 - ▶ nonsmooth reformulation,
 - ▶ using normal cones to obtain conditions on multipliers.

What did we do this week?



TECHNISCHE
UNIVERSITÄT
DARMSTADT

