

# 04 Proportion Testing

November 20, 2022

```
[1]: import pandas as pd
import numpy as np
import matplotlib.pyplot as plt
from scipy import stats
```

## 1 Bernoulli trial and proportion testing

$$P(S_n = k) = \binom{n}{k} * p^k * (1 - p)^{n-k}, \quad \text{for } 0 \leq k \leq n$$

### 1.1 Hypothesis

$$H_0 : p_0 = p$$

$$H_1 : p > p_0$$

### 1.2 Test statistics

Assume the following things are given:

- $k$ : E. g. number of failed products
- $n$ : Total number of observations
- $p_0$ : Assumed probability that a product fails during production

$$T = \frac{\bar{x} - p}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

where  $\bar{x} = \frac{k}{n}$

```
[2]: k = 46
n = 2000
p0 = 0.02
alpha = 0.01

x_bar = k / n
s_x = p0 * (1 - p0)

# Calculate test statistic
t = (x_bar - p0) / (np.sqrt(s_x / n))
```

```
# Calculate rejection area - right-sided
c = stats.norm().ppf(1 - alpha)

# Calculate the p-value
p_value = 1 - stats.norm().cdf(t)

print(f"Test statistic: {t}")
print(f"Rejection area: {c}")
print(f"P-Value: {p_value}")
print(f"Reject H0: {t > c}")
```

```
Test statistic: 0.9583148474999096
Rejection area: 2.3263478740408408
P-Value: 0.1689520095515027
Reject H0: False
```

```
[ ]:
```