

# Family *Flexiviridae*: A Case Study in Virion and Genome Plasticity

Giovanni P. Martelli,<sup>1</sup> Michael J. Adams,<sup>2</sup>  
Jan F. Kreuze,<sup>3,4</sup> and Valerian V. Dolja<sup>5</sup>

<sup>1</sup>Dipartimento di Protezione delle Piante e Microbiologia Applicata, Università degli Studi and Istituto di Virologia vegetale CNR, Sezione di Bari, 70126 Bari, Italy; email: martelli@agr.uniba.it

<sup>2</sup>Plant-Pathogen Interactions Division, Rothamsted Research, Harpenden, Hertfordshire AL5 2JQ, United Kingdom; email: mike.adams@bbsrc.ac.uk

<sup>3</sup>Germplasm Enhancement and Crop Improvement Division, International Potato Center, Lima 12, Peru and <sup>4</sup>Department of Plant Biology and Forest Genetics, Swedish University of Agricultural Sciences, 750 07 Uppsala, Sweden; email: j.kreuze@cgiar.org

<sup>5</sup>Department of Botany and Plant Pathology and Center for Genome Research and Biocomputing, Oregon State University, Corvallis, Oregon 97331-2902; email: doljav@science.oregonstate.edu

Annu. Rev. Phytopathol. 2007. 45:73–100

First published online as a Review in Advance on  
March 15, 2007

The *Annual Review of Phytopathology* is online at  
phyto.annualreviews.org

This article's doi:  
10.1146/annurev.phyto.45.062806.094401

Copyright © 2007 by Annual Reviews.  
All rights reserved

0066-4286/07/0908-0073\$20.00

## Key Words

positive-strand RNA viruses, comparative genomics, phylogeny, evolution, taxonomy

## Abstract

The plant virus family *Flexiviridae* includes the definitive genera *Potexvirus*, *Mandarivirus*, *Allexivirus*, *Carlavirus*, *Foveavirus*, *Capillovirus*, *Vitivirus*, *Trichovirus*, the putative genus *Citrivirus*, and some unassigned species. Its establishment was based on similarities in virion morphology, common features in genome type and organization, and strong phylogenetic relationships between replicational and structural proteins. In this review, we provide a brief account of the main biological and molecular properties of the members of the family, with special emphasis on the relationships within and among the genera. In phylogenetic analyses the potexvirus-like replicases were more closely related to tymoviruses than to carlaviruses. We postulate a common evolutionary ancestor for the family *Tymoviridae* and the two distinct evolutionary clusters of the *Flexiviridae*, i.e., a plant virus with a polyadenylated genome, filamentous virions, and a triple gene block of movement proteins. Subsequent recombination and gene loss would then have generated a very diverse group of plant and fungal viruses.

## INTRODUCTION

Historically, all organisms including viruses were classified based primarily on their morphology and reproductive cycle. Later, the rise of molecular biology and comparative genomics revolutionized understanding of virus taxonomy and evolution, and revealed unexpected relationships between apparently dissimilar viruses, as well as surprising genetic differences between viruses that look alike (3, 43). The recently established family of plant-infecting positive-strand RNA viruses, *Flexiviridae* (2), highlights both of these seemingly opposing tendencies. As the family name immediately suggests, its members share the flexuous morphology of their elongated helical virions. A similar morphology is found in two other plant virus families, *Potyviridae* (115) and *Closteroviridae* (31, 78). Earlier phylogenetic analysis demonstrated that capsid proteins forming filamentous virions of plant viruses are related to each other and likely originate from a single ancestral protein (29). However, whole genome comparisons showed that the virions of the three families harbor strikingly distinct genomes of the alphavirus-like (*Closteroviridae* and *Flexiviridae*) and picornavirus-like (*Potyviridae*) superfamilies (60).

The principal basis for the establishment of the family *Flexiviridae* was the strong phylogenetic relationships between the viral replicational and structural proteins and common features of genome design. All members of the family share monopartite, 3'-polyadenylated genomes that encode closely related methyltransferase, RNA helicase, and RNA polymerase domains of the viral replicase. In contrast, one or two proteases, and AlkB-homology domains are found in replicases of some, but not all, family members. There is also a remarkable diversity in the number and nature of the 3'-proximal viral genes that are expressed via formation of subgenomic mRNAs. These genes encode viral movement proteins that belong to either the "p30-like" superfamily, or to the triple gene block

movement complex (69, 91), the nucleic acid-binding proteins, and several unique proteins found in a limited number of family members. These genetic differences translate into a variety of host ranges, pathogenicity levels, and overall epidemiology exhibited by the viruses within the family *Flexiviridae*.

The currently recognized family genera are *Allexivirus*, *Capillovirus*, *Carlavirus*, *Foveavirus*, *Mandarivirus*, *Potexvirus*, *Trichovirus*, and *Vitivirus*. An additional genus, *Citricivirus* (121), with *Citrus leaf blotch virus* (CLBV) as the type species, is under scrutiny for approval by the International Committee on Taxonomy of Viruses. The unassigned family members include *Banana mild mosaic virus* (BanMMV), *Cherry green ring mottle virus* (CGRMV), *Cherry necrotic rusty mottle virus* (CNRMV), *Potato virus T* (PVT), and *Sugarcane striate mosaic-associated virus* (SCSMaV) (2). The recently described Banana virus X (BanVX) is another putative member of the family that does not seem to fit in any of the extant genera (113), whereas African oil palm ringspot virus (AOPRV) (88) and Asian prunus virus-1 (APruV-1), -2 (APruV-2), and -3 (APruV-3) (75) are additional putative members of the genus *Foveavirus*.

In this review, we provide a brief account of the main biological and molecular properties of the *Flexiviridae* with a special emphasis on the relationships within and among the component genera. Using detailed phylogenetic analysis in relation to the recent progress in understanding the basic pathways of virus-host coevolution, we attempt to develop a coherent and parsimonious scenario for the origin and evolution of this viral family.

## BIOLOGICAL PROPERTIES

### Host Range and Pathogenicity

In natural and agricultural settings, flexiviruses infect a wide variety of wild and cultivated hosts including herbaceous

and woody dicotyledonous and, less often, monocotyledonous plant species. Most of the viruses are pathogenic to agricultural crops, their infections generating mild to extremely severe diseases. Most members of the genera *Capillovirus*, *Citrivirus*, *Foveavirus*, *Mandarinivirus*, *Trichovirus*, and *Vitivirus* and the unassigned species CGRMV and CNRMV are agents of woody host diseases, whereas the genera *Potexvirus*, *Carlavirus*, and *Allexivirus* and the remaining unassigned species in the family tend to infect herbaceous plants.

Several capilloviruses, foveaviruses, and vitiviruses share the capacity to elicit modifications of the host xylem known as stem pitting or grooving, depending on whether the woody cylinder is marked by localized, shallow surface indentations (pits), or by long narrow depressions (grooves). An emblematic example of this type of disorder is the rugose wood complex of grapevine (76) in the etiology of which the vitiviruses *Grapevine virus A* (GVA) and *Grapevine virus B* (GVB), and the foveavirus *Grapevine rupestris stem pitting-associated virus* (GRSPaV) are involved, as the putative agents of Kober stem grooving (22, 41) corky bark (11), and *Rupestris* stem pitting (83), respectively. Another vitivirus, *Grapevine virus D* (GVD), is associated with an espe-

cially serious condition of certain grapevine cultivars known as corky rugose wood (10) (**Figure 1**).


Plants infected with GVA, GVB, and GRSPaV often harbor a population of viral sequence variants (45, 46, 83, 107), some of which endowed with differential pathogenicity. Thus, the comparative analysis of the 3'-terminal regions of a number of GVA isolates allowed their separation into three molecular groups that share sequence identity within groups above 90% and below this value between the groups. Only isolates of group III induced mild symptoms in the experimental host *Nicotiana benthamiana* (45). These isolates appeared not to induce in grapevine cv Shiraz a serious disorder known in South Africa as Shiraz disease (SD), whereas a variant of group I dominated in SD-affected vines (46). By contrast, one of the three sequenced GRSPaV variants was shown to infect *Vitis rupestris* without inducing symptoms (84).

*Apple stem grooving virus* (ASGV), a capillovirus, and *Apple stem pitting virus* (ASPV), a foveavirus, are elicitors of wood abnormalities (Plate 1) in some apple and pear cultivars, especially from Far Eastern countries, and may cause crippling graft incompatibility such as the Japanese apple topworking disease (27, 55). The citrus strain (tatter leaf)

**GVA:** *Grapevine virus A*

**GVB:** *Grapevine virus B*

**GVD:** *Grapevine virus D*



**Figure 1**

(a) Field-grown vine affected by rugose wood; (b) stem grooving induced on Kober 5BB by *Grapevine virus A*; (c) stem pitting induced on Virginia Crab by *Apple stem grooving virus*; (d) mild stem pitting induced on *Malus platicarpa* by *Apple stem pitting virus*. (c and d courtesy of Dr. L. Giunchedi)

**PVX:** *Potato virus X*

**PVY:** *Potato virus Y*

**PVS:** *Potato virus S*

of ASPV is strongly pathogenic to grafted trifoliolate rootstocks, which react with bud union crease and stem grooving (101). As its name suggests, the tentative capillovirus *Nandina stem pitting virus* (NSPV) is associated with a stem pitting in *Nandina domestica* (heavenly bamboo) (4), and the unassigned CGRMV and CNRMV with a wood pitting of cherry (27, 126).

*Apricot latent virus* (ApLV), a foveavirus inducing symptomless infections in most apricot cultivars (94), possesses two molecular variants pathogenic to peach, in which they cause foliar diseases called peach asteroid mosaic and peach sooty ringspot, respectively (27, 42). Similar symptoms of chlorotic leaf spotting and fruit deformation are associated with peach infection by Asian prunus viruses (75), while AOPRV is held responsible for a lethal disease of African oil palms (88). None of the putative flexiviruses infecting banana (BanMMV and BanVX) seems to be a serious pathogen (40, 113).

Whereas infection by CLBV generates bud union crease in plants grafted on trifoliolate rootstocks (39), *Indian citrus ringspot virus* (ICRSV), the only known member of the genus *Mandarivirus*, induces foliar yellow ringspots and a rapid decline of affected trees (18).

Trichoviruses are mainly pathogens of stone fruit trees (apricot, cherry, almond, peach, and plum) and of grapevine (*Grapevine berry inner necrosis virus*, GINV). In these hosts, trichoviruses elicit a variety of symptoms, ranging from delayed bud break, stunted growth, mottling and deformation of the leaves (Peach mosaic virus, PcMV; Cherry mottle leaf virus, ChMLV), to severe damage of fruits, i.e., false plum pox, plum bark split, fruit necrosis of cherry and apricot (*Apple chlorotic leafspot virus*, ACLSV), and necrosis of grape berries (GINV) (27, 67, 111).

Viruses of the genera *Allexivirus*, *Carlavirus*, and *Potexvirus* are predominantly pathogens of herbaceous plants. The genus *Potexvirus* comprises species with important economic impact. In an inquiry

coordinated by Milne (86), the potexvirus *Potato virus X* (PVX) was among the top 10 most damaging viruses in seven of nine wide geographical areas including North America, South America, Europe, India, China, Africa, and Australia/New Zealand. However, although infection of solanaceous crops such as potato, tomato, pepper, and tobacco by PVX alone could result in diseases of some consequence, effects were much worse when this virus was in mixture with others [*Potato virus Y* (PVY), in particular]. In the same inquiry, the carlavirus *Potato virus S* (PVS) was reported as a pathogen of concern only in South America, in line with the notion that carlavirus infections are frequently symptomless. Symptoms, when shown, are usually of the mosaic type, much the same as with potexviruses (33, 100, 123). The host ranges of allexiviruses are limited to *Allium* species, in which they are symptomless or induce mild diseases (6).

There is only limited information on the molecular determinants of flexivirus pathogenicity. Two examples include the 25-kDa protein encoded by ORF2 of PVX and the 10-kDa protein encoded by ORF5 of GVA, both of which are involved in symptom development and suppression of RNA silencing (14, 20, 37, 50, 127).

## Epidemiology

With a few exceptions, members of the family *Flexiviridae* are transmissible from natural to experimental hosts by mechanical inoculation. Viruses that invade and multiply in parenchymatous tissues, especially members of the genera *Potexvirus*, *Carlavirus*, *Allexivirus*, and *Trichovirus*, are more readily transmissible in this way than vitiviruses, which are phloem-restricted in the natural host, or foveaviruses, which are transmitted with difficulty (ASPV and ApLV) or not at all (GRSPaV).

In vegetatively propagated hosts, virus dissemination over long distances takes place primarily through the distribution of infected

propagative materials. Thus, members of the family that do not have a vector rely exclusively on graft-transmission for their survival and dissemination. All flexiviruses infecting woody hosts, regardless of the genus, are efficiently spread by nursery productions, which are largely responsible for the worldwide deterioration of the sanitary status of fruit tree crops. Infected bulbs, tubers, offshoots, and the like are similarly responsible for the dispersal of potex-, carla-, and allexiviruses (48). An exception is *Lilac chlorotic leafspot virus* (LiCLSV), which, although being transmitted in a semipersistent manner by aphids (15, 16), is currently classified as a capillovirus (2). However, the taxonomic allocation of LiCLSV is under scrutiny and removal from the list of approved species is being considered.

Although seeds represent important natural routes for virus dissemination, seed transmission is hardly relevant to the economy of the family *Flexiviridae*. Examples are few, represented by a capillovirus (the citrus strain of ASGV), three potexviruses [PVX, *Clover yellow mosaic virus* (CIYMV), and *White clover mosaic virus* (WCIMV)], and four carlaviruses [*Hop mosaic virus* (HpMV), *Pea streak virus* (PeSV), *Red clover vein mosaic virus* (RCVMV), and some isolates of *Cowpea mild mottle virus* (CPMMV)]. Transmission rates of individual viruses are generally low and do not exceed 10% (81, 87).

Flexivirus infection is introduced at a site by mechanical contact or from the activity of vectors. The first of these modes of transmission is, by and large, typical of potexviruses, which do not have vectors, are very stable, extremely infectious, and multiply to high titers in their hosts. Contact between an infected and a healthy plant is thought to be sufficient for propagating the infection (100). Some carlaviruses (e.g., PVS) behave in a similar way (59), although their natural transmission occurs by aphids in a nonpersistent manner (123). An exception is CPMMV, which is the only known carlavirus transmitted by the whitefly *Bemisia tabaci* (17). The molecu-

lar mechanism underlying aphid transmission of carlaviruses has not been elucidated.

Most members of the genera *Allexivirus* and at least three of the genus *Trichovirus* are mite-borne; the vectors in all cases are eriophyd mites (97). *Aceria tulipae* is the alleged vector of the allexiviruses *Garlic virus C* (GarV-C), *Garlic virus D* (GarV-D), and *Shallot virus X* (ShVX) and the tentative species *Garlic mite-borne latent virus* (GarMbLV), *Shallot mite-borne latent virus* (ShMbLV), and *Onion mite-borne latent virus* (OmbLV) (116, 117).

Among trichoviruses, PcMV is transmitted by *Eriophyes insidiosus* (58), ChMLV by *Eriophyes inequalis* (97), and GINV by *Colomerus vitis* (64). The mode of transmission of these viruses is still undetermined. Notwithstanding the lack of definitive experimental evidence, Oldfield & Proeseler (97) suggested that transmission of eriophyd-borne viruses of *Allium* species and woody perennials is of the persistent type.

Vitiviruses are the only members of the family that are vectored by mealybugs. GVA was the first RNA virus shown to be transmissible by *Pseudococcus longispinus*, a pseudococcid mealybug (102) and, ten years later, transmission of GVB by *Planococcus ficus* was obtained (11). Although GVA and GVB remain the only mealybug-transmissible vitiviruses, the number of vectors has grown to six different species in the genera *Planococcus*, *Pseudococcus*, and *Helicoccus* and one species in the soft scale insect genus *Neopukvinaria* (80). As experimentally determined with *Ps. longispinus*, transmission of GVA is semipersistent (66).

## Relations with Cells and Tissues

Modifications induced in host cells by flexiviruses have been studied either in the natural hosts for potexviruses, carlaviruses, and allexiviruses, or in experimentally infected herbaceous indicators for capilloviruses, foveaviruses, trichoviruses, and vitiviruses (23, 34, 70, 77, 96).



In general, flexivirus infections do not give rise to cytopathic structures (e.g., inclusion bodies) specific enough to be diagnostic at the genus level (77). Rather, it seems that cytoplasmic and, sometimes, nuclear accumulations of virions in various forms are the most frequent, if not the only intracellular trait common to all members of the family. Differences, however, exist in the outward appearance of these accumulations. Thus, potex-, viti-, and capillovirus aggregates have a flexuous appearance, whereas those of carla-, tricho-, and foveaviruses are rather stiff (34, 70, 96). The size of the aggregates can vary, with PVX accumulations being always massive (70, 77). Large, bundle-like viral aggregates are also seen in cells infected by vitiviruses and carlaviruses, whereas accumulations of allexi-, capillo-, and trichoviruses are relatively small (34, 96).

Species-specific cytopathic structures are elicited by some potex- and carlaviruses. Examples include (i) PVX-induced “laminated inclusions”; (ii) amorphous inclusions made up of excess coat protein in the cells infected by CIYMV; (iii) striking crystalline cytoplasmic and nuclear inclusions associated with Argentine plantago virus and Boussingaultia mosaic virus (70). As to carlaviruses, the whitefly-transmitted CPMV was reported to elicit unusual brush-like particle aggregates in the infected cells (70).

The cytopathological features common to vitiviruses are more extensive and consist of

(i) virion aggregates in the shape of bundles, whorls, banded bodies, or stacked layers; (ii) cell wall thickenings; (iii) proliferation and accumulation of endomembranes; (iv) vesicular evaginations of the tonoplast protruding into the vacuole (34, 80). Movement proteins of GVA and GVB were identified by gold immunolabelling in association with cell walls and plasmodesmata (105).


## VIRION STRUCTURE

Particles of flexiviruses are formed by a single type of capsid protein subunit with a mol. wt. ranging from 21 kDa in GVD to 41 kDa in CLBv, and encapsidate a single RNA molecule of ~6 to ~9 kb (**Table 1**) constituting ~5% of the particle weight (2). Studies of the two other families of plant-infecting filamentous viruses, closteroviruses (99, 106) and potyviruses (112), revealed that their virions possess terminal tail-like structures formed by additional viral proteins. Because these proteins were also implicated in virus transport, the tails were proposed to represent specialized movement devices (28). The ability of PVX RNA and virions to associate with the viral 25-kDa movement protein *in vitro* (57) suggests that the virions of flexiviruses may also possess the tail-like appendages required for virus transport in infected plants.

Filamentous virions of flexiviruses have modal lengths ranging from ~470 nm in potexviruses to ~1000 nm in citriviruses, a

**Table 1** Properties of the family *Flexiviridae* genera

Genus	Virion length (nm)	Pitch of the helix (nm)	Genome size (kb)	CP (kDa)	ORFs	MP type	Rep (kDa)
<i>Potexvirus</i>	470–580	3.3–3.6	5.9–7.0	22–27	5	TGB	146–191
<i>Mandarivirus</i>	650	~3.4	7.5	36	6	TGB	187
<i>Allexivirus</i>	~800	~3.2	8.1–8.8	26–28	6	TGB	175–194
<i>Carlavirus</i>	610–700	~3.4	7.4–8.5	31–40	6	TGB	226–238
<i>Foveavirus</i>	723–800+	3.6–4.0	8.7–9.3	28–45	5 or 6	TGB	244–247
<i>Capillovirus</i>	640–700	3.4–3.8	6.5–7.4	25–27	2 or 3	30 K	214–241
<i>Vitivirus</i>	725–785	3.4–3.6	7.4–7.6	21–23	5	30 K	195–196
<i>Trichovirus</i>	640–760	3.8	7.5–8.1	21–22	3 or 4	30 K	216–217
<i>Citrivirus</i>	960	UD	8.7	41	3	30 K	227



**Figure 2**

Virus particles of some genera of the family *Flexiviridae*. (a) *Potato virus X* (*Potexvirus*). (b) *Grapevine virus B* (*Vitivirus*). (c) *Apple chlorotic leaf spot virus* (*Trichovirus*). (d) *Apple stem pitting virus* (*Foveavirus*). (e) *Carnation latent virus* (*Carlavirus*). (f) *Indian citrus ringspot virus* (*Mandarivirus*). Bars = 50 nm. (a, e, f courtesy of Dr. R.G. Milne; b, c, courtesy of Dr. M.A. Castellano; d, courtesy of Dr. A. De Stradis.)

diameter of 12–14 nm (**Table 1**) and, when discernible, show an axial canal of ~1.5 nm in diameter. There are 8.9 subunits per turn of the helix in PVX virions (98, 108, 114). Depending on the genus, flexiviruses exhibit varying degrees of flexibility, with potexviruses and carlaviruses being the least flexuous. In fact, the latter are semirigid and show a tendency to curve to one side (85). Citri- and capilloviruses have an intermediate flexibility, whereas virions of all remaining genera are extremely flexuous, with an open structure readily visible in the electron microscope (**Figure 2**).


The surface of carlavirus and potexvirus particles is marked by intersecting sets of distinct grooves (85), one of which runs along the particle length and the other follows the viral helix (98). The particles of all family members

exhibit distinct cross-banding. This feature is outstanding in the species of the very flexuous genera and is less evident in potexviruses and carlaviruses (85). Cross-banding spacing corresponds to the pitch of the primary helix (12, 114). Carlavirus and potexvirus particles are marked also by near-longitudinal lines along their length that correspond to rows of subunits lying on secondary helices (85, 114).

## GENOME ORGANIZATION AND EXPRESSION

Genome organization of the typical *Flexiviridae* family members is summarized in **Figure 3**. Although the number of ORFs (open reading frames) varies between genera and sometimes between species, the first and largest ORF is always a replicase gene.

**ORF:** open reading frame



**Figure 3**

Diagram showing genome organization for the genera of the family *Flexiviridae*. Blocks represent predicted open reading frames (ORFs). The methyltransferase (M), AlkB (A), OTu-like peptidase (O), papain-like protease (P), RNA helicase (H), and RNA-dependent RNA polymerase (R) domains of the replicase are shown. RNA-binding proteins are present in some, but not all, members of the genus *Trichovirus*.

This is likely the only ORF translated directly from the capped genomic RNA, although it was suggested for GVA that a shorter-than-genomic RNA could also be involved in replicase expression (36). The remaining ORFs are expressed from 3'-coterminal subgenomic (sg) mRNAs, some of which could be bicistronic. At least in some potexviruses, sgRNAs are packaged (7).

### Replicase ORF

Although the size of the replicase protein varies between the family members from ~150 to ~250 kDa, this protein invariably contains a set of functional domains whose

amino acid sequences and order are conserved in all viruses of the alphavirus-like superfamily of the positive-strand RNA viruses (60). A methyltransferase type 1 domain (Met; **Figure 3**) is located near the N terminus of the replicase (104). This domain functions in capping the viral RNAs and in the case of *Bamboo mosaic virus* (BaMV, genus *Potexvirus*) was shown to exhibit both GTP methyltransferase and guanylyltransferase activities (53). A catalytic RNA-dependent RNA polymerase domain (RdRp; **Figure 3**) with a characteristic core motif S/TG<sub>x3</sub>TX<sub>3</sub>NS/TX<sub>22</sub>GDD is present near the C terminus of the replicase. It was shown that the BaMV RdRp is capable of recognizing the 3'-terminal

**sg:** subgenomic

**RdRp:**  
RNA-dependent  
RNA polymerase  
domain



pseudoknot structure to initiate minus-strand RNA synthesis (21). An RNA helicase domain of superfamily 1 (Hel; **Figure 3**) is localized upstream from the RdRp domain (44). Similar to Met and RdRp, Hel is indispensable for viral RNA replication, as demonstrated for PVX (26). Besides its presumed function in unwinding RNA, BaMV Hel was also implicated in cap formation (71).

In addition to the Met-Hel-RdRp module, replicases of the animal viruses of the alphavirus-like superfamily universally possess a papain-like cysteine protease domain (P-Pro) that processes the replicase polyprotein (60). The distribution of P-Pro among plant alphavirus-like viruses is more limited. In the family *Flexiviridae*, P-Pro is located upstream from Hel in carla-, capillo-, fovea-, citri-, and trichoviruses, and in BanMMV, CGRMV, CNRMV and SCSMaV, as was first demonstrated by (68) for *Blueberry scorch virus* (BlScV), genus *Carlavirus* (**Figure 3**). In BlScV, the autocatalytic processing by P-Pro appears to be required for RNA replication. In potexviruses and other genera lacking P-Pro, the replicase polyprotein functions in a nonprocessed form (7).

Subsequently, a second proteinase domain located immediately upstream from the papain-like protease was recognized in a subset of flexiviruses (13, 74). This domain belongs to a new superfamily of the proteases distantly related to papain and typified by the Ovarian Tumor (OTU) gene involved in oocyte morphogenesis in *Drosophila*. In addition to eukaryotic organisms, OTU proteases are found in several pathogens including diverse DNA, negative-strand RNA, and positive-strand RNA viruses where they were suggested to play a significant role in virus-host interactions, possibly involving the ubiquitin-dependent pathway (74). Our search of the PFAM database confirmed the presence of OTU domains in most carla-, fovea-, and fovea-like viruses (J.K., unpublished data). The exact function of this predicted OTU protease in the infection cycles of

the flexiviruses presents an interesting problem for future research.

Another unexpected recent finding was the identification of the AlkB-like domain in the replicase polyproteins of several flexiviruses and closteroviruses (5, 13). This domain is located downstream from the Met, similar to OTU, and its function in virus reproduction is currently unknown. Although the AlkB domain was identified in most of the *Flexiviridae* genera, it exhibits nontrivial distribution among the viruses within genera (see below).

## ORFs Encoding Movement Proteins

Downstream of the replicase ORF, flexiviruses encode one or more proteins involved in viral cell-to-cell movement. Five of the family genera, *Allexivirus*, *Carlavirus*, *Foveavirus*, *Mandavivirus*, and *Potexvirus*, as well as BanMMV, CGRMV, CNRMV, and SCSMaV, possess a set of three partially overlapping ORFs known as a triple gene block (TGB) (89). Most likely, these ORFs are expressed from two species of the 3'-coterminal sgRNAs (90, 118). A TGB module is also found in members of the genera *Benyvirus*, *Hordeivirus*, *Pechivirus*, and *Pomovirus*, alphavirus-like plant viruses with rod-shaped virions. However, the first and third proteins of the TGB of these viruses are considerably larger than the corresponding proteins in the family *Flexiviridae* (91). Functions of the TGB proteins were recently reviewed and are not discussed in detail here (see 91, 119). In brief, a ~25-kDa TGBp1 has ATPase, RNA-binding and RNA-helicase activities, and is believed to increase the size exclusion limit of plasmodesmata. The smaller proteins TGBp2 and TGBp3 have, respectively, two and one transmembrane domains, and localize to endomembranes and cell walls. The three proteins are believed to act in concert to enable cell-to-cell movement of the viral genomes. Although the exact mechanism by which the movement occurs remains unknown, it was found that TGBp1 of PVX possesses RNA silencing suppressor activity (122). Moreover, it was shown that this activity

---

**OTU:** Ovarian Tumor

**TGB:** triple gene block

---

**TMV:** Tobacco mosaic virus

**CP:** capsid protein

is essential, but not sufficient, for PVX movement (8).

The *Flexiviridae* genera *Capillivirus*, *Citricivirus*, *Trichovirus* and *Vitivirus* have a single movement protein that is a member of the 'p30-like' superfamily typified by the 30-kDa movement protein of *Tobacco mosaic virus* (TMV) (69, 82, 92). The p30-like movement proteins occur in a large number of diverse plant viruses with RNA and DNA genomes. Mutational study confirmed the role of the p30-like protein in the cell-to-cell movement of GVA (38, 105). It was also found recently that the 50-kDa movement protein of ACLSV suppresses systemic, but not local, RNA silencing (125). Therefore, the functional profile of this protein overlaps that of the unrelated PVX movement protein, TGBp1, emphasizing the role of silencing suppression in virus transport and mechanistic convergence among the two classes of the flexiviral movement proteins.

### 3'-Proximal ORFs Encoding Capsid and Other Proteins

The ORF coding for the capsid protein (CP) is either the 3'-most or penultimate ORF in the flexiviral genomes (**Figure 3**). Typically, CP is expressed from an abundant, functionally monocistronic sgRNA (30, 38). In the genus *Capillivirus*, the CP ORF is contiguous with the replicase ORF, but analysis of the RNA patterns suggested that the CP may be expressed from a sgRNA (73).

In the genera *Allexivirus*, *Carlavirus*, *Man-darivirus* and *Vitivirus*, the 3'-most ORFs encode RNA-binding proteins with a zinc ribbon motif (20, 37, 47). Similar proteins are found in some, but not all, viruses in the genus *Trichovirus* (54). It was recently shown that the nucleic acid-binding protein of GVA is able to suppress RNA silencing (127), suggesting that this function may be conserved in other related proteins of flexiviruses.

Viruses in the genera *Allexivirus* and *Vitivirus* possess additional ORFs encoding unique proteins of unknown functions

(**Figure 3**). Among these, the 42-kDa product of the allexivirus ORF4 was implicated in virion assembly (120), while the 19-kDa product of the vitivirus ORF2 was suggested to aid virus transmission by the mealybugs (38).

## EVOLUTIONARY HISTORY

The availability of entire genome sequences for a large number of viruses and their host organisms has enabled large-scale phylogenetic analyses and provided unprecedented opportunities for data-informed reconstructions of viral origins and evolution in relation to evolution of cellular life forms (24, 35, 61). It is now widely accepted that the positive-strand RNA viruses share a universally conserved RdRp and represent an ancient, monophyletic, evolutionary lineage (3, 56, 60). The absence of viral RdRp homologs among cellular organisms was recently interpreted to support its direct origin from a precellular RNA-protein world (62). Furthermore, it was argued that the stark contrast between the scarcity of the prokaryote-infecting positive-strand RNA viruses and the enormous abundance of such viruses in eukaryotes suggests that the diverse lineages of the extant positive-strand RNA viruses have evolved at the time of eukaryogenesis (62). This notion finds increasing support in the discovery of marine positive-strand RNA viruses that infect evolutionary distant unicellular eukaryotes (25, 65, 93, 110). Indeed, the fact that these marine viruses belong to picornavirus-like and flavivirus-like superfamilies indicates that these superfamilies have evolved in early eukaryotes. Much less plausible is an alternative scenario that would call for massive introduction of viruses that infect land plants and animals into marine plankton communities. Alphavirus-like viruses of unicellular eukaryotes are yet to be found. Because the deep evolutionary roots of the alphavirus-like superfamily cannot thus be revealed, we focused on the later phase in evolution that yielded current genomic diversity within the family *Flexiviridae*. The evolutionary scenario

for flexiviruses was deduced using extensive phylogenetic analysis of the viral replicases, CPs, and movement proteins.

## Phylogenetic Analyses of the Viral Proteins


**The replicase.** Because RdRp is the only protein universally encoded in the nondefective positive-strand RNA viruses, its phylogeny is the principal determinant of the evolutionary framework in this vast virus class. In addition, the Met and Hel domains are conserved throughout the alphavirus-like superfamily (60), therefore facilitating further refinement of the viral phylogenies. In agreement with the original finding (103), our expanded analysis of the three replication-related domains revealed that within the alphavirus-like superfamily, families *Flexiviridae* and *Tymoviridae* are most closely related to each other to the exclusion of other plant virus families such as *Closteroviridae* and *Bromoviridae* and unassigned genera such as *Tobamovirus* and *Benyvirus* (Figure 4a). Moreover, our analysis shows that the *Flexiviridae* can be divided into two clusters, one containing potex-, alllexi-, and mandariviruses (collectively called here potexvirus-like viruses), and the other containing carla-, viti-, capillo-, fovea-, and trichoviruses (carlavirus-like viruses). Surprisingly, the replicases of the potexvirus-like viruses are more closely related to those of the *Tymoviridae* than to the carlavirus-like replicases (Figure 4b).

In addition to plant flexi- and tymoviruses, this evolutionary lineage of the viral replicases includes three fungal viruses, *Botrytis cinerea* virus F (BCV-F), *Botrytis* virus X (BVX), and *Sclerotinia sclerotiorum* debilitation-associated RNA virus (SsDRV); two flexuous filamentous viruses; and a capsid-less virus, respectively (51, 52, 124). Conspicuously, our analysis of the complete replicase sequences (Figure 4b), as well as separate Hel domains (not shown), suggested that the filamentous BCV-F is related more closely to the *Tymoviridae* than to any of the *Flexiviridae* clus-

ters. This conclusion is also supported by the similarity of the papain-like proteinases found in BCV-F and tymoviruses that both belong to the C21 PFAM family of endopeptidases (results of HMMER search of PFAM database performed in this study), unlike proteases of carlavirus-like viruses that are of C23 or C34 PFAM families (potexvirus-like viruses lack proteases altogether). In contrast, BVX shows highest similarity to allexiviruses, whereas SsDRV is a peripheral member of the potexvirus-like cluster (Figure 4). Such remarkable evolutionary affinities between the fungal and plant RNA viruses are not without a precedent: The capsid-less viruses of the family *Hypoviridae* are the evolutionary kin of the plant family *Potyviridae* (95). Collectively, these observations suggest the amazing propensity of the filamentous plant viruses to cross the host kingdom barriers and spawn their evolutionary descendants to plant-pathogenic fungi. This direction of transfer seems to be supported by the higher diversity of relevant plant viruses and affinity of their fungal cousins to particular plant virus lineages.

**The capsid proteins.** As noted above, CPs of the flexiviruses share the conserved structural core and evolutionary origin with CPs of two other families of filamentous plant viruses, *Closteroviridae* and *Potyviridae* (19, 29). A phylogenetic tree of the flexiviral core CP sequences reveals two clusters with good bootstrap support (Figure 5a). The first and largest cluster contains potex-, alllexi-, mandari-, carla-, fovea-, and citriviruses, as well as the filamentous fungal virus BVX. The second cluster includes capillo-, tricho-, and vitiviruses along with another fungal virus, BCV-F. However, even though both the replicase tree (Figure 4) and the CP tree (Figure 5a) contain two major clusters each, the sets of the genera that fall into these clusters are distinct.

**The movement proteins.** As discussed above, flexiviruses possess two unrelated types





of movement proteins that belong to either the TGB complex or the p30-like superfamily. In accord with previous work (reviewed in 91), our phylogenetic analysis shows that flexivirus TGB proteins represent a compact cluster distinct from their evolutionary counterparts in the rod-shaped plant viruses of the alphavirus-like superfamily (**Figure 5b**). Therefore, the monophyletic origin of the flexiviral TGB proteins seems to be well supported.

The p30-like movement proteins of plant viruses are generally quite variable (92). Due to this and the availability of fewer sequences, support for the monophyletic relationship of the flexiviral p30-like proteins is not very robust beyond the genus level, yet sufficient to claim their common origin (data not shown; see also 82).

## AN EVOLUTIONARY SCENARIO FOR THE FLEXIVIRIDAE/TYMOVIRIDAE LINEAGE

We used comparisons of the phylogenetic trees for viral proteins and the genome designs of the extant viruses to reconstruct a hypothetical common ancestor and to develop the most parsimonious scenario for the diversification of the family *Flexiviridae*. The former task presents an immediate challenge due to the undisputable clustering of the potexvirus-like replicases with those of tymoviruses to the exclusion of carlaviruses-


like replicases (**Figure 4**). It appears that the most sensible way of solving this paradox is recognizing that the family *Tymoviridae* and two related, but distinct, evolutionary clusters comprising the family *Flexiviridae* shared a common evolutionary ancestor.

In an attempt to reconstruct an ancestor of the *Flexi-/Tymoviridae* lineage, we postulated that it was a plant virus encoding a minimal set of proteins required for RNA replication, encapsidation, and virus movement from cell to cell. Because all flexi- and tymoviruses possess closely related Met, Hel, and Pol domains, these domains must have been present in the ancestor's replicase (**Figure 6**). The presence of a Met domain implies that the ancestor's genome was 5'-capped. Because the genomes of all but one extant genus in this lineage are polyadenylated, the common ancestor was likely to possess a poly(A) tail that was replaced by a tRNA-like structure in the ancestor of the genus *Tymovirus* (32).

It also appears likely that the ancestral replicase harbored a P-Pro domain. Indeed, this domain is present in all tymoviruses and flexiviruses with carlavirus-like replicase with a single exception of the genus *Vitivirus* (**Figure 4**). Moreover, related P-Pro is universally conserved in the animal viruses of the alphavirus-like superfamily. The simplest explanation of such a distribution pattern is that P-Pro was present in a common ancestor of


**Figure 4**

Phylogenetic trees of replicase amino acid sequences. Genera within the *Flexiviridae* are indicated in different colors. (a) Tree created from the fused alignments of methyl-transferase, helicase, and polymerase conserved core domains of plant infecting "Alpha"-like viruses. Deletion of the variable regions between the conserved domains was required to obtain a robust alignment; (b) tree created from the alignment of the entire replicase sequence of flexiviruses, tymoviruses, and related viruses. Alignments and phylogenetic analysis were performed using the Mega 3.1 package (63). Presented trees were created by neighbor joining using the JTT matrix and pair-wise gap deletion. Numbers on branches indicate percentage of bootstrap support out of 1000 bootstrap replications. Branches with less than 70% bootstrap support were collapsed. Similar phylogenies were obtained using the Maximum Parsimony method, and the corresponding bootstrap values are indicated between brackets in branches above the genus level. Scales indicate JTT amino acid distances. Distantly related genera and families that formed well-supported monophyletic clades were collapsed into a triangle, the size of which corresponds to the variation found within the clade. Accession numbers of sequences used for constructing the trees are in **Table 2**.



**Figure 5**

Phylogenetic analyses of the flexivirus protein sequences: (a) capsid protein; (b) triple gene block protein 1 helicase domain. Colors and methods are as in **Figure 4**. Accession numbers of sequences used for constructing the trees are in **Table 2**.



**Figure 6**

Evolutionary scenario for the families *Flexiviridae* and *Tymoviridae*. Boxes represent open reading frames. Replicase gene domains are as in **Figure 3**. M, methyltransferase; A, AlkB; O, OTU-like peptidase; P, papain-like protease; H, RNA helicase; R, RNA-dependent RNA polymerase.

the entire alphavirus-like superfamily (60), as well as in the common ancestor of the flexi-/tymoviral lineage with independent loss of P-Pro in the potexvirus-like viruses and vitiviruses.

Since two of the three major clusters within the *Flexi-/Tymoviridae* lineage possess filamentous capsids, it is reasonable to assume that the common ancestor of this lineage was also a filamentous virus. This notion is supported by the fact that two fungal viruses with evolutionary affinities to either tymoviruses (BCV-F) or potexvirus-like viruses (BVX) also have filamentous particles. Under this scenario, the last common ancestor of the family *Tymoviridae* acquired an unrelated CP, forming icosahedral particles after divergence from the BCV-F ancestor (Figure 6). The alternative, implying that the common ancestor had an icosahedral particle, would require CP replacement on three independent occasions involving ancestors of the potexvirus-like and carlavirus-like viruses, as well as BVF, which seems less likely.

It is a much harder proposition to define the ancestral type of a movement protein for the *Flexi-/Tymoviridae* lineage. Some viruses within the family *Tymoviridae* possess a unique type of movement protein with no homologs outside the family, whereas in others there is no recognizable movement protein at all (79). As described above, the flexiviruses employ either TGB-type or p30-like movement proteins. However, TGB proteins are present in both potexvirus-like and carlavirus-like clusters of flexiviruses, whereas p30-like proteins are present only in the latter cluster. Furthermore, TGBp1 of the flexiviruses are significantly smaller than those found outside this family. This observation may suggest that flexiviral TGBp1 is closer to an ancestral form of this protein, while accretion of additional domains in TGBp1 of rod-shaped viruses occurred later in evolution.

Taken together, these considerations suggest that the common ancestor of the *Flexi-/Tymoviridae* lineage possessed a TGB movement complex that has been lost in

tymoviruses concomitant with the CP replacement. We further propose that after the separation of two flexiviral evolutionary clusters, the potexvirus-like viruses retained TGB, whereas p30-like movement proteins have replaced TGB proteins in a subset of the carlavirus-like genera (Figure 6).

Phylogenetic analysis of the flexiviral CPs reveals strong correlation of the CP evolution with the type of movement protein present in a viral genome (Figure 5a). The larger of the two major CP lineages includes viruses that possess TGB, while the smaller includes capillo-, tricho-, and vitiviruses, all of which have p30-like proteins. This phylogenetic pattern is indicative of coevolution of CPs and movement proteins, possibly due to a coupling of the encapsidation and movement functions in the viral life cycle. The only exception from such protein coevolution in flexiviruses is CLBV, which possesses a p30-like protein in combination with a CP that clusters with those of TGB-harboring viruses. This anomaly can be explained by a relatively recent recombination event that replaced one of the components of the usual movement protein-CP couple.

The distribution pattern of the AlkB domain within replicases of numerous positive-strand RNA viruses represents an interesting evolutionary puzzle. So far, this domain was found in all but one (*Capillovirus*) genera of the *Flexiviridae* (Figure 7) and in none of the *Tymoviridae* genera (Figure 6). Conspicuously, AlkB is present in some, but not all, viruses that belong to the same genus. For instance, among *Potexvirus* members, *Papaya mosaic virus* (PapMV) and CIYMV possess AlkB, whereas PVX and WCIMV do not. Likewise, in the genus *Carlavirus*, AlkB is found in BLSv and *Poplar mosaic virus* (PopMV), but not in *Potato virus M* (PVM) or PVS. There are, however, only a handful of known AlkB-containing viruses outside the family *Flexiviridae* (Figure 7). These include *Grapevine leafroll-associated virus-3* (GLRaV-3) and Little cherry virus-1 (LChV-1) (family *Closteroviridae*), Blackberry virus Y (BVY) (family



<b>Carlavirus</b>	<b>CGRMV</b>	S R F S C L V I Q I Y E F G S K L G F H S D D E S C Y D D D L E V L T V N L F G E A C I A F R R	6	L S D G E F L L M P K G F Q Q G F Q H S V K Y A S N R V S L T F R
	<b>CNRMV</b>	D R F S C L V Q H Y T K A A T L G L H K D D E D C Y E K D H E V M T I N L F C K A T L F F T S	6	L S H G E Y I L M P R G F Q Q S Y K Y H G V K D T Q A G R I S L T F R
	<b>AclV</b>	- K Y D C C L V Q V Y E A E A I G F H A D D E T I F E P G E S V L T I N L E G R A S F G I S C	6	L N G P L Q T M P R G F Q A D H K H C V R G T A G R A S I T F R
	<b>Bisv</b>	T - Y D C M L A Q R Y G A Q K I G F H A D N E E I F M R G A P V H T V S M D G N A D F G T E C	8	L R C N V Q T M P S G F Q E T H K H A V R N T T A G R V S Y T F R
	<b>LSV</b>	K - Y N C M L A Q K Y Q V N S C L G F H A D D E A L F V A G E S I L T V N I T G E A D F K V T C	25	L Q A Q M F E M P P G F Q Q T H K H A V A N C T A G R I S Y T F R
	<b>PopMV</b>	V G Y D C L A Q E Y A Q G K L G F R D D E P N L D V G A P I F T V N L Y C E A T F M L K G	18	L K P S Q C F T M P H F Q E S H K H A V E G S K G R V S L T F R
	<b>SLV</b>	N Y F N C V L F Q Q Y D G H G I G F H S D D E L F E K N S K I L T I C I Q D C E F R F R C	8	M E A P K Q F M M P E G F Q E S H K H A V R G C S P G R I S A T F R
	<b>SPCFV</b>	- Y Y D S V L V Q E Y R A G G I N F H K D D E I F E R G A K V L T V N L N G Q C Y F S F S S	8	L V E D S Y F E M P R D F Q E K Y Y H G V Q G S M G R I S N T F R
	<b>DVS</b>	N Y S C L V Q V Y E A D A T L N W H S D D E S L F E K G R I A T V N L S G A V F H V K C	8	L V G F G F E M P A D F Q S T H K H K L S N P S N G R E S Y T F R
	<b>HplV</b>	- K Y D S C L V Q I Y E A G A L G Y H A D D E L F E Q E S I A T V N L S C A A E F G V K C	8	L R G P Q Q F E M P A F Q V T H K H S V W G S Q R E S Y T F R
<b>Potexvirus</b>	<b>NCLV</b>	- K Y N S C L V Q V Y K A G S S T A L H S D L G M F I P E E G F L T V N L G S A T L S T E C	8	L D C A V F F E M P H G L Q S T H K Y S V D C S P G R A S I T F R
	<b>AlsVX</b>	F Q A N A V L I Q V Y E A N A G I P W H R D E K C Y E K - D A I L T L N F - G T A T F E F S N	8	L T D G C H F I N - S G E W L D L K H R V T N C S A D R M S L T F R
	<b>AlmV</b>	P H F N Q C L I Q K Y Q R G A S I P F H S D N E P C Y D V D H Q V L T I N L T G E A E K T S C	8	L V E N Q F H L S P P G F Q K T H K H S V S L S A G R V S L T F R
	<b>PapMV</b>	P E F N Q C L V Q Q F K L Q A A I P F H R D D E P C Y K G H Q V L T I N H S C E C L T Q I A C	8	M G F G D Y L S P V G F Q E S H K H A V S N T T G R V S L T F R
	<b>CIYMV</b>	T D F D H C L I Q R Y Q N G Y H L R P H S D N E P C Y E A N P I L T I N T E G Q A E F I I S R	8	L G P N S W L L M P S G L Q E T H K H E V I A M S E G R T S I T F R
	<b>CVX</b>	E E H D H C L F V F Q Q A A I N F H S D D E P L I Q P G S P I T L S I - G H C E L L T R D	13	L S G P I I Y T M P S G F Q E T H Q H S V R S L Q K N R L S I T F R
	<b>OpVX</b>	D F F D H C L F Q V F E A S A S I G L H Q D D E A L I A P D T E I V T L N F - G S A D L T T L D	5	L N G T V K Y S M P P G F Q K T H K H A V K S N S A K R I S I T F R
	<b>ZyVX</b>	N E H D H C L F Q V F D K Q S S I Q F H A D D E A L I R P N S I I T V S I - C H C E L L T K E	10	L S G P V V H T M P E G F Q S T H L S I K S L Q E G R L S I T F R
	<b>RSPaV</b>	N Y N S C L A Q I Y E E N S K L A L H K D D E S C Y E I G H K V L T I N L I G S A T F T I S K	10	I G P N E F F E M P R G M Q C N Y F H G V S N C T P G R V S I T F R
	<b>ASPV</b>	N F Y N Q C L V Q E Y S T G H G L S M H K D D E S I Y D I N H Q V L T V N Y S G D A L F C I E C	10	L S G P Q M L L M P F G F Q K E H R H G I K S P S K G R I S L T F R
<b>Trichovirus</b>	<b>ChMLV</b>	V P F N A A L V Q V Y E K D C I L G M H K D N E S C Y G - N H P I L T V N V S G K A V F S T D C	10	L D S G D E L L M P E D F Q R K F R H G V K S I T D G R M S V T L R
	<b>ACLSV</b>	I D F N S A L I Q V Y N D C R L P L H S D N E E C Y D - D D G I L T I N V V G D A K F H T T C	10	L K O G N E I L M P A G Y Q K K N R H A V E A S E G R T S V T L R
<b>Cittrivirus</b>	<b>APCLSV</b>	I D F N A S L I Q I Y E K S I I G M H K D N E E C Y D - D D G V L T L N V K G N A T F S V S C	12	L K E G N E L L M P P G Y Q K K F K H G V K S E S E G R I S V T L R
	<b>CLBV</b>	- D F N C A L I N F Y E A N S S L G F H R D N E R V Y N - D D P I L T V C T F G E G R F T I E F	12	M T A G S F F L M P K G F Q K K A R H S V S N E - M S R V S I T F R
<b>Allexivirus</b>	<b>SnVX</b>	T T Y N Q C L V K Y E Q S R I G F H S D E A I Y P K G N K I L T V N A A G S G T F G I K C	12	L E D G D Y F Q M P S G F Q E T H K H N V V A V - T P R L S F T F R
<b>Vitivirus</b>	<b>GVA</b>	- S Y D H C L I Q R Y T A G S I G F H A D D E P C Y L P G G S V T V N L H C D A T F E V K E	10	L H D G D V Y V M G P G M Q Q T H K H R V T S H T D G R C S I T L R
	<b>GVB</b>	- D F D H C L A Q I Y E E D K I N F N A D D E P C Y T D P - E V V T V N L N G N A N F H L K C	7	L S D G D V L I M P K G F Q K T H K H A V T G T S A G R I S L T F R
<b>Mandarivirus</b>	<b>ICRSV</b>	G K F N T C L A Q T H D Q C A R I G Y H A D D E D C Y D K D V T V A T V N L T G N A T F S L K T	8	L K P G D F I V L K P G A Q Q C T K H A I S D C T T N R T S L T F R

Conserved Invariable

Figure 7

Alignment of AlkB domains from flexiviruses. Conserved amino-acids of the AlkB domain are highlighted in blue, invariable residues in yellow.

**Table 2** Abbreviations, names, taxonomic position and sequence accession numbers of the viruses reported in this article. Only names of definitive virus species are italicized.

Abbreviations	Virus names	Genus	Family	Acc. No
ACLSV	<i>Apple chlorotic leaf spot virus</i>	<i>Trichovirus</i>	<i>Flexiviridae</i>	m58152
AcLV	<i>Aconitum latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	ab051848
AlsVX	<i>Alstroemeria virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ab206396
AltMV	<i>Alternanthera mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay863024
AOPRV	African oil palm ringspot virus	<i>Foveavirus</i>	<i>Flexiviridae</i>	af468004
APCLSV	<i>Apricot pseudo-chlorotic leaf spot virus</i>	<i>Trichovirus</i>	<i>Flexiviridae</i>	ay713379
ApLV	<i>Apricot latent virus</i>	<i>Foveavirus</i>	<i>Flexiviridae</i>	af057035
ApruV-1	Asian prunus virus 1	<i>Foveavirus</i>	<i>Flexiviridae</i>	dq205236
ApruV-2	Asian prunus virus 2	<i>Foveavirus</i>	<i>Flexiviridae</i>	dq205237
ApruV-3	Asian prunus virus 3	<i>Foveavirus</i>	<i>Flexiviridae</i>	dq205238
ASGV	<i>Apple stem grooving virus</i>	<i>Capillovirus</i>	<i>Flexiviridae</i>	d14995
ASPV	<i>Apple stem pitting virus</i>	<i>Foveavirus</i>	<i>Flexiviridae</i>	d21829
AYVV	<i>Anagryis vein yellowing virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	ay751780
BaMV	<i>Bamboo mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	d26017
BanMMV	<i>Banana mild mosaic virus</i>	Unassigned	<i>Flexiviridae</i>	af314662
BanVX	Banana virus X	Unassigned	<i>Flexiviridae</i>	ay710267
BCV-F	Botrytis cinerea virus F	Unassigned ssRNA+ virus		af238884
BVY	Blackberry virus Y	Unassigned	<i>Potyviridae</i>	ay994084
BRpNV	Black raspberry necrosis virus	<i>Sadwavirus</i>	Unassigned	Not available
BlSeV	<i>Blueberry scorch virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	l25658
BMLV	Bombyx mori Macula-like latent virus	<i>Maculavirus</i>	<i>Tymoviridae</i>	ab186123
BVX	Botrytis virus X	Unassigned ssRNA+ virus	<i>Flexiviridae</i>	ay055762
CGRMV	Cherry green ring mottle virus	Unassigned	<i>Flexiviridae</i>	af017780
ChMLV	Cherry mottle leaf virus	<i>Trichovirus</i>	<i>Flexiviridae</i>	af170028
ChMV	<i>Chayote mosaic virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	af195000
CLBV	<i>Citrus leaf blotch virus</i>	<i>Citricivirus</i>	<i>Flexiviridae</i>	aj318061
CLV	<i>Carnation latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj010697
CIYMV	<i>Clover yellow mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	d29630
CNRMV	Cherry necrotic rusty mottle virus	Unassigned	<i>Flexiviridae</i>	af237816
CoLV	<i>Cole latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	ay340584
CPMMV	<i>Cowpea mild mottle virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	af024629
CsCMV	<i>Cassava common mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	u23414
CSDaV	Citrus sudden death-associated virus	<i>Marafivirus</i>	<i>Tymoviridae</i>	ay884005
CTV	<i>Citrus tristeza virus</i>	<i>Closterovirus</i>	<i>Closteroviridae</i>	ay170468
CVA	<i>Cherry virus A</i>	<i>Capillovirus</i>	<i>Flexiviridae</i>	x82547
CVB	Chrysanthemum virus B	<i>Carlavirus</i>	<i>Flexiviridae</i>	s60150
CVX	<i>Cactus virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	af308158
CymMV	<i>Cymbidium mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	u62963
DuMV	<i>Dulcamara mottle virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	ay789137
DVS	Daphne virus S	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj620300

(Continued)

Table 2 (Continued)

Abbreviations	Virus names	Genus	Family	Acc. No
EMV	<i>Eggplant mosaic virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	j04374
ErLV	<i>Erysimum latent virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	af098523
FoMV	<i>Foxtail mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	m62730
GarV-A	<i>Garlic virus A</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	ab010300
GarV-B	<i>Garlic virus B</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	ab010301
GarV-C	<i>Garlic virus C</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	ab010302
GarV-D	<i>Garlic virus D</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	ab010303
GarV-E	<i>Garlic virus E</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	aj292230
GarV-X	<i>Garlic virus X</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	aj292229
GarMblV	Garlic mite-borne latent virus	<i>Allexivirus</i>	<i>Flexiviridae</i>	Not available
GCLV	<i>Garlic common latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	x81138
GFkV	<i>Grapevine fleck virus</i>	<i>Maculavirus</i>	<i>Tymoviridae</i>	aj309022
GINV	<i>Grapevine berry inner necrosis virus</i>	<i>Trichovirus</i>	<i>Flexiviridae</i>	d88448
GLRaV-3	<i>Grapevine leafroll associated virus-3</i>	<i>Ampelovirus</i>	<i>Closteroviridae</i>	af037268
GRSPaV	<i>Grapevine rupestris stem pitting-associated virus</i>	<i>Foveavirus</i>	<i>Flexiviridae</i>	af026278
GVA	<i>Grapevine virus A</i>	<i>Vitivirus</i>	<i>Flexiviridae</i>	x75433
GVB	<i>Grapevine virus B</i>	<i>Vitivirus</i>	<i>Flexiviridae</i>	x75448
GVD	<i>Grapevine virus D</i>	<i>Vitivirus</i>	<i>Flexiviridae</i>	y07764
HdRSV	<i>Hydrangea ringspot virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay707100
HpLV	<i>Hop latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	ab032469
HpMV	<i>Hop mosaic virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	ab051109
HVX	<i>Hosta virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	aj620114
ICRSV	<i>Indian citrus ringspot virus</i>	<i>Mandarinivirus</i>	<i>Flexiviridae</i>	af406744
KLV	<i>Kalanchoe latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj293570
KYMV	<i>Kennedya yellow mosaic virus</i>	<i>Tymovirus</i>	<i>Tymovirus</i>	d00637
LiCLSV	<i>Lilac chlorotic leafspot virus</i>	<i>Capillovirus</i>	<i>Flexiviridae</i>	Not available
LSV	<i>Lily symptomless virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj516059
LVX	<i>Lily virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	aj633822
LChV-1	Little cherry virus 1	Unassigned	<i>Closteroviridae</i>	y10237
MRFV	<i>Maize rayado fino virus</i>	<i>Marafivirus</i>	<i>Tymoviridae</i>	af265566
MXV	Mint virus X	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay789138
NCLV	<i>Narcissus common latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	am158439
NeLV	<i>Nerine latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	dq098905
NMV	<i>Narcissus mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	d13747
NRNV	Nemesia ring necrosis virus	<i>Tymovirus</i>	<i>Tymoviridae</i>	ay751778
NSV	Narcissus symptomless virus	<i>Carlavirus</i>	<i>Flexiviridae</i>	am182569
NSPV	Nandina stem pitting virus	<i>Capillovirus</i>	<i>Flexiviridae</i>	Not available
NVX	<i>Nerine virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ab219105
OBdV	<i>Oat blue dwarf virus</i>	<i>Marafivirus</i>	<i>Tymoviridae</i>	u87832
OmbLV	Onion mite-borne latent virus	<i>Allexivirus</i>	<i>Flexiviridae</i>	Not available
OpVX	Opuntia virus X	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay366209
OYMV	<i>Ononis yellow mosaic virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	j04375
PAMV	<i>Potato aucuba mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	s73580

(Continued)

Table 2 (Continued)

Abbreviations	Virus names	Genus	Family	Acc. No
PapMV	<i>Papaya mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	d13957
PcMV	<i>Peach mosaic virus</i>	<i>Trichovirus</i>	<i>Flexiviridae</i>	dq117579
PepMV	<i>Pepino mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	af484251
PeSRSV	<i>Peach sooty ringspot virus</i>	<i>Foveavirus</i>	<i>Flexiviridae</i>	af318062
PeSV	<i>Pea streak virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	af354652
PhyMV	<i>Physalis mottle virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	y16104
PLAMV	<i>Plantago asiatica mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	z21647
PnMV	<i>Poinsettia mosaic virus</i>	Unassigned	<i>Tymoviridae</i>	aj271595
PopMV	<i>Poplar mosaic virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	x65102
PVM	<i>Potato virus M</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	d14449
PVS	<i>Potato virus S</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj863509
PVT	<i>Potato virus T</i>	Unassigned	<i>Flexiviridae</i>	d10172
PVX	<i>Potato virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	x05198
PVY	<i>Potato virus Y</i>	<i>Potyvirus</i>	<i>Potyviridae</i>	x12456
RCVMV	<i>Red clover vein mosaic virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	Not available
ScaVX	<i>Scallion virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	aj316085
SchVX	<i>Schlumbergera virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay366207
ScrMV	<i>Scrophularia mottle virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	ay751777
SCSMaV	<i>Sugarcane striate mosaic-associated virus</i>	Unassigned	<i>Flexiviridae</i>	af315308
ShMbLV	<i>Shallot mite-borne latent virus</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	Not available
ShVX	<i>Shallot virus X</i>	<i>Allexivirus</i>	<i>Flexiviridae</i>	m97264
SLV	<i>Shallot latent virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	aj292226
SMYEV	<i>Strawberry mild yellow edge virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	d12517
SPCFV	<i>Sweet potato chlorotic fleck virus</i>	<i>Carlavirus</i>	<i>Flexiviridae</i>	ay461421
SsDRV	<i>Sclerotinia sclerotiorum</i> debilitation-associated RNA virus	Unassigned ssRNA+ virus		ay147260
TMV	<i>Tobacco mosaic virus</i>	<i>Tobamovirus</i>	Unassigned	v01408
TVX	<i>Tulip virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ab066288
TYMV	<i>Turnip yellow mosaic virus</i>	<i>Tymovirus</i>	<i>Tymoviridae</i>	x07441
WCIMV	<i>White clover mosaic virus</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	x06728
ZyVX	<i>Zygocactus virus X</i>	<i>Potexvirus</i>	<i>Flexiviridae</i>	ay366208

*Potyviridae* (109), and Black raspberry necrosis virus (BRpNV) (genus *Sadwavirus*) (49).

Close inspection of the list of AlkB-containing viruses (Figure 7) allows two generalizations: (i) almost exclusively, the AlkB domain is found in replicases of the three families of filamentous plant viruses; (ii) most, if not all, AlkB-containing viruses infect woody or perennial plants. The latter tendency may be attributed to the functional advantages provided by AlkB for the long-term survival

of viruses within the single infected plant. Indeed, the cellular AlkB was implicated in repair of RNA methylation damage (1), an activity that would provide an obvious benefit of higher genetic stability for viral RNA genomes in woody or perennial plants.

A patchy distribution of AlkB among viral genomes makes it difficult to fathom its exact evolutionary history that would necessarily require multiple events of acquisition and/or loss within different virus taxa.



Because AlkB is most frequently present in the viruses of carlaviruses-like cluster of *Flexiviridae*, we marginally favor its inclusion into a common ancestor of the viruses in this cluster (**Figure 6**). Since AlkB is widely conserved in cellular organisms from bacteria to humans, but found only in a minority of plant-infecting viruses, it was most likely acquired into virus genomes via recombination with the corresponding host mRNA. The following evolution of AlkB has likely included its repeated loss in certain carlaviruses-like viruses and acquisition by some potexvirus-like viruses via recombination of the viruses coinfecting the same host plant. It also does not seem unreasonable to suggest that *Flexiviridae* donated AlkB to a few AlkB-containing viruses outside this family.

A similar line of reasoning applies to the mosaic distribution of the OTU-like protease among the subset of carlaviruses-like viruses. Under the most parsimonious scenario, the OTU domain was acquired by the common ancestor of the carla- and fovea-viruses and donated to citrivirus via horizontal gene transfer (**Figure 6**).

Another protein of uncertain evolutionary history is the Zn-ribbon-containing, RNA-binding protein present in five flexiviral genera (**Figure 3**). These proteins are so variable in sequence that alignments have proven unsuitable for robust phylogenetic reconstruction (data not shown). An apparently random distribution of these five genera in both evolutionary clusters of the flexiviruses (**Figure 6**) is compatible with two scenarios. According to the first, the RNA-binding protein was present in the common ancestor and was lost on multiple occasions. According to a second scenario, this protein was acquired by one of the genera (e.g., *Carlavirus*), and then distributed to other flexiviruses through horizontal gene transfer. It appears that flexiviruses may have donated a related RNA-binding protein to a single virus within the genus *Closterovirus*, *Citrus tristeza virus* (CTV), where it functions as a suppressor of RNA silencing (20, 72).

## CONCLUSIONS AND TAXONOMIC IMPLICATIONS

The most prominent aspect of the evolutionary scenario devised here and summarized in **Figure 6** is the unusual fluidity of the flexiviral genomes compared to those of the related rod-shaped or icosahedral RNA viruses of the alphavirus-like superfamily. This fluidity could be explained by the relative ease with which the filamentous virions can accommodate the increase or reduction in the size of the viral genome. Indeed, the only other family of plant viruses that exhibits comparable genomic variability is an evolutionarily distant family of filamentous alphavirus-like viruses, *Closteroviridae* (31).

There is no region of flexiviral genomes that is left untouched by genetic variation. Even the most conservative genomic core that encodes the RNA replicase exhibits multiple events of acquisition or loss of such domains as P-Pro, OTU-like protease, or AlkB. Another major determinant of the viral life style, the virion architecture that is sometimes defined as nothing less than “the viral self” (9), underwent a dramatic transformation at the descent of the *Tymoviridae* family from a common ancestor with potexvirus-like viruses. The levity with which flexiviruses switch the modes of their cell-to-cell movement between TGB and p30-like proteins is also amusing. This levity seems to be even more extreme in the *Tymoviridae* lineage where some viruses possess exotic movement proteins or no apparent movement protein at all, as in the genus *Marafivirus*. The genes for nucleic acid-binding proteins that wander among flexiviral genomes complete the picture of genetic unrest that seems to reign within the *Flexi-/Tymoviridae* evolutionary lineage.

Do flexiviruses benefit from their genetic plasticity? It seems that the answer is yes if one considers the ability of flexiviruses to find novel ecological niches, e.g., by evolving into fungal viruses and sometimes changing the capsid architecture or even leaving it behind, as is the case for SsDRV. Another aspect of

flexiviruses that stands out is their propensity for infecting woody plants. A rapidly growing list of flexiviruses found in trees, shrubs, and vines, from poplars to blueberries to hops (**Table 2**) provides a testimony to this notion.

In addition to revealing the vagaries of virus evolution, our analysis illustrates the difficulty of classifying viruses that unceasingly evolve via exchange of genetic modules. If taxonomy is to reflect the likely evolutionary pathways, it should be based on phylogenetic analysis of the only character shared by all positive-strand RNA viruses, which is the RdRp. We generally followed this line of thought even though we expanded our analysis by inclusion of the Met and Hel domains conserved in the alphavirus-like superfamily (**Figure 4**). A radical taxonomic interpretation of this analysis would call for separa-

tion of the existing family *Flexiviridae* into two families corresponding to potexvirus-like and carlavirus-like lineages. A more conformist alternative to this interpretation would take into account additional evolutionary considerations. One of these is a shared origin and morphology of the capsid. Another consideration is that the genomes of flexiviruses are peppered with such genetic modules as TGB and genes encoding a nucleic acid-binding protein or an AlkB domain. Importantly, none of these modules is found in the otherwise closely related family *Tymoviridae*, and, except for TGB and 30 k, these proteins are seldom seen outside flexiviruses. Taken together, our phylogenetic reconstructions seem to favor retaining the family *Flexiviridae*, with the possible establishment of two subfamilies, to account for the bifurcated evolutionary history of the viral replicases.

## ACKNOWLEDGMENTS

The research in V.V.D. laboratory is supported in part by grants from NIH (GM015390) and BARD (IS-3784-05). Rothamsted Research receives grant-aided support from the Biotechnology and Biological Sciences Research Council of the United Kingdom. J.F.K. is funded by the Belgium Directorate General for Development Co-operation.

## LITERATURE CITED

1. Aas PA, Otterlei M, Falnes PO, Vagbo CB, Skorpen F, et al. 2003. Human and bacterial oxidative demethylases repair alkylation damage in both RNA and DNA. *Nature* 421:859–63
2. Adams MJ, Accotto GP, Agranovsky AA, Bar-Joseph M, Boscia D, et al. 2005. Family *Flexiviridae*. In *Virus Taxonomy. Eighth Rep. Int. Comm. Taxon. Viruses*, ed. CM Fauquet, MA Mayo, J Maniloff, U Desselberger, LA Ball, pp. 1089–124. Amsterdam: Elsevier/Academic
3. Ahlquist P. 2006. Parallels among positive-strand RNA viruses, reverse-transcribing viruses and double-stranded RNA viruses. *Nat. Rev. Microbiol.* 4:371–82
4. Ahmed N, Christie S, Zettler F. 1983. Identification and partial characterization of a closterovirus infecting *Nandina domestica*. *Phytopathology* 73:470–75
5. Aravind L, Koonin EV. 2001. The DNA repair protein AlkB, EGL-9, and leprecan define new families of 2-oxoglutarate- and iron-dependent dioxygenases. *Genome Biol.* 2(3):research007.1–research007.8
6. Barg E, Lesemann DE, Vetten HJ, Green SK. 1994. Identification, partial characterization and distribution of viruses infecting allium crops in south and south-east Asian countries. *Acta Hort.* 358:251–58

7. Batten JS, Yoshinari S, Hemnway C. 2003. *Potato virus X*: a model system for virus replication, movement and gene expression. *Mol. Plant Pathol.* 4:125–31
8. Bayne EH, Rakitina DV, Morozov SY, Baulcombe DC. 2005. Cell-to-cell movement of potato potexvirus X is dependent on suppression of RNA silencing. *Plant J.* 44:471–82
9. Benson SD, Bamford JKH, Bamford DH, Burnett RM. 2004. Does common architecture reveal a viral lineage spanning all three domains of life? *Mol. Cell* 16:673–85
10. Bonavia M, Digiario M, Boscia D, Boari A, Bottalico G, et al. 1996. Studies on “corky rugose wood” of grapevine and the diagnosis of grapevine virus B. *Vitis* 35:53–58
11. Boscia D, Savino V, Minafra A, Namba S, Elicio V, et al. 1993. Properties of a filamentous virus isolated from grapevines affected by corky bark. *Arch. Virol.* 130:109–20
12. Boscia D, Minafra A, Martelli GP. 1997. Filamentous viruses of the grapevine: putative trichoviruses and capilloviruses. See Ref. 87a, pp. 19–28
13. Bratlie MS, Drabløs F. 2005. Bioinformatic mapping of AlkB homology domains in viruses. *BMC Genomics* 6:1–15, doi:10.1186/1471-2164-6-1
14. Brigneti G, Voinnet O, Li WX, Ji LH, Ding SW, et al. 1998. Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*. *EMBO J.* 17:6739–46
15. Brunt AA. 1978. The occurrence, hosts and properties of lilac chlorotic leafspot virus, a newly recognized virus from *Syringa vulgaris*. *Ann. Appl. Biol.* 88:383–91
16. Brunt AA, Stace Smith R. 1978. The intracellular location of lilac chlorotic leafspot virus. *J. Gen. Virol.* 39:63–71
17. Brunt AA, Crabtree K, Dallwitz MJ, Gibbs AJ, Watson L, eds. 1996. *Viruses of Plants. Descriptions and Lists from the VIDE Database*. Wallingford, UK: CAB Int. 1484 pp.
18. Byadgi AS, Ahlawat YS, Chakraborty NK, Varma A, Srivastava M, et al. 1993. Characterization of a filamentous virus associated with citrus ringspot in India. *Proc. Conf. Int. Org. Citrus Virol.*, 12th, Riverside, pp. 184–87
19. Callaway A, Giesman-Cookmeyer D, Gillock ET, Sit TL, Lommel SA. 2001. The multifunctional capsid proteins of plant RNA viruses. *Annu. Rev. Phytopathol.* 39:419–60
20. Chiba M, Reed JC, Prokhnovsky AI, Chapman EJ, Mawassi M, et al. 2006. Diverse suppressors of RNA silencing enhance agroinfection by a viral replicon. *Virology* 346:7–14
21. Cheng JH, Peng CW, Hsu YH, Tsai CH. 2002. The synthesis of minus-strand RNA of Bamboo mosaic potexvirus initiates from multiple sites within the poly(A) tail. *J. Virol.* 76:6114–20
22. Chevalier S, Greif C, Clauzel JM, Walter B, Fritsch C. 1995. Use of an immunocapture-polymerase chain reaction procedure for the detection of grapevine virus A in Kober stem grooving-infected grapevines. *J. Phytopathol.* 143:368–73
23. Christie RG, Edwardon JR. 1977. Light and electron microscopy of plant virus inclusions. *Fla. Agric. Exp. Stn. Monogr. Ser.* No 9, 155 pp.
24. Claverie JM. 2006. Viruses take center stage in cellular evolution. *Genome Biol.* 7(6):110
25. Culley AI, Lang AS, Suttle CA. 2006. Metagenomic analysis of coastal RNA virus communities. *Science* 312:1795–98
26. Davenport GF, Baulcombe DC. 1997. Mutation of the GKS motif of the RNA-dependent RNA polymerase from potato virus X disables or eliminates virus replication. *J. Gen. Virol.* 78:1247–51
27. Desvignes JC, Boyé R, Cornaggia D, Grasseau N. 1999. *Maladies à Virus des Arbres Fruitières*. Paris: Cent. Tech. Interprof. Fruits Legumes. 202 pp.
28. Dolja VV. 2003. Beet yellows virus: the importance of being different. *Mol. Plant Pathol.* 4:91–98

29. Dolja VV, Boyko VP, Agranovsky AA, Koonin EV. 1991. Phylogeny of capsid proteins of rod-shaped and filamentous RNA plant viruses: two families with distinct patterns of sequence and probably structure conservation. *Virology* 184:79–86
30. Dolja VV, Grama DP, Morozov SY, Atabekov JG. 1987. Potato virus X-related single- and double-stranded RNAs. *FEBS Lett.* 214:308–12
31. Dolja VV, Kreuze JF, Valkonen JPT. 2006. Comparative and functional genomics of closteroviruses. *Virus Res.* 117:38–51
32. Dreher TW. 1999. Functions of the 3'-untranslated regions of positive strand RNA viral genomes. *Annu. Rev. Phytopathol.* 37:151–74
33. Falk BW, Duffus JE. 1988. Ecology and control. See Ref. 86, 4:275–96
34. Faoro F. 1997. Cytopathology of closteroviruses and trichoviruses infecting grapevines. See Ref. 87a, pp. 29–47
35. Forterre P. 2006. The origin of viruses and their possible roles in major evolutionary transitions. *Virus Res.* 117:5–16
36. Galiakparov N, Goszczynski DE, Che X, Batuman O, Bar-Joseph M, et al. 2003. Two classes of subgenomic RNA of *Grapevine virus A* produced by internal controller elements. *Virology* 312:434–48
37. Galiakparov N, Tanne E, Mawassi M, Gafny R, Sela I. 2003. ORF 5 of grapevine virus A encodes a nucleic acid-binding protein and affects pathogenesis. *Virus Genes* 27:257–62
38. Galiakparov N, Tanne E, Sela I, Gafny R. 2003. Functional analysis of the grapevine virus A genome. *Virology* 306:42–50
39. Galipienso L, Vives MC, Moreno P, Milne RG, Navarro L, et al. 2001. Partial characterization of citrus leafblotch virus, a new virus from Nagami kumquat. *Arch. Virol.* 146:357–68
40. Gambley CF, Thomas JE. 2001. Molecular characterization of Banana mild mosaic virus, a new filamentous virus in *Musa* spp. *Arch. Virol.* 146:1369–79
41. Garau R, Prota VA, Piredda R, Boscia D, Prota U. 1994. On the possible relationship between Kober stem grooving and grapevine virus A. *Vitis* 33:161–63
42. Gentit P, Foissac X, Svanella-Dumas L, Candresse T. 2001. Variants of *Apricot latent foveavirus* (ALV) isolated from south European orchards associated with peach asteroid spot and peach sooty ringspot diseases. *Acta Hort.* 550:213–20
43. Goldbach R. 1987. Genome similarities between plant and animal RNA viruses. *Microbiol. Sci.* 4:197–202
44. Gorbalenya AE, Koonin EV. 1993. Helicases: amino acid sequence comparisons and structure-function relationship. *Curr. Opin. Struct. Biol.* 3:419–29
45. Goszczynski DE, Joste AEC. 2003. Identification of divergent variants of *Grapevine virus A*. *Eur. J. Plant Pathol.* 109:397–403
46. Goszczynski DE. 2006. Molecular variants of *Grapevine virus A* (GVA) associated with Shiraz disease in South Africa. *Extended Abstr. 15th Meet. ICVG, Stellenbosch*, pp. 72–73
47. Gramstat A, Courtpozanis A, Rohde W. 1990. The 12 kDa protein of potato virus M displays properties of a nucleic acid-binding regulatory protein. *FEBS Lett.* 276:34–38
48. Hadidi A, Khetarpal RK, Koganezawa H, eds. 1998. *Plant Virus Disease Control*. St. Paul, MN: APS Press
49. Halgren A, Tzanetakis IE, Martin RR. 2007. Identification, characterization and detection of Black raspberry necrosis virus. *Phytopathology* 97:44–50
50. Haviv S, Goszczynski DE, Mawassi M. 2006. The p10 of *Grapevine virus A* affects pathogenicity on *Nicotiana benthamiana* plants. *Extended Abstr. 15th Meet. ICVG, Stellenbosch*, pp. 69–71



51. Howitt RLJ, Beever RE, Pearson MN, Forster RLS. 2001. Genome characterization of Botrytis virus F, a flexuous rod-shaped mycovirus resembling plant 'potex-like' viruses. *J. Gen. Virol.* 82:67–78
52. Howitt RL, Beever RE, Pearson MN, Forster RL. 2006. Genome characterization of a flexuous rod-shaped mycovirus, Botrytis virus X, reveals high amino acid identity to genes from plant 'potex-like' viruses. *Arch. Virol.* 151:563–79
53. Huang YL, Han YT, Chang YT, Hsu YH, Meng M. 2004. Critical residues for GTP methylation and formation of the covalent m7GMP-enzyme intermediate in the capping enzyme domain of Bamboo mosaic virus. *J. Virol.* 78:1271–80
54. James D, Varga H, Croft H, Rast H, Thompson D, et al. 2006. Molecular characterization, phylogenetic relationships, and specific detection of *Peach mosaic virus*. *Phytopathology* 96:137–44
55. Jelkmann W. 1997. Apple stem pitting virus. See Ref. 87a, pp. 133–42
56. Kamer G, Argos P. 1984. Primary structural comparison of RNA-dependent polymerases from plant, animal and bacterial viruses. *Nucleic Acids Res.* 12:7269–82
57. Karpova OV, Zayakina OV, Arkhipenko MV, Sheval EV, Kiselyova OI, et al. 2006. Potato virus X RNA-mediated assembly of single-tailed ternary coat protein-RNA-movement protein complexes. *J. Gen. Virol.* 87:2731–40
58. Keifer HH, Wilson NS. 1955. A new species of eriophyd mite responsible for the vection of peach mosaic virus. *Bull. Calif. Dep. Agric.* 44:145–46
59. Koenig R. 1982. Carlavirus group. *CMI/AAB Descriptions of Plant Viruses*, No. 259
60. Koonin EV, Dolja VV. 1993. Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Crit. Rev. Biochem. Mol. Biol.* 28:375–430
61. Koonin EV, Dolja VV. 2006. Evolution of complexity in the viral world: the dawn of a new vision. *Virus Res.* 117:1–4
62. Koonin EV, Senkevich TG, Dolja VV. 2006. The ancient virus world and evolution of cells. *Biol. Direct* 1:1–27, doi:10.1186/1745-6150-1-29
63. Kumar S, Tamura K, Nei M. 2004. MEGA3: Integrated software for molecular evolutionary genetics analysis and sequence alignment. *Briefings Bioinform.* 5:150–63
64. Kunugi Y, Asari S, Terai Y, Shinkai A. 2000. Studies on the grapevine berry inner necrosis virus disease. 2. Transmission of berry inner necrosis virus by the grape erineum mite *Colomerus vitis* in Yamanashi. *Bull. Yamanashi Fruit Tree Exp. Stn.* 10:57–63
- 64a. Kurstak E, ed. 1981. *Handbook of Plant Virus Infections and Comparative Diagnosis*. Amsterdam: Elsevier/North Holland
65. Lang AS, Culley AI, Suttle CA. 2004. Genome sequence and characterization of a virus (HaRNAV) related to picorna-like viruses that infects the marine toxic bloom-forming alga *Heterosigma akashiwo*. *Virology* 320:206–17
66. La Notte P, Buzkan N, Choueiri E, Minafra A, Martelli GP. 1997. Acquisition and transmission of Grapevine virus A by the mealybug *Pseudococcus longispinus*. *J. Plant Pathol.* 79:79–85
67. Larsen HJ, Oldfield GN. 1995. Peach mosaic. In *Compendium of Stone Fruits Diseases*, ed. JM Ogawa, EI Zher, GW Bird, DF Ritchie, K Uriu, et al., pp. 67–68. St. Paul, MN: APS Press
68. Lawrence DM, Rozanov MN, Hillman BI. 1995. Autocatalytic processing of the 223-kDa protein of blueberry scorch carlavirus by a papain-like proteinase. *Virology* 207:127–35
69. Lazarovitz SG, Beachy RN. 1999. Viral movement proteins as probes for intracellular and intercellular trafficking in plants. *Plant Cell* 11:535–48

70. Lesemann DE. 1988. Cytopathology. See Ref. 86, 4:179–235
71. Li YI, Shih TW, Hsu YH, Han YT, Huang YL, et al. 2001. The helicase-like domain of plant potexvirus replicase participates in formation of RNA 5' cap structure by exhibiting RNA'-triphosphatase activity. *J. Virol.* 75:12114–20
72. Lu R, Folimonov AS, Shintaku M, Li WX, Falk BW. 2004. Three distinct suppressors of RNA silencing encoded by a 20-kb viral RNA genome. *Proc. Natl. Acad. Sci. USA* 101:15742–47
73. Magome H, Terauchi H, Yoshikawa N, Takahashi T. 1997. Analysis of double-stranded RNA in tissues infected with apple stem grooving capillovirus. *Ann. Phytopathol. Soc. Jpn.* 63:450–54
74. Makarova KS, Aravind L, Koonin EV. 2000. A novel superfamily of predicted cysteine proteases from eukaryotes, viruses and *Chlamydia pneumoniae*. *Trends Biochem. Sci.* 25:50–52
75. Marais A, Svanella-Dumas L, Foissac X, Gentit P, Candresse T. 2006. Asian prunus viruses: new related members of the family *Flexiviridae* in *Prunus* germplasm of Asian origin. *Virus Res.* 120:176–83
76. Martelli GP, ed. 1993. *Graft-Transmissible Diseases of Grapevines. Handbook for Detection and Diagnosis*. Rome: FAO Publ. Div. 263 pp.
77. Martelli GP, Russo M. 1984. The use of thin sectioning for visualization and identification of plant viruses. *Meth. Virol.* 8:143–224
78. Martelli GP, Agranovsky AA, Bar-Joseph M, Boscia D, Candresse T, et al. 2002. The family *Closteroviridae* revised. *Arch. Virol.* 147:2039–44
79. Martelli GP, Sabanadzovic S, Abou-Ghanem Sabanadzovic N, Edwards MC, Dreher T. 2002. The family *Tymoviridae*. *Arch. Virol.* 147:1837–46
80. GP M, Boudon-Padieu E. 2006. Directory of major infectious diseases of the grapevine. *Options Mediterr.* B55:11–201
81. Maury Y, Duby C, Khetarpal RH. 1998. Seed certification for viruses. See Ref. 48, pp. 237–48
82. Melcher U. 2000. The '30K' superfamily of viral movement proteins. *J. Gen. Virol.* 81:257–66
83. Meng B, Zhu HY, Gonsalves D. 1999. Rupestris stem pitting-associated virus 1 consists of a family of sequence variants. *Arch. Virol.* 144:2071–85
84. Meng B, Li C, Wang W, Goszczynski D, Gonsalves D. 2005. Complete genome sequence of two new variants of Grapevine rupestris stem pitting-associated virus and comparative analysis. *J. Gen. Virol.* 86:1555–60
85. Milne RG. 1984. Electron microscopy for the identification of plant viruses in in vitro preparations. *Meth. Virol.* 7:87–120
86. Milne RG, ed. 1988. *The Plant Viruses*. Vol. 4: *The Filamentous Plant Viruses*. New York: Plenum. 423 pp.
87. Mink GI. 1993. Pollen- and seed-transmitted viruses and viroids. *Annu. Rev. Phytopathol.* 31:375–402
- 87a. Monette PL, ed. 1997. *Filamentous Viruses of Woody Plants*. Trivandrum: Res. Signpost
88. Morales FJ, Lozano I, Velasco AC, Arroyave GA. 2002. Detection of a fovea-like virus in African oil palm affected by a lethal "ringspot" disease in South America. *J. Phytopathol.* 150:611–15
89. Morozov SY, Dolja VV, Atabekov JG. 1989. Probable reassortment of genomic elements among elongated RNA-containing plant viruses. *J. Mol. Evol.* 29:52–62
90. Morozov SY, Miroshnichenko NA, Solovyev AG, Fedorkin ON, Zelenina DA, et al. 1991. Expression strategy of the potato virus X triple gene block. *J. Gen. Virol.* 72:2039–42

91. Morozov SY, Solovyev AG. 2003. Triple gene block: modular design of a multifunctional machine for plant virus movement. *J. Gen. Virol.* 84:1351–66
92. Mushegian AR, Koonin EV. 1993. Cell-to-cell movement of plant viruses. *Arch. Virol.* 133:239–57
93. Nagasaki K, Shirai Y, Takao Y, Mizumoto H, Nishida K, et al. 2005. Comparison of genome sequences of single-stranded RNA viruses infecting the bivalve-killing dinoflagellate *Heterocapsa circularisquama*. *Appl. Environ. Microbiol.* 71:8888–94
94. Nemchinov LG, Shamloul AM, Zemtchik EZ, Verderevskaya TD, Hadidi A. 2000. Apricot latent virus: a new species in the genus Foveavirus. *Arch. Virol.* 145:1801–13
95. Nuss DL. 2005. Hypovirulence: mycoviruses at the fungal-plant interface. *Nat. Rev. Microbiol.* 3:632–42
96. Ohki ST. 1997. Cytopathology of filamentous viruses infecting fruit trees and woody ornamentals. See Ref. 87a, pp. 171–77
97. Oldfield GN, Proeseler G. 1996. Eriophyoid mites as vectors of plant viruses. In *Eriophyoid Mites, Their Biology, Natural Enemies and Control*, ed. EE Lindquist, MW Sabelis, J Bruin, pp. 259–75. Amsterdam: Elsevier
98. Parker L, Kendall A, Stubbs G. 2002. Surface feature of Potato virus X from fiber diffraction. *Virology* 300:291–95
99. Peremyslov VV, Andreev IA, Prokhnevsky AI, Duncan GH, Taliansky ME, et al. 2004. Complex molecular architecture of beet yellows virus particles. *Proc. Natl. Acad. Sci. USA* 101:5030–35
100. Purcifull DE, Edwardson JR. 1981. Potexviruses. See Ref. 64a, pp. 628–93
101. Roistacher CN. 1991. *Graft-Transmissible Diseases of Citrus. Handbook for Detection and Diagnosis*. Rome: FAO Publ. Div. 286 pp.
102. Rosciglione B, Castellano MA, Martelli GP, Savino V, Cannizzaro G. 1983. Mealybug transmission of grapevine virus A. *Vitis* 22:331–47
103. Rozanov MN, Morozov SY, Skryabin KG. 1990. Unexpected close relationship between the large nonvirion proteins of filamentous potexviruses and spherical tymoviruses. *Virus Genes* 3:373–79
104. Rozanov MN, Koonin EV, Gorbalenya AE. 1992. Conservation of the putative methyltransferase domain: a hallmark of the ‘Sindbis-like’ supergroup of positive-strand RNA viruses. *J. Gen. Virol.* 73:2129–34
105. Saldarelli P, Minafra A, Castellano MA, Martelli GP. 2000. Immunodetection and sub-cellular localization of the proteins encoded by ORF 3 of grapevine viruses A and B. *Arch. Virol.* 145:1535–42
106. Satyanarayana T, Gowda S, Ayllon MA, Dawson WO. 2004. Closterovirus bipolar virion: evidence for initiation of assembly by minor coat protein and its restriction to the genomic RNA 5' region. *Proc. Natl. Acad. Sci. USA* 101:799–804
107. Shi BJ, Habili N, Gafny R, Symons RH. 2004. Extensive variations of sequence within isolates of Grapevine virus B. *Virus Genes* 29:279–85
108. Stubbs G, Parker L, Junn J, Kendall A. 2005. Flexible filamentous virus structures from fiber diffraction. *Fiber Diffract. Rev.* 13:38–42
109. Susaimuthu J, Tzanetakis IE, Gergerich RC, Martin RR. 2006. Blackberry virus Y: a potyvirus component of blackberry yellows disease. *Phytopathology* 96:S112
110. Takao Y, Mise K, Nagasaki K, Okuno T, Honda D. 2006. Complete nucleotide sequence and genome organization of a single-stranded RNA virus infecting the marine fungoid protist *Schizochytrium* sp. *J. Gen. Virol.* 87:723–33

111. Terai Y, Yanase H. 1992. Induction of berry necrosis in Kyoho back-inoculated with the virus isolate from grapevine mosaic diseases clones and renaming to grapevine berry inner necrosis. *Ann. Phytopathol. Soc. Jpn.* 58:617
112. Torrance L, Andreev IA, Gabrenaitė-Verhovskaya R, Cowan G, Makinen K, et al. 2005. An unusual structure at one end of potato potyvirus particles. *J. Mol. Biol.* 357:1–8
113. Teycheney PY, Marais A, Svanella-Dumas L, Dulucq MJ, Candresse T. 2005. Molecular characterization of banana virus X (BVX), a novel member of the *Flexiviridae* family. *Arch. Virol.* 150:1715–27
114. Tollin T, Wilson HR. 1988. Particle structure. See Ref. 86, 4:51–83
115. Urcuqui-Inchima S, Haenni AL, Bernardi F. 2001. Potyvirus proteins: a wealth of functions. *Virus Res.* 74:157–75
116. Van Dijk P, Veerbek M, Bos L. 1991. Mite-borne viruses isolates from cultivated *Allium* species and their classification into two new rymoviruses in the family *Potyviridae*. *Neth. J. Plant Pathol.* 97:381–99
117. Van Dijk P, Van der Vlugt RAA. 1994. New mite-borne viruses isolated from rakkyo, shallot and wild leek species. *Eur. J. Plant Pathol.* 100:269–77
118. Verchot J, Angell SM, Baulcombe DC. 1998. In vivo translation of the triple gene block of potato virus X requires two subgenomic mRNAs. *J. Virol.* 72:8316–20
119. Verchot-Lubicz J. 2005. A new cell-to-cell transport model for potexviruses. *Mol. Plant Microbe Interact.* 18:283–90
120. Vishnichenko VK, Stelmashchuk VY, Zavriev SK. 2002. The 42K protein of the Shallot virus X participates in formation of virus particles. *Mol. Biol.* 36:1080–84
121. Vives MC, Galipienso L, Navarro L, Moreno P, Guerri J. 2001. The nucleotide sequence and menome organisation of citrus leaf blotch virus: candidate type species for a new virus genus. *Virology* 287:225–33
122. Voinnet O, Ledere C, Baulcombe DC. 2000. A viral movement protein prevents spread of the gene silencing signal in *Nicotiana benthamiana*. *Cell* 103:157–67
123. Wetter C, Milne RG. 1981. Carlaviruses. See Ref. 64a, pp. 696–730
124. Xie J, Wei D, Jiang D, Fu Y, Li G, et al. 2006. Characterization of debilitation-associated mycovirus infecting the plant-pathogenic fungus *Sclerotinia sclerotiorum*. *J. Gen. Virol.* 87:241–49
125. Yaegashi H, Takahashi T, Isogai M, Kobori T, Ohki S, Yoshikawa N. 2007. Apple chlorotic leaf spot virus 50 kDa movement protein acts as a suppressor of systemic silencing without interfering with local silencing in *Nicotiana benthamiana*. *J. Gen. Virol.* 88:316–24
126. Zhang YP, Uyemoto JK, Kirkpatrick BC. 1998b. Analysis of double stranded RNAs from cherry trees with stem pitting in California. *Plant Dis.* 82:871–74
127. Zhou SZH, Dell’Orco M, Saldarelli P, Turturo C, Minafra A, et al. 2006. Identification of an RNA silencing suppressor in the genome of *Grapevine virus A*. *J. Gen. Virol.* 87:2387–95



# Contents

Tell Me Again What It Is That You Do <i>R. James Cook</i> .....	1
Noel T. Keen—Pioneer Leader in Molecular Plant Pathology <i>Alan Collmer and Scott Gold</i> .....	25
Structure and Function of Resistance Proteins in Solanaceous Plants <i>Gerben van Ooijen, Harrold A. van den Burg, Ben J.C. Cornelissen, and Frank L. W. Takken</i> .....	43
Family <i>Flexiviridae</i> : A Case Study in Virion and Genome Plasticity <i>Giovanni P. Martelli, Michael J. Adams, Jan F. Kreuze, and Valerian V. Dolja</i> .....	73
Cell Wall-Associated Mechanisms of Disease Resistance and Susceptibility <i>Ralph Hüchelboven</i> .....	101
Genomic Insights into the Contribution of Phytopathogenic Bacterial Plasmids to the Evolutionary History of Their Hosts <i>George W. Sundin</i> .....	129
Identifying Microorganisms Involved in Specific Pathogen Suppression in Soil <i>James Borneman and J. Ole Becker</i> .....	153
Safety of Virus-Resistant Transgenic Plants Two Decades After Their Introduction: Lessons from Realistic Field Risk Assessment Studies <i>Marc Fuchs and Dennis Gonsalves</i> .....	173
Disease Cycle Approach to Plant Disease Prediction <i>Erick D. De Wolf and Scott A. Isard</i> .....	203
Virus-Induced Disease: Altering Host Physiology One Interaction at a Time <i>James N. Culver and Meenu S. Padmanabhan</i> .....	221
Bacteriophages for Plant Disease Control <i>J.B. Jones, L.E. Jackson, B. Balogh, A. Obradovic, F.B. Iriarte, and M. T. Momol</i> .....	245



Reniform in U.S. Cotton: When, Where, Why, and Some Remedies <i>A. Forest Robinson</i> .....	263
Flax Rust Resistance Gene Specificity is Based on Direct Resistance-Avirulence Protein Interactions <i>Jeffrey G. Ellis, Peter N. Dodds, and Gregory J. Lawrence</i> .....	289
Microarrays for Rapid Identification of Plant Viruses <i>Neil Boonham, Jenny Tomlinson, and Rick Mumford</i> .....	307
Transcript Profiling in Host-Pathogen Interactions <i>Roger P. Wise, Matthew J. Moscou, Adam J. Bogdanove,        and Steven A. Whitam</i> .....	329
The Epidemiology and Management of Seedborne Bacterial Diseases <i>Ronald Gitiatis and Ronald Walcott</i> .....	371
Elicitors, Effectors, and <i>R</i> Genes: The New Paradigm and a Lifetime Supply of Questions <i>Andrew F. Bent and David Mackey</i> .....	399
Magnaporthe as a Model for Understanding Host-Pathogen Interactions <i>Daniel J. Ebbole</i> .....	437
Challenges in Tropical Plant Nematology <i>Dirk De Waele and Annemie Elsen</i> .....	457

## Errata

An online log of corrections to *Annual Review of Phytopathology* articles may be found at <http://phyto.annualreviews.org/>