

First Draft of CVX Paper

alexa tyszka

(Manuscript last compiled on September 15, 2021.)

Abstract

Potexviruses (Alphaflexiviridae) are positive-sense single-stranded RNA viruses known to infect multiple species across plants, including species of Cactaceae. Cactus Virus X, Zygocactus Virus X, Schlumbergera Virus X, Pitaya Virus X, and Opuntia Virus X are five of the 48 currently described Potexvirus species, which all infect valuable ornamental and crop plants, often causing production losses. The taxonomic naming schemes often employ outdated plant name synonyms, complicating taxonomic assignments. Also, the source of infections in cultivated plants is unclear, as is the distribution of and significance of infections in wild species of cacti. The lack of clarity is partly related to low sampling across the family. Here, we report results of original RNA-seq experiments and archived sequence deposits, aimed at detecting Potexviruses in cacti, assembling whole genomes, estimating their phylogenetic relationships, and delimiting viral species. The data suggests novel modes of transmission, based on expression analyses across tissues, particularly pollen. We also perform molecular evolutionary analyses to detect genomic regions under different modes of selection. Finally, we examine and discuss the implications of our analyses for the taxonomy of Potexviruses across cacti.

Introduction

1. Introduce CVX, History of problem, previous work and diversity of system The plant virome represents a fundamentally complex and intertwined evolutionary interaction between eukaryotic host and viral vector (Delwart [2007]). This coincides with the rise of faster, cheaper, high-throughput environmental metagenomic techniques (Delwart [2007], Lefeuvre et al. [2019], Schulz et al. [2017]): greenhouse-raised and lab-grown organisms that are commonly analyzed in experiments actually represent a small fraction of living diversity. Metagenomic studies aim to remedy this by sampling hundreds of thousands of genomes, and have vastly expanded both the cellular tree of life (Schulz et al. [2017], Hug et al. [2016]) and the viral tree of life (Gregory et al. [2019], Lefeuvre et al. [2019], Shi et al. [2016]). However, referring to viruses as a unified whole is not entirely correct or representative of the vast morphological diversity and innovative strategies present within the viral realms, many of which infect plants (Delwart [2007], Lefeuvre et al. [2019]). Metagenomic analysis of plants, particularly understudied or non-crop plants have both enriched our genetic knowledge of plants and uncovered novel insights on viral evolution, adaptation, and transmission. A careful study of the plant virome provides a view into biological realities that are not presently reflected in the literature.

The study of plant viruses often suffers from poor nomenclature due to the erroneous practice of naming a virus based on hosts. The term "plant virus", for example, implies that plant infection is monophyletic rather than homoplastic (Lefeuvre et al. [2019]). There is strong evidence to suggest that viruses frequently have "jumped" from fungal or invertebrate hosts to their associated plants (Lefeuvre et al. [2019]). Many agricultural plant viruses are named for the common name of a plant, for example, Pitaya Virus X. The common names "pitaya" and "dragonfruit" can both refer to two separate genera within Echinocereaceae (?): *Selenicereus* (which has 31 species) (Korotkova et al. [2017], Guerrero et al. [2019]) and *Stenocereus*, which has 23 putative species (Guerrero et al. [2019]). Further, the implication that one viral species exclusively infects all plants with the common name "pitaya" is incorrect since there are reports of multiple Potexviruses infecting the same host plant (Li* 2014). These problems persist throughout Potexvirus and are especially prominent with regards to cactus-infecting Potexviruses, underlining the necessity of a more informed taxonomy.

2. study system

Cactus Virus X, Zygocactus Virus X, Schlumbergera Virus X, Pitaya Virus X, and Opuntia Virus X are all Potexviruses (family Alphaflexiviridae) that are grouped broadly by their infections of certain cacti: *Selenicereus undatus* and *S. polyrhizus* (?Peng et al. [2016]); *Opuntia* spp. especially tuna (Koenig et al. [2004], Duarte et al. [2008]) and *monacantha* (Attathom et al. [1978] Sammons 1961 Duarte 2008); *Schlumbergera* (previously *Zygocactus*) *truncata* and *S. bridgesii* (Duarte et al. [2008], Koenig et al. [2004]), *Parodia* (previously *Notocactus*) *leninghausii* (Park et al. [2018]), *Echinopsis chamaecereus* f. *cristata*, *E. pectinatus* f. *cristata*, *E. jusbertainii*, and *E. macrogona* (Maliarenko and Mudrak [2013]); *Mammillaria elongata* f. *cristata* (Maliarenko and Mudrak [2013]); and multiple other species within many genera (Evallo et al. [2021]). Of these viruses, only CVX has been reported on wild *Ferocactus cylindraceus* (previously *Ferrocactus acanthodes*) (Attathom et al. [1978]) although this report predates DNA records confirming viral identity. Additionally, although they are originally found on cacti, the viruses are frequently manipulated with serological experiments and have been found to produce lesions (which indicate infection) on: *Chenopodium murale* L. (Maliarenko and Mudrak [2013]) and *C. quinoa* (Attathom [1978], Attathom et al. [1978], ?; *Nicotiana glauca* Link. et al. Otto (Maliarenko and Mudrak [2013]); Four species of *Amaranthaceae* (Attathom [1978]); *Escobaria vivipara* (Attathom [1978]); and other cactaceae (Attathom [1978]). All cactus-infecting potexviruses consist of roughly 6,600 bp of positive-sense single-stranded RNA. They have similar rod-shaped filamentous virions and share the same division of five primary open reading frames (ORFs): Replicase (Rep), Triple gene block (TGB), Coat protein (CP), coded in the 5' direction as well as two smaller overlapping ORFs coded in the 3' direction: ORF6 and ORF7 (Evallo et al. [2021], Liou et al. [2004], Martelli et al. [2007]).

They are closely related to other Potexviruses such as Alternanthera Mosaic Virus and Papaya Mosaic Virus (Martelli et al. [2007], Park et al. [2018], Liou et al. [2004]). These viruses produce a wide range of symptomatic and damaging infections in cacti. Reports of symptomatic plants range from 5.5 percent of wild *Ferocactus cylindraceus* (Attathom et al. [1978]) up to 44 percent of crop plants on Hainan Island, China (Peng et al. [2016]). However, many infected plants do not show external signs of viral infection (Liou et al. [2004], ?). The most commonly recognized symptoms of disease are mosaic, mottling, stunted growth and distortion (Maliarenko and Mudrak [2013], Peng et al. [2016], Attathom et al. [1978]). It is unclear what the method of transmission from infected plant to new host is. Some reports specify that cactus-infecting Potexviruses can only be transmitted through grafting (Duarte et al. [2008], Martelli et al. [2007]) but most agree that transmission can occur through other mechanical contact such as sap inoculation (Liou et al. [2004], Maliarenko and Mudrak [2013], Park et al. [2018]) and external tissue contact. Grafting is a primary means of propagation among crop cacti (Park et al. [2018]), and *Selenicereus* is a commonly chosen graft stock. However, there are reports of other members within *Alphaflexiviridae* transmitting via insect and seed vectors (Martelli et al. [2007]), and pre-DNA studies suggest that CVX may transmit via pollen in the wild (Attathom et al. [1978]).

3. problems in existing research and knowledge gaps

4. what is gained from studying this? Knowledge about cactus-infecting potexviruses contribute to a growing yet biased study of plant viruses. The pursuit of wild cactus-infecting potexviruses serves to expand our evolutionary knowledge of viral evolution, host selection, and transmission mechanics.

5. Specific objectives and questions/summary In this study we present the largest to date phylogeny of cactus-infecting Potexviruses. We attempt to use this expanded phylogeny to answer relevant questions about Potexvirus evolutionary relationships as well as revisiting the utility of decades-old taxonomy in current virus research.

Materials and Methods

Plant Material Collection: Pistils, pollen, and styles from *Schlumbergera truncata* individuals were collected from B. Igie’s personal collection, which contained specimens displaying symptoms typical to Potexvirus infection (Chessin [1972]): frail, reddened cladodes, and poor growth. The plants were initially obtained from various sources, then kept in a shared growing condition.

RNA Extraction and Sequencing: Tissues were collected and immediately submerged in 1.5 ml of RNeasy lysis solution (Qiagen). Submerged samples were held at room temperature for thirty minutes and then stored at -80° C.

NCBI Data Collection and Compilation: We collected publicly available genomes, complete proteins, gene annotations, and available metadata from Potexviruses (NCBI:txid12176) (NCBI: www.ncbi.nlm.nih.gov/, accession numbers provided in Supplemental Data). The untranslated regions (UTRs) were trimmed from the sequences to provide consistency.

We also searched the NCBI Sequence Read Archive (SRA) database (www.ncbi.nlm.nih.gov/sra) for RNA-sequencing (RNA-seq) data within Caryophyllales (NCBI:txid3524) that had been sequenced using the Illumina library sequencing platform. For each identified SRA run accession (SRR), any viral RNA that matched sample cactus-infecting Potexvirus RNA (accession numbers provided in Supplemental Data) was identified, extracted, and assembled using the kakapo 0.7.3-dev pipeline (<http://flightless.one>) with Kraken2 viral filters disabled. The .sam files produced through kakapo were loaded through Geneious 11.1.5 along with the *Schlumbergera* reads. These sequences were annotated using the Geneious 11.1.5 "Find ORFs" function.

The complete dataset comprises: 37 existing Potexvirus genomes and proteins, 4 new viral sequences located within original *Schlumbergera truncata* RNA-seq data, and 52 viral sequences found within NCBI Caryophyllales RNA-seq data.

Alignment and Phylogenetic Analyses Sequence alignments were performed through MAFFT v7.429 (Katoh [2002]) using the full dataset. The aligned sequences were divided by ORF using the annotations to produce five partial sequence alignments corresponding to each ORF to accompany the full-sequence alignment. The individual proteins were exported to .FASTA files, then gaps at the start of the sequence and stop codons were removed manually. Phylogenetic relationships and bootstrap values were inferred using IQtree v1.6.12 (Nguyen et al. [2015]), ModelFinder (Kalyaanamoorthy et al. [2017]), and UFBoot (Hoang et al. [2018]) for both the individual gene/protein alignments and the full sequence alignment. Trees were visualized in R version 4.0.3 using ggtree v2.4.2 (?). Host information was obtained through reported metadata and mapped onto the phylogeny. Species groupings were determined using the existing species boundaries when compared to the phylogenetic branch lengths within the Potexvirus genus. This was generally consistent with most recent branch lengths over 0.1 subs/site and this value was therefore used as a cutoff. Pairwise distance analysis was conducted on the sequence alignments in R using the ape v5.5 dist.dna() function with a raw model. For each defined clade, nonzero pairwise distances between each possible combination of tips was averaged. Expanded phylogenetic trees and individual gene/protein trees are available in the Supplementary Data.

Results

Phylogenetic analysis The phylogenetic tree places the new viral sequences from *Schlumbergera* and *Selenicereus* near existing viral species within Potexvirus (Figure 1). The *S. truncata* samples were located within the Cactus Virus X clade and appear to represent the only known discovery of Cactus Virus X on members of *Schlumbergera*. The publicly available data which was collected from NCBI produced 52 new viral sequences which were dispersed among viral species. These samples were exclusively representative

of *Selenicereus undatus* and *Selenicereus polyrhizus* hosts. The Cactus Virus X species clade is expanded by a factor of 8. The only putative species that was not expanded by either the Schlumbergera data or the *Selenicereus* data was *Opuntia Virus X*, which appears to be an outgroup to the other cactus-infecting Potexviruses.

Divisions within species The five species within the phylogeny appear to be generally characterized by long (*) branch lengths separating clusters of closely related (short branch lengths) tips. Cactus Virus X, as a putative species, appears to have two time-separated evolutionary distant sets of tips. These have been marked as different colors in Figure 1. The other group that has outliers is *Zygocactus Virus X*, which appears to have a longer evolutionary distance between the tip "Zygocactus virus X KM288845.1" and the rest of the samples either identified as *Zygocactus Virus X* or identified as closely related to ZyVX. The relationship and its bootstrap value have both been marked with different colors in Figure 1.

Pairwise distances between species were calculated for six groups, with a value of 0 indicating identical sequences and a value of 1 indicating completely divergent sequences (Figure 2). The average nonzero pairwise distance between all included potexvirus tips was 0.256, maximum = 0.4925. When the outgroup (including *Plantago asiatica* MV, *Alternanthera* MV, *Papaya* MV, etc.) was excluded from pairwise analysis, the average nonzero pairwise distance value was 0.177, maximum = 0.326. When these cactus-infecting potexviruses were subdivided into six groups of relatively recent diversification, the average nonzero pairwise distance for full-genome sequences was always above 0.015. For the genes RNA-dependent RNA polymerase (RdRp) and Coat protein (CP), which the ICTV recommends be analyzed for species delimitation, the average nonzero pairwise distance was always above 0.02 (Figure 2). This correlates to roughly greater than 97.5 and 98 nucleotide identity.

Gene phylogenies The assembled phylogenies for each of the five genes are available in their fully annotated versions in the Supplemental Data. The gene trees generally supported the species groupings of the full-genome tree and do not change the interpretation of the "main" tree.

Host range The host from attached metadata for each species is noted in Figure 1. There are two unique samples collected from NCBI that report non-plant hosts of cactus-infecting potexviruses: "Cactus virus X SCM51431", and "Mytcor Virus 1 MG210801". These have the hosts Diptera and Bivalve, respectively.

Discussion

References

- S. Attathom. Identification and Characterization of a Potexvirus from California Barrel Cactus. *Phytopathology*, 68(10):1401, 1978. ISSN 0031949X. doi: 10.1094/Phyto-68-1401. URL http://www.apsnet.org/publications/phytopathology/backissues/Documents/1978Abstracts/Phyto68_1401.htm.
- S. Attathom, L. G. Weathers, and D. J. Gumpf. Occurrence and distribution of a virus induced disease of barrel cactus in California. *Plant disease reporter*, 1978. ISSN 0032-0811. URL https://scholar.google.com/scholar_lookup?title=Occurrence+and+distribution+of+a+virus+induced+disease+of+barrel+cactus+in+California&author=Attathom%2C+S.&publication_year=1978.
- M. Chessin. Distribution of Cactus Viruses in Wild Plants. *Phytopathology*, 62(1):97, 1972. ISSN 0031949X. doi: 10.1094/Phyto-62-97. URL http://www.apsnet.org/publications/phytopathology/backissues/Documents/1972Abstracts/Phyto62_97.htm.
- Eric L. Delwart. Viral metagenomics. *Reviews in Medical Virology*, 17(2):115–131, March 2007. ISSN 10529276, 10991654. doi: 10.1002/rmv.532. URL <https://onlinelibrary.wiley.com/doi/10.1002/rmv.532>.
- L M L Duarte, M A V Alexandre, E B Rivas, R Harakava, S R Galleti, and M M Barradas. POTEXVIRUS DIVERSITY IN CACTACEAE FROM SÃO PAULO STATE IN BRAZIL. *Journal of Plant Pathology*, page 9, 2008.
- Edzel Evallo, John Darby Taguam, and Mark Angelo Balendres. A brief review of plant diseases caused by Cactus virus X. *Crop Protection*, 143:105566, May 2021. ISSN 02612194. doi: 10.1016/j.cropro.2021.105566. URL <https://linkinghub.elsevier.com/retrieve/pii/S0261219421000363>.
- Ann C. Gregory, Ahmed A. Zayed, Nádia Conceição-Neto, Ben Temperton, Ben Bolduc, Adriana Alberti, Mathieu Ardyna, Ksenia Arkhipova, Margaux Carmichael, Corinne Cruaud, Céline Dimier, Guillermo Domínguez-Huerta, Joannie Ferland, Stefanie Kandels, Yunxiao Liu, Claudie Marec, Stéphane Pesant, Marc Picheral, Sergey Pisarev, Julie Poulain, Jean-Éric Tremblay, Dean Vik, Silvia G. Acinas, Marcel Babin, Peer Bork, Emmanuel Boss, Chris Bowler, Guy Cochrane, Colomban de Vargas, Michael Follows, Gabriel Gorsky, Nigel Grimsley, Lionel Guidi, Pascal Hingamp, Daniele Iudicone, Olivier Jaillon, Stefanie Kandels-Lewis, Lee Karp-Boss, Eric Karsenti, Fabrice Not, Hiroyuki Ogata, Stéphane Pesant, Nicole Poulton, Jeroen Raes, Christian Sardet, Sabrina Speich, Lars Stemmann, Matthew B. Sullivan, Shinichi Sunagawa, Patrick Wincker, Marcel Babin, Chris Bowler, Alexander I. Culley, Colomban de Vargas, Bas E. Dutilh, Daniele Iudicone, Lee Karp-Boss, Simon Roux, Shinichi Sunagawa, Patrick Wincker, and Matthew B. Sullivan. Marine DNA Viral Macro- and Microdiversity from Pole to Pole. *Cell*, 177(5):1109–1123.e14, May 2019. ISSN 0092-8674. doi: 10.1016/j.cell.2019.03.040. URL <https://www.sciencedirect.com/science/article/pii/S0092867419303411>.
- Pablo C Guerrero, Lucas C Majure, Amelia Cornejo-Romero, and Tania Hernández-Hernández. Phylogenetic Relationships and Evolutionary Trends in the Cactus Family. *Journal of Heredity*, 110(1):4–21,

- January 2019. ISSN 0022-1503, 1465-7333. doi: 10.1093/jhered/esy064. URL <https://academic.oup.com/jhered/article/110/1/4/5205122>.
- Diep Thi Hoang, Olga Chernomor, Arndt von Haeseler, Bui Quang Minh, and Le Sy Vinh. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, 35(2):518–522, February 2018. ISSN 0737-4038, 1537-1719. doi: 10.1093/molbev/msx281. URL <https://academic.oup.com/mbe/article/35/2/518/4565479>.
- Laura A. Hug, Brett J. Baker, Karthik Anantharaman, Christopher T. Brown, Alexander J. Probst, Cindy J. Castelle, Cristina N. Butterfield, Alex W. Hernsdorf, Yuki Amano, Kotaro Ise, Yohey Suzuki, Natasha Dudek, David A. Relman, Kari M. Finstad, Ronald Amundson, Brian C. Thomas, and Jillian F. Banfield. A new view of the tree of life. *Nature Microbiology*, 1(5):16048, May 2016. ISSN 2058-5276. doi: 10.1038/nmicrobiol.2016.48. URL <http://www.nature.com/articles/nmicrobiol201648>.
- Subha Kalyaanamoorthy, Bui Quang Minh, Thomas K F Wong, Arndt von Haeseler, and Lars S Jermiin. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*, 14(6): 587–589, June 2017. ISSN 1548-7091, 1548-7105. doi: 10.1038/nmeth.4285. URL <http://www.nature.com/articles/nmeth.4285>.
- K. Katoh. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, 30(14):3059–3066, July 2002. ISSN 13624962. doi: 10.1093/nar/gkf436. URL <https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkf436>.
- R. Koenig, H. Aust, J. Schiemann, C. W. A. Pleij, S. Loss, and W. Burgermeister. Molecular characterisation of potexviruses isolated from three different genera in the family Cactaceae. *Archives of Virology*, 149(5): 903–914, May 2004. ISSN 0304-8608, 1432-8798. doi: 10.1007/s00705-003-0268-5. URL <http://link.springer.com/10.1007/s00705-003-0268-5>.
- Nadja Korotkova, Thomas Borsch, and Salvador Arias. A phylogenetic framework for the Hylocereeae (Cactaceae) and implications for the circumscription of the genera. *Phytotaxa*, 327(1):1, November 2017. ISSN 1179-3163, 1179-3155. doi: 10.11646/phytotaxa.327.1.1. URL <https://biotaxa.org/Phytotaxa/article/view/phytotaxa.327.1.1>.
- Pierre Lefeuvre, Darren P. Martin, Santiago F. Elena, Dionne N. Shepherd, Philippe Roumagnac, and Arvind Varsani. Evolution and ecology of plant viruses. *Nature Reviews Microbiology*, 17(10):632–644, October 2019. ISSN 1740-1526, 1740-1534. doi: 10.1038/s41579-019-0232-3. URL <http://www.nature.com/articles/s41579-019-0232-3>.
- M. R. Liou, Y. R. Chen, and R. F. Liou. Complete nucleotide sequence and genome organization of a Cactus virus X strain from *Hylocereus undatus* (Cactaceae). *Archives of Virology*, 149(5):1037–1043, May 2004. ISSN 0304-8608, 1432-8798. doi: 10.1007/s00705-003-0251-1. URL <http://link.springer.com/10.1007/s00705-003-0251-1>.
- Valentyna M. Maliarenko and Tatyana P. Mudrak. Cactus viruses in fasciated plants. *Biologija*, 59(2), November 2013. ISSN 2029-0578, 1392-0146. doi: 10.6001/biologija.v59i2.2754. URL <http://lmaleidykla.lt/ojs/index.php/biologija/article/view/2754>.
- Giovanni P. Martelli, Michael J. Adams, Jan F. Kreuze, and Valerian V. Dolja. Family *Flexiviridae* : A Case Study in Virion and Genome Plasticity. *Annual Review of Phytopathology*, 45(1):73–100, September 2007. ISSN 0066-4286, 1545-2107. doi: 10.1146/annurev.phyto.45.062806.094401. URL <http://www.annualreviews.org/doi/10.1146/annurev.phyto.45.062806.094401>.
- Lam-Tung Nguyen, Heiko A. Schmidt, Arndt von Haeseler, and Bui Quang Minh. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution*, 32(1):268–274, January 2015. ISSN 1537-1719, 0737-4038. doi: 10.1093/molbev/msu300. URL <https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msu300>.
- Chung Hwa Park, Eun Gyeong Song, and Ki Hyun Ryu. Detection of Co-Infection of *Notocactus leninghausii* f. *cristatus* with Six Virus Species in South Korea. *The Plant Pathology Journal*, 34(1):65–70, February 2018. ISSN 1598-2254, 2093-9280. doi: 10.5423/PPJ.NT.08.2017.0187. URL <http://ppjonline.org/journal/view.php?doi=10.5423/PPJ.NT.08.2017.0187>.
- C. Peng, N. T. Yu, Z. W. Luo, H. Y. Fan, F. He, X. H. Li, Z. L. Zhang, and Z. X. Liu. Molecular Identification of Cactus virus X Infecting *Hylocereus polyrhizus* (Cactaceae) in Hainan Island, China. *Plant Disease*, 100(9):1956, September 2016. ISSN 0191-2917, 1943-7692. doi: 10.1094/PDIS-01-16-0048-PDN. URL <https://apsjournals.apsnet.org/doi/10.1094/PDIS-01-16-0048-PDN>.
- Frederik Schulz, Emiley A. Eloie-Fadrosch, Robert M. Bowers, Jessica Jarett, Torben Nielsen, Natalia N. Ivanova, Nikos C. Kyrpides, and Tanja Woyke. Towards a balanced view of the bacterial tree of life. *Microbiome*, 5(1):140, December 2017. ISSN 2049-2618. doi: 10.1186/s40168-017-0360-9. URL <https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-017-0360-9>.
- Mang Shi, Xian-Dan Lin, Jun-Hua Tian, Liang-Jun Chen, Xiao Chen, Ci-Xiu Li, Xin-Cheng Qin, Jun Li, Jian-Ping Cao, John-Sebastian Eden, Jan Buchmann, Wen Wang, Jianguo Xu, Edward C. Holmes, and Yong-Zhen Zhang. Redefining the invertebrate RNA virosphere. *Nature*, 540(7634):539–543, December 2016. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature20167. URL <http://www.nature.com/articles/nature20167>.