

Bus Charging Schedule Simulated Annealing with MILP Constraints

Alexander Brown

March 26, 2023

Contents

1 Preliminaries	1
1.1 Mixed Integer Linear Program	1
1.2 Fuzzy Sets Theory	1
1.2.1 Fuzzy Sets	1
1.2.2 Fuzzy Arithmetic	2
1.2.3 Comparing Fuzzy Numbers	2
1.3 Fully Fuzzy Linear Programming (FFLP)	3

1 Preliminaries

1.1 Mixed Integer Linear Program

A mixed integer linear programming (MILP) problem is a class of constrained optimization in which one seeks to find a set of continuous or integer values that maximizes or minimizes an objective function while satisfying a set of constraints [2]. Given an objective function J , decision variables (i.e. variables of optimization) $x_j \in \mathbb{R}$ and $y_k \in \mathbb{Z}^+$, and input parameters $c_j, d_k, a_{ij}, g_{ik}, b_i \in \mathbb{R}$, a MILP has the mathematical structure represented in Equation 1 [2].

$$\begin{aligned} \text{Maximize } & J = \sum_j c_j x_j + \sum_k d_k y_k \\ \text{subject to } & \sum_j a_{ij} x_j + \sum_k g_{ik} y_k \leq b_i \quad (i = 1, 2, \dots, m) \\ & x_j \geq 0 \quad (j = 1, 2, \dots, n) \\ & y_k \in \mathbb{Z}^+ \quad (k = 1, 2, \dots, n) \end{aligned} \tag{1}$$

This formulation of the MILP is also referred to as “crisp”. By this it is meant that each variable in the formulation acts as an injective mapping to its number representation. In other words, no values on the formulation are fuzzy [5].

1.2 Fuzzy Sets Theory

This section introduces the notion of fuzzy numbers and some basic definitions. Concepts from this section are pulled from [8, 3, 7, 1].

1.2.1 Fuzzy Sets

Let’s begin with what a fuzzy number is not. A classical (crisp) set is defined as a collection of elements $x \in X$. Crisp sets are binary, either an element either belongs in the set, or it does not [8]. For a fuzzy set, what is known as the characteristic functions applies various degrees of membership for elements of a given set.

[8].

Definition 1.1 *Let X be a collection of objects (often called the universe of discourse [1]). If X is denoted generically by x , then a fuzzy set \tilde{A} in X is a set of ordered pairs as shown in Equation 2.*

$$\tilde{A} = \{(x, \mu_{\tilde{A}}(x)) | x \in X\} \quad (2)$$

$\mu_{\tilde{A}}$ is called the membership function where $\mu_{\tilde{A}}$ is the mapping $\mu_{\tilde{A}} : X \rightarrow [0, 1]$; which assigns a real number to the interval $[0, 1]$. The value of $\mu_{\tilde{A}}$ represents the degree of membership of x in \tilde{A} .

This paper will use fuzzy sets defined on the real numbers \mathbb{R} . The membership function describes the shape of the fuzzy number. As an example, consider the following definition.

Definition 1.2 A fuzzy number that is represented by $\tilde{A} = (a, b, c)$ is said to be triangular if its membership function is defined as Equation 3.

$$\mu_{\tilde{A}}(x) = \begin{cases} \frac{(x-a)}{(b-a)} & a \leq x \leq b \\ \frac{(d-x)}{(d-b)} & c \leq x \leq d \\ 0 & \text{else} \end{cases} \quad (3)$$

Definition 1.3 The fuzzy set \tilde{A} in \mathbb{R} is normal if $\max_x \mu_{\tilde{A}}(x) = 1$.

Definition 1.4 A fuzzy set \tilde{A} in \mathbb{R} is convex if and only if the membership function of \tilde{A} satisfies the inequality

$$\mu_{\tilde{A}}[\beta x_1 + (1 - \beta)x_2] \geq \min[\mu_{\tilde{A}}(x_1), \mu_{\tilde{A}}(x_2)] \quad \forall x_1, x_2 \in \mathbb{R} \quad \beta \in [0, 1]$$

Definition 1.5 A fuzzy number is a normal convex fuzzy set in \mathbb{R} .

Definition 1.6 The triangular fuzzy number \tilde{A} is nonnegative $\iff a \geq 0$.

1.2.2 Fuzzy Arithmetic

If two triangular fuzzy numbers $\tilde{a}_1 = \{a_1, a_2, a_3\}$ and $\tilde{b}_1 = \{b_1, b_2, b_3\}$ are nonnegative then the following operations are defined

$$\begin{aligned} \tilde{a} \oplus \tilde{b} &= (a_1 + b_1, a_2 + b_2, a_3 + b_3) \\ \tilde{a} \ominus \tilde{b} &= (a_1 + b_3, a_2 + b_2, a_3 + b_1) \end{aligned} \quad (4)$$

1.2.3 Comparing Fuzzy Numbers

Fuzzy numbers do not directly provide a method of ordering nor do they always provide an ordered set like real numbers [1]. There are multiple methods for ordering fuzzy numbers, each coming with advantages and disadvantages. Different properties have been applied to justify comparison of fuzzy numbers, such as: preference, rationality, and robustness [4, 1, 5]. Commonly, including in this work, the First index of Yager [6] is used (Equation 5).

$$\Re(\tilde{A}) = \frac{\sum_i a_i}{|\tilde{A}|} \quad (5)$$

where $|\cdot|$ represents the cardinality of the fuzzy number. In words, Equation 5 is merely the average of the values in the fuzzy number.

1.3 Fully Fuzzy Linear Programming (FFLP)