

b -flavour tagging in pp collisions

Alex Birnkraut on behalf of the LHCb collaboration

Basics

Introduction

Measurements of flavour oscillations and time-dependent CP asymmetries in neutral B meson systems require knowledge of the b quark flavour at production. This identification is performed by the Flavour Tagging (FT). [1,2]

Two independent classes of algorithms

- same side taggers (SS)**


use charged particles created in the fragmentation process of the b quark of the signal B meson

- kaon for B_s^0 → SS kaon / SS kaon nnet
- pion for B^0 → SS pion
- proton for B^0 → SS proton

- opposite side taggers (OS)**

exploit the non-signal b quark of the initial $b\bar{b}$ pair

- overall charge of the secondary vertex (SV) → OS vertex charge
- lepton from semi-leptonic b hadron decays → OS muon / OS electron
- kaon from the $b \rightarrow c \rightarrow s$ decay chain → OS kaon
- D meson from the $b \rightarrow c$ decay chain → OS charm (New!)



Each tagger provides a decision d on the initial flavour ("tag") and a probability to be wrong, η .

Flavour Tagging characteristics

- mistag**

fraction of events with a wrong tagging decision

$$\omega = \frac{N_{\text{wrong}}}{N_{\text{right}} + N_{\text{wrong}}}$$

- tagging efficiency**

fraction of events with a tagging decision

$$\epsilon_{\text{tag}} = \frac{N_{\text{right}} + N_{\text{wrong}}}{N_{\text{all}}}$$

- effective tagging efficiency**

represents the statistical reduction factor of a sample in a tagged analysis

$$\epsilon_{\text{eff}} = \epsilon_{\text{tag}} (1 - 2\omega)^2$$

Calibration

Mistag calibration


$$\omega(\eta) = p_0 + p_1(\eta - \langle \eta \rangle)$$

↑ calibrated ev-by-ev mistag ↑ estimated ev-by-ev mistag ↑ mean estimated mistag

Several flavour-specific decay channels are used

- $B^+ \rightarrow J/\psi K^+$, $B^+ \rightarrow D^0 \pi^+$
charged channels: extract ω by comparing tag decision with charge of the final state
- $B^0 \rightarrow J/\psi K^{*0}$, $B^0 \rightarrow D^{*-} \mu^+ \nu_\mu$, $B_s^0 \rightarrow D_s^- \pi^+$, ...
neutral channels: full time-dependent analysis to extract ω from the mixing asymmetry

$$\mathcal{A}_{\text{mix}}(t) \propto (1 - 2\omega) \cos(\Delta m_{d/s} t)$$



Flavour Tagging in Run I

Usage in analyses


- one calibration per tagger valid for all channels
- systematic uncertainties from
 - calibration methods
 - results in different control channels
- "ad-hoc" calibration using best-suited control channels for analyses dominated by FT uncertainty

Highlights of flavour-tagged measurements


- Measurements of ϕ_s**

Decay mode	$\epsilon_{\text{eff}, 2011}$	$\epsilon_{\text{eff}, \text{Run I}}$
$B_s^0 \rightarrow J/\psi K^+ K^-$	3.13 [3]	3.73 [4]
$\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$	2.43 [5]	3.89 [6]
$\bar{B}_s^0 \rightarrow D_s^+ D_s^-$	-	5.33 [7]

- newest analyses profited from:
 - including SS kaon nnet tagger
 - re-optimisation of OS algorithms




- CP violation in $B_s^0 \rightarrow D_s^\mp K^\pm$**



- analysis on 2011 data: $\epsilon_{\text{eff}} = 5.07\%$
- SS kaon nnet adds more than 1.3 % to ϵ_{eff} [8]

- CP violation in $B^0 \rightarrow J/\psi K_s^0$ ($\sin 2\beta$)**


- analysis on 2011 data: $\epsilon_{\text{eff}} = 2.38\%$ [9]
- full Run I analysis: $\epsilon_{\text{eff}} = 3.02\%$ [10]
 - SS pion tagger adds more than 0.376 % to ϵ_{eff}



- precision analysis → "ad-hoc" FT calibration
 - OS algorithms calibrated with $B^+ \rightarrow J/\psi K^+$
 - SS pion calibrated with $B^0 \rightarrow J/\psi K^{*0}$

- CP violation in $B_s^0 \rightarrow J/\psi K_s^0$**

- not possible to exclude B^0 events in selection



- B_s^0 events: $\epsilon_{\text{eff}} = 4.00\%$ [11]
- B^0 events: $\epsilon_{\text{eff}} = 2.62\%$ [11]
 - small tagging power of SS kaon for B^0 :
 - same-side protons misidentified as kaons
 - kaons from same-side K^* (892)
 - kaons have opposite charge for B^0 : tagging decision has to be inverted

Developments

OS charm tagger (preliminary)

- reconstruct $D^0/D^\pm/D^*$ decays related to OS b decay

Decay mode	Relative ϵ_{tag}	Relative ϵ_{eff}
$D^0 \rightarrow K^- \pi^+$	10.0 %	24.0 %
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	5.9 %	8.4 %
$D^+ \rightarrow K^- \pi^+ \pi^+$	10.3 %	2.6 %
$D^0, D^+ \rightarrow K^- \pi^+ X$	69.7 %	61.5 %
$D^0, D^+ \rightarrow K^- e^+ X$	0.5 %	0.2 %
$D^0, D^+ \rightarrow K^- \mu^+ X$	3.4 %	0.3 %
$\Lambda_c^+ \rightarrow p^+ K^- \pi^+$	0.2 %	2.4 %


- one boosted decision tree (BDT) for each mode [12]
- clean measure of B meson flavour (low mistag)
- stand-alone tagging power of $\epsilon_{\text{eff}} = 0.30\%$ to 0.40%

SS pion calibration

- calibration performed with $B^0 \rightarrow J/\psi K^{*0}$
- full evaluation of systematic uncertainties
- used for the first time in the measurements of
 - $\sin(2\beta)$ with $B^0 \rightarrow J/\psi K_s^0$
 - ⇒ $\epsilon_{\text{eff}}^{\text{SS}\pi} = 0.38\%$
 - $\sin(2\beta_{\text{eff}})$ with $B^0 \rightarrow J/\psi \pi^+ \pi^-$
 - ⇒ $\epsilon_{\text{eff}}^{\text{SS}\pi} = 0.54\%$

SS kaon tagging using neural nets (NN)

- basic idea: use two NN
 - first NN distinguishes between:
 - fragmentation tracks
 - ⇒ signal for SS kaon nnet
 - underlying event tracks



- second NN:
 - assigns final tag and mistag based on multiple candidates [13]
- SS kaon nnet tagger is a great success, compared to the previous cut-based SS kaon it gives
 - $B_s^0 \rightarrow D_s^- \pi^+$: 50 % relative improvement in ϵ_{eff}
 - $B_s^0 \rightarrow J/\psi \phi$: 41 % relative improvement in ϵ_{eff}

References

- [1] LHCb Collaboration, R. Aaij et. al., *Opposite-side flavour tagging of B mesons at the LHCb experiment*, Eur.Phys.J. C72 (2012) 2022
- [2] LHCb Collaboration, R. Aaij et. al., *Optimization and calibration of the same-side kaon tagging algorithm using hadronic B_s^0 decays in 2011 data*, LHCb-CONF-2012-033
- [3] LHCb Collaboration, R. Aaij et. al., *Measurement of CP violation and the B_s^0 meson decay width difference with $B_s^0 \rightarrow J/\psi K^+ K^-$ and $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Rev. D87 (2013) 11, 112010
- [4] LHCb Collaboration, R. Aaij et. al., *Precision measurement of CP violation in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays*, Phys.Rev.Lett. 114 (2015) 4, 041801
- [5] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP -violating phase ϕ_s in $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Lett. B713 (2012) 378-386
- [6] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP -violating phase ϕ_s in $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Lett. B736 (2014) 186-195
- [7] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP -violating phase ϕ_s in $\bar{B}_s^0 \rightarrow D_s^+ D_s^-$ decays*, Phys.Rev.Lett. 113 (2014) 21, 211801
- [8] LHCb Collaboration, R. Aaij et. al., *Measurement of CP asymmetry in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays*, JHEP 1411 (2014) 060
- [9] LHCb Collaboration, R. Aaij et. al., *Measurement of the time-dependent CP asymmetry in $B^0 \rightarrow J/\psi K_s^0$ decays*, Phys.Lett. B721 (2013) 24-31
- [10] LHCb Collaboration, R. Aaij et. al., *Measurement of CP violation in $B^0 \rightarrow J/\psi K_s^0$ decays*, Phys.Rev.Lett. 115 (2015) 3, 031601
- [11] LHCb Collaboration, R. Aaij et. al., *Measurement of the time-dependent CP asymmetries in $B_s^0 \rightarrow J/\psi K_s^0$* , JHEP 1506 (2015) 131
- [12] LHCb Collaboration, R. Aaij et. al., *B flavor tagging using reconstructed charm decays at the LHCb experiment*, LHCb-PAPER-2015.027
- [13] G. A. Krocker, *Development and calibration of a same side kaon tagging algorithm and measurement of the $B_s^0 - \bar{B}_s^0$ oscillation frequency Δm_s at the LHCb experiment*, PhD thesis, Heidelberg U., Sep. 2013, CERN-THESIS-2013-213