

***b*-flavour tagging in *pp* collisions**

Alex Birnkrout*

TU Dortmund

E-mail: a.birnkrout@cern.ch

In the system of neutral B^0 mesons CP -violating processes can be measured using time dependent analyses, as performed at the LHCb experiment. For such analyses the knowledge of the initial flavour of the mesons is mandatory. This information is provided by the Flavour Tagging, which exploits a variety of different algorithms. This article shows the good performance in different analyses which allowed high precision measurements and also the improvements which were made throughout Run I. Finally some very recent developments and new algorithms to tag the initial flavour are described.

*The European Physical Society Conference on High Energy Physics
22–29 July 2015
Vienna, Austria*

*Speaker.

indicates then indicates the initial flavour of the b quark. The OS kaon and the OS charm taggers use quite similar approaches by analysing the decay product of the decay chain $b \rightarrow c \rightarrow s$ or the decay product of the decay chain $b \rightarrow c$, respectively. The charge of the final state kaon in case of the OS kaon or the D meson in case of the OS charm tagger are used to get information about the initial b flavour.

In contrast to this method the OS vertex charge tagger does not use single tracks but a weighted charge of the whole secondary vertex to come to a tag decision. In order to do so, the secondary vertex is reconstructed first out of two tracks which have the highest probability to originate from the b hadron. After this, more tracks are added to the vertex and the weighted charge is calculated as

$$Q_{\text{vtx}} = \frac{\sum_i p_T^k(i) Q_i}{\sum_i p_T^k(i)} \quad (2.1)$$

The opposite side tagging is independent of the signal B meson as it only investigates the propagation of the opposite b quark.

2.2 Same side tagging

For the same side tagging one has to distinguish between B_d^0 and B_s^0 mesons as the accompanying quark is a d or a s quark, respectively. In case of a B_d^0 meson an additional d quark which can hadronise with an u quark into a pion emerges. Also pions from excited states as B^* and B^{**} have the same charge as pions from the direct fragmentation process with a B meson. If the signal B meson is a B_s^0 an kaon can be formed out of the additional s quark and an u quark. In a new development the fragmentation kaon is selected using neural nets. The so called SS kaon neural net tagger will be described more in detail in section 5.

2.3 Flavour Tagging characteristics

Each tagger provides for each candidate a tag d and the probability η that this tag is wrong. The tag can be 1 for a \bar{b} quark, -1 for a b quark and 0 if the tagger could not provide a tag decision. The mistag probability is defined to smaller then 0.5 as the tag could be flipped otherwise. This mistag is just a self-estimation of the tagger and the true mistag is defined as

$$\omega = \frac{N_{\text{wrong}}}{N_{\text{right}} + N_{\text{wrong}}} \quad (2.2)$$

where N_{wrong} is the number of wrong tagged candidates and N_{right} the number of correctly tagged candidates. Also the performance of the different taggers is described by their efficiencies

$$\epsilon_{\text{tag}} = \frac{N_{\text{right}} + N_{\text{wrong}}}{N_{\text{all}}} \quad (2.3)$$

with the number of wrong tagged (N_{wrong}) correctly tagged (N_{right}) and all (N_{all}) candidates and the effective tagging efficiency

$$\epsilon_{\text{eff}} = \epsilon_{\text{tag}} (1 - 2\omega)^2. \quad (2.4)$$

The effective tagging efficiency indicates the statistical degradation of the information and can thus be used to compare the performance of different tagging algorithms with each other and also in different decay modes.

3. Calibration of the Flavour Tagging

The mistag estimate η provided by the different tagging algorithms has to be corrected and transformed into the true mistag probability ω . This is done with a linear calibration function

$$\omega(\eta) = p_0 + p_1 (\eta - \langle \eta \rangle) \quad (3.1)$$

with the mean mistag estimate $\langle \eta \rangle$. The parameters p_0 and p_1 of this calibration function are extracted in two different ways. With charged decay modes as $B^+ \rightarrow J/\psi K^+$ and $B^+ \rightarrow D^0 \pi^+$ the true mistag ω can be extracted by comparing the tag with the charge of the kaon or pion in the final state. In neutral decay modes as $B^0 \rightarrow J/\psi K^{*0}$, $B^0 \rightarrow D^{*-} \mu^+ \nu_\mu$ or $B_s^0 \rightarrow D_s^- \pi^+$ a full time-dependent analysis is necessary to extract ω from the mixing asymmetry:

$$A_{\text{mix}}(t) \propto (1 - 2\omega) \cos(\Delta m_{d/s} t) \quad (3.2)$$

In both cases the calculation of ω is done in bins of the mistag estimate η and the linear function 3.1 is fitted to the (ω, η) pairs. Figure 2 shows for one bin the time dependent mixing asymmetry and the linear calibration function for the calibration mode $B^0 \rightarrow J/\psi K^{*0}$.




Figure 2: Mixing asymmetry (left) and calibration function (right) for the OS tagger combination in the calibration mode $B^0 \rightarrow J/\psi K^{*0}$.

4. Flavour Tagging in Run I

5. Developments

6. Conclusion

References

- [1] LHCb Collaboration, R.Aaij et. al., *Opposite-side flavour tagging of B mesons at the LHCb experiment*, *Eur.Phys.J. C* 72 (2012) 2022

- [2] LHCb Collaboration, R. Aaij et. al., *Optimization and calibration of the same-side kaon tagging algorithm using hadronic B_s^0 decays in 2011 data*, LHCb-CONF-2012-033
- [3] LHCb Collaboration, R. Aaij et. al., *Measurement of CP violation and the B_s^0 meson decay width difference with $B_s^0 \rightarrow J/\psi K^+ K^-$ and $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Rev. D87 (2013) 11, 112010
- [4] LHCb Collaboration, R. Aaij et. al., *Precision measurement of CP violation in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays*, Phys.Rev.Lett. 114 (2015) 4, 041801
- [5] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP-violating phase ϕ_s in $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Lett. B713 (2012) 378-386
- [6] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP-violating phase ϕ_s in $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays*, Phys.Lett. B736 (2014) 186-195
- [7] LHCb Collaboration, R. Aaij et. al., *Measurement of the CP-violating phase ϕ_s in $\bar{B}_s^0 \rightarrow D_s^+ D_s^-$ decays*, Phys.Rev.Lett. 113 (2014) 21, 211801
- [8] LHCb Collaboration, R. Aaij et. al., *Measurement of CP asymmetry in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays*, JHEP 1411 (2014) 060
- [9] LHCb Collaboration, R. Aaij et. al., *Measurement of the time-dependent CP asymmetry in $B^0 \rightarrow J/\psi K_s^0$ decays*, Phys.Lett. B721 (2013) 24-31
- [10] LHCb Collaboration, R. Aaij et. al., *Measurement of CP violation in $B^0 \rightarrow J/\psi K_s^0$ decays*, Phys.Rev.Lett. 115 (2015) 3, 031601
- [11] LHCb Collaboration, R. Aaij et. al., *Measurement of the time-dependent CP asymmetries in $B_s^0 \rightarrow J/\psi K_s^0$* , JHEP 1506 (2015) 131
- [12] LHCb Collaboration, R. Aaij et. al., *B flavor tagging using reconstructed charm decays at the LHCb experiment*, LHCb-PAPER-2015.027
- [13] G. A. Krockner, *Development and calibration of a same side kaon tagging algorithm and measurement of the $B_s^0 - \bar{B}_s^0$ oscillation frequency Δm_s at the LHCb experiment*, PhD thesis, Heidelberg U., Sep, 2013, CERN-THESIS-2013-213