

3. (25%)


(a) The recurrence relation for $T(n)$ of the modified algorithm is:

$$T(n) = \begin{cases} C_0 & \text{if } x = 1 \\ 5T(\frac{n}{5}) + cn & \text{otherwise} \end{cases}$$

This is the same as:

$$T(n) = c_0n + 5 \sum_{k=0}^{\log_5(\frac{n}{5})-1} cn$$

(b) Recurrence Tree:

(c) We wish to show that $T(n) = \Theta(n \log(n))$

$$\begin{aligned}
 T(n) &= c_0n + 5 \sum_{k=0}^{\log_5(\frac{n}{5})-1} cn \\
 &= n(c_0 + 5c \log_5(\frac{n}{5}) - c) \\
 &= n(c_0 + 5c \log_5(n) - \log_5(5) - c) \\
 &= n(c_0 + 5c \log_5(n) - 1 - c) \\
 &= n(c_1 + 5c \log_5(n)) \\
 &= n(c_1 + 5c \left(\frac{\log_2(n)}{\log_2(5)} \right)) \\
 &= n(c_1 + 5c_2 \log_2(n)) \\
 &= nc_1 + c_3n \log_2(n) \\
 &= \Theta(n \log(n))
 \end{aligned}$$

(d) By dividing the problem into m subproblems, we divide n into m pieces at each level, and repeat each subproblem m times. This results in a different logarithmic base and leading coefficient. Because of the change of base, this changes the constant coefficient multiplier, which doesn't alter the asymptotic result.