

# Eigenvalues of Schrödinger operators perturbed by dissipative barriers

Alexei Stepanenko

PhD supervisors: Jonathan Ben-Artzi and Marco Marletta

Cardiff University

February 2, 2021

## Introduction

Consider the Schrödinger operator

$$H_R := -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \quad \text{on} \quad L^2(\mathbb{R}_+) \quad (R > 0),$$

endowed with a Dirichlet boundary condition at 0, where:

1.  $q \in L^1(\mathbb{R}_+)$  (background potential)
  2.  $\gamma > 0$ .
- Multiplication operator  $i\gamma\chi_{[0,R]}$  is referred to as a *dissipative barrier*.

## Introduction

Consider the Schrödinger operator

$$H_R := -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \quad \text{on} \quad L^2(\mathbb{R}_+) \quad (R > 0),$$

endowed with a Dirichlet boundary condition at 0, where:

1.  $q \in L^1(\mathbb{R}_+)$  (background potential)
  2.  $\gamma > 0$ .
- ▶ Multiplication operator  $i\gamma\chi_{[0,R]}$  is referred to as a *dissipative barrier*.
  - ▶ Dissipative barriers are useful in numerical analysis and arise in many settings.

## Introduction

Consider the Schrödinger operator

$$H_R := -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \quad \text{on} \quad L^2(\mathbb{R}_+) \quad (R > 0),$$

endowed with a Dirichlet boundary condition at 0, where:

1.  $q \in L^1(\mathbb{R}_+)$  (background potential)
  2.  $\gamma > 0$ .
- ▶ Multiplication operator  $i\gamma\chi_{[0,R]}$  is referred to as a *dissipative barrier*.
  - ▶ Dissipative barriers are useful in numerical analysis and arise in many settings.

**Question:** How do the eigenvalues of  $H_R$  behave for large  $R$ ?

[1] Spectral inclusion and pollution for a class of dissipative perturbations, S., Journal of Mathematical Physics **62**, 013501 (2021) [arXiv:2006.10097](#)

## Introduction

Consider the Schrödinger operator

$$H_R := -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \quad \text{on} \quad L^2(\mathbb{R}_+) \quad (R > 0),$$

endowed with a Dirichlet boundary condition at 0, where:

1.  $q \in L^1(\mathbb{R}_+)$  (background potential)
  2.  $\gamma > 0$ .
- ▶ Multiplication operator  $i\gamma\chi_{[0,R]}$  is referred to as a *dissipative barrier*.
  - ▶ Dissipative barriers are useful in numerical analysis and arise in many settings.

**Question:** How do the eigenvalues of  $H_R$  behave for large  $R$ ?

[1] Spectral inclusion and pollution for a class of dissipative perturbations, S., Journal of Mathematical Physics **62**, 013501 (2021) [arXiv:2006.10097](#)

[2] Bounds for Schrödinger operators on the half-line perturbed by dissipative barriers, S., Submitted. (2020) [arXiv:2010.05663](#)

## Spectral inclusion

We have (as  $R \rightarrow \infty$ )

$$H_R = -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \xrightarrow{s} -\frac{d^2}{dx^2} + q + i\gamma =: H_\infty.$$

## Spectral inclusion

We have (as  $R \rightarrow \infty$ )

$$H_R = -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \xrightarrow{s} -\frac{d^2}{dx^2} + q + i\gamma =: H_\infty.$$

**Question:** Is  $\sigma(H_\infty)$  approximated by  $\sigma(H_R)$ ?

## Spectral inclusion

We have (as  $R \rightarrow \infty$ )

$$H_R = -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \xrightarrow{s} -\frac{d^2}{dx^2} + q + i\gamma =: H_\infty.$$

**Question:** Is  $\sigma(H_\infty)$  approximated by  $\sigma(H_R)$ ?

**Theorem 1** ([1])

- (a) *For any eigenvalue  $\lambda$  of  $H_\infty$ , there exists eigenvalues  $\lambda_R$  of  $H_R$  and constants  $C_0, \beta, R_0 > 0$  such that*

$$|\lambda_R - \lambda| \leq C_0 e^{-\beta R} \quad (R \geq R_0).$$



## Spectral inclusion

We have (as  $R \rightarrow \infty$ )

$$H_R = -\frac{d^2}{dx^2} + q + i\gamma\chi_{[0,R]} \xrightarrow{s} -\frac{d^2}{dx^2} + q + i\gamma =: H_\infty.$$

**Question:** Is  $\sigma(H_\infty)$  approximated by  $\sigma(H_R)$ ?

### Theorem 1 ([1])

- (a) *For any eigenvalue  $\lambda$  of  $H_\infty$ , there exists eigenvalues  $\lambda_R$  of  $H_R$  and constants  $C_0, \beta, R_0 > 0$  such that*

$$|\lambda_R - \lambda| \leq C_0 e^{-\beta R} \quad (R \geq R_0).$$

- (b) *If  $\exists \varepsilon > 0 : \int e^{\varepsilon t} |q(t)| dt < \infty$ , then for (almost<sup>1</sup>) any  $\mu \in \sigma_{\text{ess}}(H_\infty)$  there exists eigenvalues  $\lambda_R$  of  $H_R$  and constants  $C_0, R_0 > 0$  such that*

$$|\lambda_R - \mu| \leq \frac{C_0}{R} \quad (R \geq R_0).$$

<sup>1</sup>Any  $\mu \in \sigma_{\text{ess}}(H_\infty)$  which is not an embedded resonance or the band-end.

## Theorem 1 ([1])


- (a) For any eigenvalue  $\lambda$  of  $H_\infty$ , there exists eigenvalues  $\lambda_R$  of  $H_R$  and constants  $C_0, \beta, R_0 > 0$  such that

$$|\lambda_R - \lambda| \leq C_0 e^{-\beta R} \quad (R \geq R_0).$$


- (b) If  $\exists \varepsilon > 0 : \int e^{\varepsilon t} |q(t)| dt < \infty$ , then for (almost) any  $\mu \in \sigma_{\text{ess}}(H_\infty)$  there exists eigenvalues  $\lambda_R$  of  $H_R$  and constants  $C_0, R_0 > 0$  such that

$$|\lambda_R - \mu| \leq \frac{C_0}{R} \quad (R \geq R_0).$$

## Example




## Example



Next steps:

- Enclosures for the eigenvalues of  $H_R$

## Example



### Next steps:


- ▶ Enclosures for the eigenvalues of  $H_R$ .
- ▶ Estimates for the number of eigenvalues  $N(H_R)$ .

## Enclosures

### Theorem 2 ([2])

- (a)  $\exists$  constant  $X = X(q, \gamma) > 0$  such that  $\forall R > 0$  the eigenvalues of  $H_R$  lie in

$$\Gamma := B_X(0) \cup ([0, \infty) + i[0, \gamma]).$$



## Enclosures


### Theorem 2 ([2])

- (a)  $\exists$  constant  $X = X(q, \gamma) > 0$  such that  $\forall R > 0$  the eigenvalues of  $H_R$  lie in

$$\Gamma := B_X(0) \cup ([0, \infty) + i[0, \gamma]).$$

- (b)  $\exists$  constant  $R_0 = R_0(q, \gamma) > 0$  such that any eigenvalue  $\lambda_R$  of  $H_R$  satisfies

$$\sqrt{|\lambda_R - i\gamma|} \leq \frac{5\gamma R}{\log R} \quad (R \geq R_0).$$



## Enclosures

### Theorem 2 ([2])


- (a)  $\exists$  constant  $X = X(q, \gamma) > 0$  such that  $\forall R > 0$  the eigenvalues of  $H_R$  lie in


$$\Gamma := B_X(0) \cup ([0, \infty) + i[0, \gamma]).$$

- (b)  $\exists$  constant  $R_0 = R_0(q, \gamma) > 0$  such that any eigenvalue  $\lambda_R$  of  $H_R$  satisfies

$$\sqrt{|\lambda_R - i\gamma|} \leq \frac{5\gamma R}{\log R} \quad (R \geq R_0).$$

- To compare to (b), application of a *sharp* enclosure of Frank, Laptev and Seiringer (2011) gives  $\sqrt{|\lambda_R|} = O(R)$ .





*Ideas in proof of Theorem 2.*

1. **Use large- $|\lambda|$  Levinson asymptotics:** Solutions  $\psi_{\pm}(\cdot, \lambda)$  to  $-\psi'' + q\psi = \lambda\psi$  such that

$$\psi_{\pm}(x, \lambda) = e^{\pm i\sqrt{\lambda}x}(1 + E_{\pm}(x, \lambda))$$

where

$$|E_{\pm}(x, \lambda)| = O\left(\frac{1}{\sqrt{|\lambda|}}\right) \quad \text{as } |\lambda| \rightarrow \infty.$$



*Ideas in proof of Theorem 2.*

1. **Use large- $|\lambda|$  Levinson asymptotics:** Solutions  $\psi_{\pm}(\cdot, \lambda)$  to  $-\psi'' + q\psi = \lambda\psi$  such that

$$\psi_{\pm}(x, \lambda) = e^{\pm i\sqrt{\lambda}x}(1 + E_{\pm}(x, \lambda))$$

where

$$|E_{\pm}(x, \lambda)| = O\left(\frac{1}{\sqrt{|\lambda|}}\right) \quad \text{as } |\lambda| \rightarrow \infty.$$

2. Construct analytic function  $f_R$  such that

$$\lambda \text{ eigenvalue of } H_R \iff f_R(\lambda) = 0.$$

$f_R$  has form

$$\begin{aligned} f_R(\lambda) = & \psi_{-}(0, \lambda - i\gamma) \left( \sqrt{\lambda} - \sqrt{\lambda - i\gamma} + \mathcal{E}_1(R, \lambda) \right) e^{i\sqrt{\lambda - i\gamma}R} \\ & - \psi_{+}(0, \lambda - i\gamma) \left( \sqrt{\lambda} + \sqrt{\lambda - i\gamma} + \mathcal{E}_2(R, \lambda) \right) e^{-i\sqrt{\lambda - i\gamma}R} \end{aligned}$$

where  $|\mathcal{E}_1(R, \lambda)| + |\mathcal{E}_2(R, \lambda)| \leq C(q, \gamma)$ .

## Counting eigenvalues

*Notation.*  $N(H_R)$  = Number of eigenvalues of  $H_R$ .

# Counting eigenvalues

*Notation.*  $N(H_R)$  = Number of eigenvalues of  $H_R$ .

## Theorem 3 ([2])

- (a) *If  $q$  is compactly supported, then there exists constant  $R_0 = R_0(q, \gamma) > 0$  such that*

$$N(H_R) \leq \frac{11}{\log 2} \frac{\gamma R^2}{\log R} \quad (R \geq R_0).$$

# Counting eigenvalues

*Notation.*  $N(H_R)$  = Number of eigenvalues of  $H_R$ .

## Theorem 3 ([2])

- (a) *If  $q$  is compactly supported, then there exists constant  $R_0 = R_0(q, \gamma) > 0$  such that*

$$N(H_R) \leq \frac{11}{\log 2} \frac{\gamma R^2}{\log R} \quad (R \geq R_0).$$

- (b) *If  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$ , then*

$$N(H_R) \leq C \frac{\sqrt{X} + a}{a^2} \frac{\gamma R^3}{(\log R)^2} \quad (R \geq R_0)$$

*where  $X = X(q, \gamma) > 0$  appeared in Th. 2 (a) and  $C = 88788$ .*

# Counting eigenvalues

*Notation.*  $N(H_R)$  = Number of eigenvalues of  $H_R$ .

## Theorem 3 ([2])

(a) If  $q$  is compactly supported, then there exists constant  $R_0 = R_0(q, \gamma) > 0$  such that

$$N(H_R) \leq \frac{11}{\log 2} \frac{\gamma R^2}{\log R} \quad (R \geq R_0).$$

(b) If  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$ , then

$$N(H_R) \leq C \frac{\sqrt{X} + a}{a^2} \frac{\gamma R^3}{(\log R)^2} \quad (R \geq R_0)$$

where  $X = X(q, \gamma) > 0$  appeared in Th. 2 (a) and  $C = 88788$ .

► Comparing to application of results in the literature:

	Th. 3	Application of literature
Compactly supported	$N(H_R) = O\left(\frac{R^2}{\log R}\right)$	$N(H_R) = O(R^2)$ Korotyaev (2020)

# Counting eigenvalues

*Notation.*  $N(H_R)$  = Number of eigenvalues of  $H_R$ .

## Theorem 3 ([2])

- (a) If  $q$  is compactly supported, then there exists constant  $R_0 = R_0(q, \gamma) > 0$  such that

$$N(H_R) \leq \frac{11}{\log 2} \frac{\gamma R^2}{\log R} \quad (R \geq R_0).$$

- (b) If  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$ , then

$$N(H_R) \leq C \frac{\sqrt{X} + a}{a^2} \frac{\gamma R^3}{(\log R)^2} \quad (R \geq R_0)$$

where  $X = X(q, \gamma) > 0$  appeared in Th. 2 (a) and  $C = 88788$ .

- Comparing to application of results in the literature:


	Th. 3	Application of literature
Compactly supported	$N(H_R) = O\left(\frac{R^2}{\log R}\right)$	$N(H_R) = O(R^2)$ Korotyaev (2020)
Naimark	$N(H_R) = O\left(\frac{R^3}{(\log R)^2}\right)$	$N(H_R) = O(R^4)$ Frank, Laptev, Safronov (2016)

	Th. 3	Application of literature
Compactly supported	$N(H_R) = O\left(\frac{R^2}{\log R}\right)$	$N(H_R) = O(R^2)$ Korotyaev (2020)
Naimark	$N(H_R) = O\left(\frac{R^3}{(\log R)^2}\right)$	$N(H_R) = O(R^4)$ Frank, Laptev, Safronov (2016)

*Ideas in proof of Theorem 3.*

1. Construct analytic function  $g_R$  on  $\mathbb{C}_+$  such that

$$g_R(z) = 0 \iff z^2 \text{ eigval of } H_R.$$




	Th. 3	Application of literature
Compactly supported	$N(H_R) = O\left(\frac{R^2}{\log R}\right)$	$N(H_R) = O(R^2)$ Korotyaev (2020)
Naimark	$N(H_R) = O\left(\frac{R^3}{(\log R)^2}\right)$	$N(H_R) = O(R^4)$ Frank, Laptev, Safronov (2016)


*Ideas in proof of Theorem 3.*

1. Construct analytic function  $g_R$  on  $\mathbb{C}_+$  such that

$$g_R(z) = 0 \iff z^2 \text{ eigval of } H_R.$$



2. If  $q$  compactly supported,  $g_R$  admits analytic continuation to  $\mathbb{C}$ .






	Th. 3	Application of literature
Compactly supported	$N(H_R) = O\left(\frac{R^2}{\log R}\right)$	$N(H_R) = O(R^2)$ Korotyaev (2020)
Naimark	$N(H_R) = O\left(\frac{R^3}{(\log R)^2}\right)$	$N(H_R) = O(R^4)$ Frank, Laptev, Safronov (2016)

*Ideas in proof of Theorem 3.*


1. Construct analytic function  $g_R$  on  $\mathbb{C}_+$  such that

$$g_R(z) = 0 \iff z^2 \text{ eigval of } H_R.$$



2. If  $q$  compactly supported,  $g_R$  admits analytic continuation to  $\mathbb{C}$ .


- Apply Jensen's formula to bound  $N(H_R)$  and prove Th. 3 (a).



*Ideas in proof of Theorem 3. (cont.)*

3.  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$

$\Rightarrow g_R$  admits analytic continuation to  $\{\Im z > -2a\}$ .




*Ideas in proof of Theorem 3. (cont.)*

3.  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$

$\Rightarrow g_R$  admits analytic continuation to  $\{\Im z > -2a\}$ .

► Enclosure Th. 2  $\Rightarrow \Im \sqrt{\lambda_R} \leq \sqrt{X}$  for any eigval  $\lambda_R$  of  $H_R$   
(recall that  $X = X(q, \gamma) > 0$ ).




*Ideas in proof of Theorem 3. (cont.)*

3.  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$

$\Rightarrow g_R$  admits analytic continuation to  $\{\Im z > -2a\}$ .

- ▶ Enclosure Th. 2  $\Rightarrow \Im \sqrt{\lambda_R} \leq \sqrt{X}$  for any eigval  $\lambda_R$  of  $H_R$  (recall that  $X = X(q, \gamma) > 0$ ).
- ▶ Prove bound for number of zeros of an arbitrary analytic function on  $\{\Im z > -2a\}$  in the region

$$D_{r,X} := \left\{ z \in \mathbb{C}_+ : \Im z \leq \sqrt{X}, |z| \leq r \right\}.$$



*Ideas in proof of Theorem 3. (cont.)*


3.  $\exists a > 0 : \int e^{4at} |q(t)| dt < \infty$

$\Rightarrow g_R$  admits analytic continuation to  $\{\Im z > -2a\}$ .

- Enclosure Th. 2  $\Rightarrow \Im \sqrt{\lambda_R} \leq \sqrt{X}$  for any eigval  $\lambda_R$  of  $H_R$  (recall that  $X = X(q, \gamma) > 0$ ).
- Prove bound for number of zeros of an arbitrary analytic function on  $\{\Im z > -2a\}$  in the region

$$D_{r,X} := \left\{ z \in \mathbb{C}_+ : \Im z \leq \sqrt{X}, |z| \leq r \right\}.$$

- Apply this bound to  $g_R$  with  $r = O(R/\log R)$ .



# Thanks for listening!

## References

- [1] *Spectral inclusion and pollution for a class of dissipative perturbations*, S., Journal of Mathematical Physics **62**, 013501 (2021) [arXiv:2006.10097](#)
- [2] *Bounds for Schrödinger operators on the half-line perturbed by dissipative barriers*, S., Submitted. (2020) [arXiv:2010.05663](#)
- [3] *A Sharp Bound on Eigenvalues of Schrödinger Operators on the Half-line with Complex-valued Potentials*, R Frank, A Laptev, R Seiringer, in Spectral Theory and Analysis, Operator Theory: Advances and Applications. Springer, Basel, pp. 39–44 (2011)
- [4] *Trace formulas for Schrödinger operators with complex potentials on a half line*, E Korotyaev, Lett Math Phys **110**, 1–20 (2020)
- [5] *On the number of eigenvalues of Schrödinger operators with complex potentials*, R Frank, A Laptev, O Safronov, Journal of the London Mathematical Society **94**, 377–390 (2016)

# **Appendix**

## Proposition

Suppose that  $f$  is an analytic function defined on an open neighbourhood of the closed semi-disc  $D_r := \overline{B}_r(0) \cap \overline{\mathbb{C}}_+$  for some  $r > 0$ . Let  $\alpha$  and  $\beta$  be any numbers in the interval  $(0, 1)$  satisfying

$$\beta \left( \frac{1 - \alpha}{\alpha + \beta} \right)^2 > \frac{Y}{\eta} \quad (1)$$

and let  $N(\alpha r)$  denote the number of zeros in the region

$$D_{\alpha r, \eta, Y} := \{z \in \mathbb{C} : \eta \leq \Im z \leq Y, |z| \leq \alpha r\} \quad (2)$$

where  $Y, \eta > 0$  are given parameters satisfying  $\eta < Y < r$ . Then,

$$N(\alpha r) \leq \frac{2}{\log \Lambda(r)} \log \left( \frac{1}{\min\{\beta, 1 - \beta\}} \frac{\sup_{z \in \partial D_r} |f(z)|}{|f(i\beta r)|} \right) \quad (3)$$

where

$$\Lambda(r) := \frac{1 + \frac{4\beta\eta}{(\alpha+\beta)^2} \frac{1}{r}}{1 + \frac{4Y}{(1-\alpha)^2} \frac{1}{r}}. \quad (4)$$

## Remark

One can always guarantee that condition (1) for  $\alpha$  and  $\beta$  is satisfied by choosing, for instance,

$$\alpha = \beta = \frac{1}{4} \frac{\eta}{2Y + \eta}. \quad (5)$$