Commutators of Riesz potential in the vanishing generalized weighted Morrey spaces with variable exponent

Vagif S. Guliyev a,b,c,*), Javanshir J. Hasanov d, Xayyam A. Badalov c

^a Dumlupinar University, Department of Mathematics, 43020 Kutahya, Turkey
 ^b S.M. Nikolskii Institute of Mathematics at RUDN University, 6 Miklukho-Maklay St, Moscow, 117198
 ^c Institute of Mathematics and Mechanics of NAS of Azerbaijan, AZ1141 Baku, Azerbaijan
 ^d Azerbaijan State Oil and Industry University, Azadlig av.20, AZ 1601, Baku, Azerbaijan

Abstract

Let $\Omega\subset\mathbb{R}^n$ be an unbounded open set. We consider the generalized weighted Morrey spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ and the vanishing generalized weighted Morrey spaces $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ with variable exponent p(x) and a general function $\varphi(x,r)$ defining the Morrey-type norm. The main result of this paper are the boundedness of Riesz potential and its commutators on the spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ and $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$. This result generalizes several existing results for Riesz potential and its commutators on Morrey type spaces. Especially, it gives a unified result for generalized Morrey spaces and variable Morrey spaces which currently gained a lot of attentions from researchers in theory of function spaces.

AMS Mathematics Subject Classification: 42B20, 42B25, 42B35Key words: Riesz potential, commutator, vanishing generalized weighted Morrey space with variable exponent, BMO space

1 Introduction

The variable exponent generalized weighted Morrey spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ over an open set $\Omega \subset \mathbb{R}^n$ was introduced and the boundedness of the Hardy-Littlewood maximal operator, the singular integral operators and their commutators on these spaces was proven in [28]. The main focus of this article is to prove that the Riesz potential and its commutators are bounded on generalized weighted Morrey spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$

^{*)} Corresponding author.

and vanishing generalized weighted Morrey spaces $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ with variable exponents. Also some Sobolev-type inequalities for Riesz potentials on spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ and $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ are proved.

The classical Morrey spaces were introduced by Morrey [38] to study the local behavior of solutions to second-order elliptic partial differential equations. Moreover, various Morrey-type spaces are defined in the process of study. Mizuhara [39] and Nakai [42] introduced generalized Morrey spaces $M^{p,\varphi}(\mathbb{R}^n)$ (see, also [21]); Komori and Shirai [33] defined weighted Morrey spaces $L^{p,\kappa}(w)$; Guliyev [22] gave a concept of the generalized weighted Morrey spaces $M^{p,\varphi}_w(\mathbb{R}^n)$ which could be viewed as extension of both $M^{p,\varphi}(\mathbb{R}^n)$ and $L^{p,\kappa}(w)$.

Vanishing Morrey spaces $VM^{p,\varphi}(\mathbb{R}^n)$ are subspaces of functions in Morrey spaces which were introduced by Vitanza [52] satisfying the condition

$$\lim_{\substack{r \to 0 \\ 0 < t < r}} \sup_{x \in \mathbb{R}^n} r^{-\frac{\lambda}{p}} ||f\chi_{B(x,t)}||_{L^{p(\cdot)}(B(x,t)} = 0$$

and applied there to obtain a regularity result for elliptic partial differential equations. Also Ragusa [44] proved a sufficient condition for commutators of fractional integral operators to belong to vanishing Morrey spaces $VM^{p,\lambda}(\mathbb{R}^n)$. The vanishing generalized Morrey spaces $VM^{p,\varphi}(\mathbb{R}^n)$ were introduced and studied by Samko in [46], see also [3, 17, 37].

As it is known, last two decades there is an increasing interest to the study of variable exponent spaces and operators with variable parameters in such spaces, we refer for instance to the surveying papers [16, 32, 47], on the progress in this field, including topics of Harmonic Analysis and Operator Theory, see also references therein. For mapping properties of maximal functions and Riesz potential on Lebesgue spaces with variable exponent we refer to [9, 10, 13, 14, 15, 31, 35, 45].

Variable exponent Morrey spaces $\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega)$, were introduced and studied in [4] and [40] in the Euclidean setting. The boundedness of Riesz potential in variable exponent Morrey spaces $\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega)$ under the log-condition on $p(\cdot),\lambda(\cdot)$ and a Sobolev type $\mathcal{L}^{p(\cdot),\lambda(\cdot)}\to\mathcal{L}^{q(\cdot),\lambda(\cdot)}$ —theorem for potential operators of variable order $\alpha(x)$ was proved in [31]. In the case of constant α , there was also proved a boundedness theorem in the limiting case $p(x)=\frac{n-\lambda(x)}{\alpha}$, when the potential operator I^{α} acts from $\mathcal{L}^{p(\cdot),\lambda(\cdot)}$ into BMO was proved in [4]. In [40] the maximal operator and potential operators were considered in a somewhat more general space, but under more restrictive conditions on p(x).

Generalized Morrey spaces of such a kind in the case of constant p were studied in [18, 21, 39, 42, 50, 51]. In the case of bounded sets the boundedness of the maximal operator, singular integral operator and the potential operators in generalized variable exponent Morrey type spaces was proved in [24, 25, 26] and in the case of unbounded sets in [27]. Also, in the case of bounded sets the boundedness of these operators in generalized variable exponent weighted Morrey spaces for the power weights was proved in [30].

In the case of constant p and λ , the results on the boundedness of potential operators go back to [1] and [43], respectively, while the boundedness of the maximal operator in the Euclidean setting was proved in [12]; for further results in the case of constant p and λ (see, for instance, [5, 19]).

In the spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ over open sets $\Omega \subset \mathbb{R}^n$ we consider the following operators:

1) the Hardy-Littlewood maximal operator

$$Mf(x) = \sup_{r>0} |B(x,r)|^{-1} \int_{\widetilde{B}(x,r)} |f(y)| dy,$$

2) Riesz potential operator

$$I^{\alpha}f(x) = \int_{\Omega} |x - y|^{\alpha - n} f(y) dy, \quad 0 < \alpha < n,$$

3) the fractional maximal operator

$$M^{\alpha} f(x) = \sup_{r>0} |B(x,r)|^{\frac{\alpha}{n}-1} \int_{\widetilde{B}(x,r)} |f(y)| dy, \ 0 \le \alpha < n.$$

We find the condition on the function $\,\varphi(x,r)\,$ for the boundedness of the Riesz potential $\,I^{\alpha}\,$ and its commutators in the generalized weighted Morrey space $\,\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)\,$ and the vanishing generalized weighted Morrey spaces $\,V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)\,$ with variable $\,p(x)\,$ under the log-condition on $\,p(\cdot)\,$.

The paper is organized as follows. In Section 2 we provide necessary preliminaries on variable exponent weighted Lebesgue, generalized weighted Morrey spaces and vanishing generalized weighted Morrey spaces. In Section 3 we prove the boundedness of Riesz potential and its commutators on the variable exponent generalized weighted Morrey spaces. In Section 4 we prove the boundedness of Riesz potential and its commutators on the variable exponent vanishing generalized weighted Morrey spaces.

The main results are given in Theorems 3.4, 3.5, 3.8, 3.9, 3.10, 4.1 and 4.2. We emphasize that the results we obtain for generalized Morrey spaces are new even in the case when p(x) is constant, because we do not impose any monotonicity type condition on $\varphi(r)$.

We use the following notation: \mathbb{R}^n is the n-dimensional Euclidean space, $\Omega \subset \mathbb{R}^n$ is an open set, $\chi_E(x)$ is the characteristic function of a set $E \subseteq \mathbb{R}^n$, $B(x,r) = \{y \in \mathbb{R}^n : |x-y| < r\}\}$, $\widetilde{B}(x,r) = B(x,r) \cap \Omega$, by c, C, c_1 , c_2 etc, we denote various absolute positive constants, which may have different values even in the same line.

2 Preliminaries on variable exponent weighted Lebesgue, generalized weighted Morrey spaces and vanishing generalized weighted Morrey spaces

We refer to the book [14] for variable exponent Lebesgue spaces but give some basic definitions and facts. Let $p(\cdot)$ be a measurable function on Ω with values in $(1,\infty)$. An open set Ω which may be unbounded throughout the whole paper. We mainly suppose that

$$1 < p_{-} \le p(x) \le p_{+} < \infty, \tag{2.1}$$

where $p_-:=\operatorname*{ess\ inf}_{x\in\Omega}p(x)$, $p_+:=\operatorname*{ess\ sup}_{x\in\Omega}p(x)$. By $L^{p(\cdot)}(\Omega)$ we denote the space of all measurable functions f(x) on Ω such that

$$I_{p(\cdot)}(f) = \int_{\Omega} |f(x)|^{p(x)} dx < \infty.$$

Equipped with the norm

$$||f||_{p(\cdot)} = \inf \left\{ \eta > 0 : I_{p(\cdot)} \left(\frac{f}{\eta} \right) \le 1 \right\},$$

this is a Banach function space. By $p'(\cdot) = \frac{p(x)}{p(x)-1}$, $x \in \Omega$, we denote the conjugate exponent.

The space $\,L^{p(\cdot)}(\Omega)\,$ coincides with the space

$$\left\{ f(x) : \left| \int_{\Omega} f(y)g(y)dy \right| < \infty \text{ for all } g \in L^{p'(\cdot)}(\Omega) \right\}$$
 (2.2)

up to the equivalence of the norms

$$||f||_{L^{p(\cdot)}(\Omega)} \approx \sup_{||g||_{L^{p'(\cdot)}} \le 1} \left| \int_{\Omega} f(y)g(y)dy \right|, \tag{2.3}$$

see [36, Proposition 2.2], see also [34, Theorem 2.3], or [48, Theorem 3.5].

For the basics on variable exponent Lebesgue spaces we refer to [49], [34]. $\mathcal{P}(\Omega)$ is the set of bounded measurable functions $p:\Omega\to[1,\infty)$; $\mathcal{P}^{log}(\Omega)$ is the set of exponents $p\in\mathcal{P}(\Omega)$ satisfying the local log-condition

$$|p(x) - p(y)| \le \frac{A}{-\ln|x - y|}, |x - y| \le \frac{1}{2} x, y \in \Omega,$$
 (2.4)

where A = A(p) > 0 does not depend on x, y;

 $\mathcal{A}^{log}(\Omega)$ is the set of bounded exponents $p:\Omega\to\mathbb{R}^n$ satisfying the condition (2.4);

 $\mathbb{P}^{log}(\Omega)$ is the set of exponents $p \in \mathcal{P}^{log}(\Omega)$ with $1 < p_- \le p_+ < \infty$; for Ω which may be unbounded, by $\mathcal{P}_{\infty}(\Omega)$, $\mathcal{P}^{log}_{\infty}(\Omega)$, $\mathbb{P}^{log}_{\infty}(\Omega)$, $\mathcal{A}^{log}_{\infty}(\Omega)$ we denote the subsets of the above sets of exponents satisfying the decay condition (when Ω is unbounded)

$$|p(x) - p(\infty)| \le \frac{A_{\infty}}{\ln(2+|x|)}, \quad x \in \mathbb{R}^n, \tag{2.5}$$

where $p_{\infty} = \lim_{x \to \infty} p(x) > 1$.

We will also make use of the estimate provided by the following lemma (see [14], Corollary 4.5.9).

$$\|\chi_{\widetilde{B}(x,r)}(\cdot)\|_{p(\cdot)} \le Cr^{\theta_p(x,r)}, \quad x \in \Omega, \ p \in \mathbb{P}_{\infty}^{log}(\Omega),$$
 (2.6)

where
$$\theta_p(x,r) = \begin{cases} \frac{n}{p(x)}, & r \leq 1, \\ \frac{n}{p(\infty)}, & r > 1 \end{cases}$$

By ω we always denote a weight, i.e. a positive, locally integrable function with domain Ω . The weighted Lebesgue space $L^{p(\cdot)}_{\omega}(\Omega)$ is defined as the set of all measurable functions for which

$$||f||_{L^{p(\cdot)}(\Omega)} = ||f\omega||_{L^{p(\cdot)}(\Omega)}.$$

Let us define the class $A_{p(\cdot)}(\Omega)$ (see [16], [35]) to consist of those weights ω for which

$$[\omega]_{A_{p(\cdot)}} \equiv \sup_{B} |B|^{-1} \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,r))} \|\omega^{-1}\|_{L^{p'(\cdot)}(\widetilde{B}(x,r))} < \infty.$$

A weight function ω belongs to the class $A_{p(\cdot),q(\cdot)}(\Omega)$ if

$$[\omega]_{A_{p(\cdot),q(\cdot)}} \equiv \sup_{B} |B|^{\frac{1}{p(x)} - \frac{1}{q(x)} - 1} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,r))} \|\omega^{-1}\|_{L^{p'(\cdot)}(\widetilde{B}(x,r))} < \infty.$$

Lemma 2.1. Let p,q satisfy condition (2.1) and $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$, then $\omega^{-1} \in A_{q'(\cdot),p'(\cdot)}(\Omega)$, with $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$.

Proof. Let p,q satisfy condition (2.1) and $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$. Then $\varphi = \omega^{-1} \in A_{q'(\cdot),p'(\cdot)}(\Omega)$. Indeed,

$$\begin{split} &|B|^{\frac{1}{q'(x)} - \frac{1}{p'(x)} - 1} \|\varphi\|_{L^{p'(\cdot)}(\widetilde{B}(x,r))} \|\varphi^{-1}\|_{L^{q(\cdot)}(\widetilde{B}(x,r))} \\ = &|B|^{\frac{1}{p(x)} - \frac{1}{q(x)} - 1} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,r))} \|\omega^{-1}\|_{L^{p'(\cdot)}(\widetilde{B}(x,r))}. \end{split}$$

Theorem 2.1. [29, Therem 1.1] Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set and $p \in \mathbb{P}^{log}_{\infty}(\Omega)$. Then $M: L^{p(\cdot)}_{\omega}(\Omega) \to L^{p(\cdot)}_{\omega}(\Omega)$ if and only if $\omega \in A_{p(\cdot)}(\Omega)$.

For unbounded sets, say $\Omega = \mathbb{R}^n$, and constant orders a the corresponding Sobolev theorem proved in [8, 9] runs as follows.

Theorem 2.2. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$ and $p \in \mathbb{P}^{log}_{\infty}(\Omega)$. Let also $p_+ < \frac{n}{\alpha}$. Then the operators M^{α} and I^{α} are bounded from $L^{p(\cdot)}(\Omega)$ to $L^{q(\cdot)}(\Omega)$ with $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$.

Let $\lambda(x)$ be a measurable function on Ω with values in [0,n]. The variable Morrey space $\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega)$ and variable weighted Morrey space $\mathcal{L}^{p(\cdot),\lambda(\cdot)}_{\omega}(\Omega)$ is defined as the set of integrable functions f on Ω with the finite norms

$$||f||_{\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega)} = \sup_{x \in \Omega, \ t > 0} t^{-\frac{\lambda(x)}{p(x)}} ||f\chi_{\widetilde{B}(x,t)}||_{L^{p(\cdot)}(\Omega)},$$

$$||f||_{\mathcal{L}^{p(\cdot),\lambda(\cdot)}_{\omega}(\Omega)} = \sup_{x \in \Omega, \ t > 0} t^{-\frac{\lambda(x)}{p(x)}} ||f\chi_{\widetilde{B}(x,t)}||_{L^{p(\cdot)}_{\omega}(\Omega)}.$$

Let ω be a nonnegative measurable function on \mathbb{R}^n such that ω^p is locally integrable on \mathbb{R}^n . Then a Radon measure μ is canonically associated with the weight $\omega(\cdot)^{p(\cdot)}$, that is,

$$\mu(E) = \int_{E} \omega(y)^{p(y)} dy.$$

We denote by $\mathcal{L}^{p(\cdot),\lambda}(\mathbb{R}^n,d\mu)$ the set of all measurable functions f with finite norm

$$||f||_{\mathcal{L}^{p(\cdot),\lambda}(\mathbb{R}^n,d\mu)} = \inf \left\{ \eta > 0: \sup_{x \in \Omega, \ t > 0} \frac{t^{\lambda}}{\mu(B(x,t))} \int_{B(x,t)} \left(\frac{|f(y)|}{\eta} \right)^{p(y)} d\mu(y) \le 1 \right\}.$$

Theorem 2.3. [41] Let $0 < \alpha < n$, $0 \le \lambda < n$, $p \in \mathbb{P}^{log}_{\infty}(\mathbb{R}^n)$, $p_+ < \frac{\lambda}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{\lambda}$, $\omega \in A_{p(\cdot)}(\mathbb{R}^n)$. Then the operator I^{α} is bounded from $\mathcal{L}^{p(\cdot),\lambda}(\mathbb{R}^n,d\mu)$ to $\mathcal{L}^{q(\cdot),\lambda}(\mathbb{R}^n,d\mu)$.

In view of the well known pointwise estimate $\,M^{\alpha}f(x) \leq C(I^{\alpha}|f|)(x)$, it suffices to treat only the case of the operator $\,I^{\alpha}$.

Corollary 2.1. ([11], [41]) Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}_{\infty}^{log}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$. Then the operators M^{α} and I^{α} are bounded from $L_{\omega}^{p(\cdot)}(\Omega)$ to $L_{\omega}^{q(\cdot)}(\Omega)$.

Let M^{\sharp} be the sharp maximal function defined by

$$M^{\sharp}f(x) = \sup_{r>0} |B(x,r)|^{-1} \int_{\widetilde{B}(x,r)} |f(y) - f_{\widetilde{B}(x,r)}| dy,$$

where $f_{\widetilde{B}(x,t)}(x) = |\widetilde{B}(x,t)|^{-1} \int_{\widetilde{B}(x,t)} f(y) dy$.

Definition 2.1. We define the $BMO(\Omega)$ space as the set of all locally integrable functions f with finite norm

$$||f||_{BMO} = \sup_{x \in \Omega} M^{\sharp} f(x) = \sup_{x \in \Omega, r > 0} |B(x, r)|^{-1} \int_{\widetilde{B}(x, r)} |f(y) - f_{\widetilde{B}(x, r)}| dy.$$

Definition 2.2. We define the $BMO_{p(\cdot),\omega}(\Omega)$ space as the set of all locally integrable functions f with finite norm

$$||f||_{BMO_{p(\cdot),\omega}} = \sup_{x \in \Omega, r > 0} \frac{||(f(\cdot) - f_{\widetilde{B}(x,r)})\chi_{\widetilde{B}(x,r)}||_{L_{\omega}^{p(\cdot)}(\Omega)}}{||\chi_{\widetilde{B}(x,r)}||_{L_{\omega}^{p(\cdot)}(\Omega)}}.$$

Theorem 2.4. [36] Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$ and ω be a Lebesgue measurable function. If $\omega \in A_{p(\cdot)}(\Omega)$, then the norms $\|\cdot\|_{BMO_{p(\cdot),\omega}}$ and $\|\cdot\|_{BMO}$ are mutually equivalent.

Everywhere in the sequel the functions $\varphi(x,r)$, $\varphi_1(x,r)$ and $\varphi_2(x,r)$ used in the body of the paper, are non-negative measurable function on $\Omega \times (0,\infty)$. We find it convenient to define the variable exponent generalized weighted Morrey spaces in the form as follows.

Definition 2.3. Let $1 \leq p(x) < \infty$, $x \in \Omega$. The variable exponent generalized Morrey space $\mathcal{M}^{p(\cdot),\varphi}(\Omega)$ and variable exponent generalized weighted Morrey space $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ are defined by the norms

$$||f||_{\mathcal{M}^{p(\cdot),\varphi}} = \sup_{x \in \Omega, r > 0} \frac{1}{\varphi(x,r)r^{\theta_p(x,r)}} ||f||_{L^{p(\cdot)}(\widetilde{B}(x,r))},$$

$$||f||_{\mathcal{M}^{p(\cdot),\varphi}_{\omega}} = \sup_{x \in \Omega, r > 0} \frac{1}{\varphi(x,r) ||\omega||_{L^{p(\cdot)}(\widetilde{B}(x,r))}} ||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,r))}.$$

According to this definition, we recover the space $\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega)$ under the choice $\varphi(x,r)=r^{\theta_p(x,r)-\frac{\lambda(x)}{p(x)}}$:

$$\mathcal{L}^{p(\cdot),\lambda(\cdot)}(\Omega) = \mathcal{M}^{p(\cdot),\varphi(\cdot)}(\Omega) \bigg|_{\varphi(x,r) = r^{\theta_p(x,r) - \frac{\lambda(x)}{p(x)}}}.$$

Definition 2.4. (Vanishing generalized weighted Morrey space) The vanishing generalized weighted Morrey space $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ is defined as the space of functions $f \in \mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ such that

$$\lim_{r\to 0}\sup_{x\in\Omega}\frac{1}{\varphi_1(x,t)\|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}}\|f\chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)}=0.$$

Everywhere in the sequel we assume that

$$\lim_{r \to 0} \frac{1}{\|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))} \inf_{x \in \Omega} \varphi(x,t)} = 0$$
(2.7)

and

$$\sup_{0 < r < \infty} \frac{1}{\|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}} \inf_{x \in \Omega} \varphi(x,t) = 0, \tag{2.8}$$

which makes the spaces $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$ non-trivial, because bounded functions with compact support belong then to this space.

Theorem 2.5. [28] Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$, $\omega \in A_{p(\cdot)}(\Omega)$ and the function $\varphi_1(x,r)$ and $\varphi_2(x,r)$ satisfy the condition

$$\sup_{t>r} \frac{\operatorname{ess inf}_{t< s<\infty} \varphi_1(x,s) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,s))}}{\|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}} \le C\varphi_2(x,r). \tag{2.9}$$

Then the maximal operator M is bounded from the space $\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ the space $\mathcal{M}^{p(\cdot),\varphi_2}_{\omega}(\Omega)$.

3 Riesz potential and its commutators in the spaces $\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$

It is well-known that the commutator is an important integral operator and it plays a key role in harmonic analysis. Let $b \in BMO(\mathbb{R}^n)$. A well known result of Chanillo [7] states that the commutator operator $[b,I^{\alpha}]f=I^{\alpha}(bf)-b\,I^{\alpha}f$ is bounded from $L_p(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$ with $1/q=1/p-\alpha/n$, $1< p< n/\alpha$.

Let $L^{\infty}(\mathbb{R}_+,v)$ be the weighted L^{∞} -space with the norm

$$||g||_{L^{\infty}(\mathbb{R}_+,v)} = \operatorname{ess \, sup}_{t>0} v(t)g(t).$$

In the sequel $\mathfrak{M}(\mathbb{R}_+)$, $\mathfrak{M}^+(\mathbb{R}_+)$ and $\mathfrak{M}^+(\mathbb{R}_+;\uparrow)$ stand for the set of Lebesgue-measurable functions on \mathbb{R}_+ , and its subspaces of nonnegative and nonnegative non-decreasing functions, respectively. We also denote

$$\mathbb{A} = \left\{ \varphi \in \mathfrak{M}^+(\mathbb{R}_+;\uparrow) : \lim_{t \to 0+} \varphi(t) = 0 \right\}.$$

Let u be a continuous and non-negative function on \mathbb{R}_+ . We define the supremal operator \overline{S}_u by

$$(\overline{S}_u g)(t) := \|u g\|_{L_1(0,t)}, \ t \in (0,\infty).$$

In the following theorem proved in [6], we use the notation

$$\widetilde{v}_1(t) = \sup_{0 < \xi < t} v_1(\xi).$$

Theorem 3.1. Suppose that v_1 and v_2 are nonnegative measurable functions such that $0 < \|v_1\|_{L_\infty(0,t)} < \infty$ for every t > 0. Let u be a continuous nonnegative function on $\mathbb R$. Then the operator \overline{S}_u is bounded from $L_\infty(\mathbb R_+,v_1)$ to $L_\infty(\mathbb R_+,v_2)$ on the cone $\mathbb A$ if and only if

$$\left\| v_2 \overline{S}_u \left(\|v_1\|_{L_{\infty}(0,\cdot)}^{-1} \right) \right\|_{L_{\infty}(\mathbb{R}_+)} < \infty.$$

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w g(t) := \int_t^1 g(s) w(s) ds, \ H_w^* g(t) := \int_t^1 \left(1 + \ln \frac{s}{t} \right) g(s) w(s) ds, \ 0 < t < \infty,$$

where w is a weight.

The following theorem was proved in [23].

Theorem 3.2. [23] Let v_1, v_2 and w be weights on $(0, \infty)$ and $v_1(t)$ be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w g(t) \le C \sup_{t>0} v_1(t) g(t)$$

holds for some C>0 for all non-negative and non-decreasing g on $(0,\infty)$ if and only if

$$B := \sup_{t>0} v_2(t) \int_t^1 \frac{w(s)ds}{\sup_{s<\tau<\infty} v_1(\tau)} < \infty.$$

Theorem 3.3. [22] Let v_1 , v_2 and w be weights on $(0, \infty)$ and $v_1(t)$ be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w^* g(t) \le C \sup_{t>0} v_1(t) g(t)$$
(3.1)

holds for some C>0 for all non-negative and non-decreasing g on $(0,\infty)$ if and only if

$$B := \sup_{t>0} v_2(t) \int_t^1 \left(1 + \ln \frac{s}{t} \right) \frac{w(s)ds}{\sup_{0 < \tau < s} v_1(\tau)} < \infty.$$

Moreover, the value C = B is the best constant for (3.1).

The following weighted local estimates are valid.

Theorem 3.4. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}^{\log}_{\infty}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$. Then

$$||I^{\alpha}f||_{L^{q(\cdot)}_{\omega}(\widetilde{B}(x,t))} \leq C||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}, \quad (3.2)$$

where C does not depend on f, x and t.

Proof. We represent f as

$$f = f_1 + f_2, \quad f_1(y) = f(y)\chi_{\widetilde{B}(x,2t)}(y), \quad f_2(y) = f(y)\chi_{\Omega\setminus\widetilde{B}(x,2t)}(y), \quad t > 0,$$
(3.3)

and have

$$I^{\alpha}f(x) = I^{\alpha}f_1(x) + I^{\alpha}f_2(x).$$

By Corollary 2.1 we obtain

$$||I^{\alpha}f_{1}||_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \leq ||I^{\alpha}f_{1}||_{L_{\omega}^{q(\cdot)}(\Omega)} \leq C||f_{1}||_{L_{\omega}^{p(\cdot)}(\Omega)} = C||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,2t))}.$$

Then

$$||I^{\alpha}f_1||_{L^{q(\cdot)}_{\omega}(\widetilde{B}(x,t))} \le C||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,2t))},$$

where the constant C is independent of f.

On the other hand,

$$\begin{split} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,2t))} &\approx |B|^{1-\frac{\alpha}{n}} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,2t))} \int_{2t}^{1} \frac{ds}{s^{n+1-\alpha}} \\ &\leq |B|^{1-\frac{\alpha}{n}} \int_{2t}^{1} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \frac{ds}{s^{n+1-\alpha}} \\ &\lesssim \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \|w^{-1}\|_{L^{p'(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{1} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \frac{ds}{s^{n+1-\alpha}} \\ &\lesssim \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{1} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega^{-1}\|_{L^{p'(\cdot)}(\widetilde{B}(x,s))} \frac{ds}{s^{n+1-\alpha}} \\ &\lesssim [\omega]_{A_{p(\cdot),q(\cdot)}} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{1} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega^{-1}\|_{L^{q(\cdot)}(\widetilde{B}(x,s))} \frac{ds}{s^{n+1-\alpha}} \end{split}$$

Taking into account that

$$||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,t))} \leq C||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

we get

$$||I^{\alpha}f_{1}||_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \leq C||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}.$$
(3.5)

When $|x-z| \le t$, $|z-y| \ge 2t$, we have $\frac{1}{2}|z-y| \le |x-y| \le \frac{3}{2}|z-y|$, and therefore

$$|I^{\alpha} f_2(x)| \leq \int_{\Omega \setminus \widetilde{B}(x,2t)} |z - y|^{\alpha - n} |f(y)| dy$$

$$\leq 2^{n - \alpha} \int_{\Omega \setminus \widetilde{B}(x,2t)} |x - y|^{\alpha - n} |f(y)| dy.$$

We obtain

$$\int_{\Omega \setminus \widetilde{B}(x,2t)} |f(y)| dy = \int_{\Omega \setminus \widetilde{B}(x,2t)} |f(y)| \left(\int_{|x-y|}^{\infty} s^{\alpha-n-1} ds \right) dy$$

$$\lesssim \int_{2t}^{\infty} s^{\alpha-n-1} \left(\int_{\{y \in \Omega: 2t \le |x-y| \le s\}} |f(y)| dy \right) ds$$

$$\lesssim \int_{t}^{\infty} s^{\alpha-n-1} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega^{-1}||_{L^{p'(\cdot)}(\widetilde{B}(x,s))} ds$$

$$\lesssim \int_{t}^{\infty} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}.$$

Hence

$$||I^{\alpha}f_{2}||_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \lesssim ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

which together with (3.5) yields (3.2).

Theorem 3.5. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}_{\infty}^{log}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$ and the functions $\varphi_1(x,t)$ and $\varphi_2(x,t)$ fulfill the condition

$$\int_{t}^{\infty} \frac{\operatorname{ess inf}_{s < r < \infty} \varphi_{1}(x, r) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} \lesssim \varphi_{2}(x, t). \tag{3.6}$$

Then the operator I^{α} is bounded from $\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ to $\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)$.

Proof. Let $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$, by condition (3.6) and Theorems 3.4, 3.2 with $v_2(r) = \varphi_2(x,r)^{-1}$, $v_1(r) = \varphi_1(x,r)^{-1} \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,r))}^{-1}$, $g(r) = \|f\|_{L^{p(\cdot)}(\widetilde{B}(x,r))}$ and $w(r) = \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,r))}^{-1} r^{-1}$ we obtain

$$\begin{split} & \|I^{\alpha}f\|_{\mathcal{M}^{q(\cdot),\varphi_{2}}_{\omega}(\Omega)} \\ & \lesssim \sup_{x \in \Omega, \ t>0} \frac{1}{\varphi_{2}(x,t)} \int_{t}^{\infty} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s} \\ & \lesssim \sup_{x \in \Omega, \ t>0} \frac{1}{\varphi_{1}(x,t) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,t))} = \|f\|_{\mathcal{M}^{p(\cdot),\varphi_{1}}_{\omega}(\Omega)}. \end{split}$$

Now we consider the commutators Riesz potential defined by

$$[b, I^{\alpha}]f(x) = \int_{\mathbb{R}^n} [b(x) - b(y)]f(y)|x - y|^{\alpha - n} dy.$$

The commutator generated by $\,M\,$ and a suitable function $\,b\,$ is formally defined by

$$[M, b]f = M(bf) - bM(f).$$

Given a measurable function b the maximal commutator is defined by

$$M_b(f)(x) := \sup_{r>0} |B(x,r)|^{-1} \int_{B(x,r)} |b(x) - b(y)| |f(y)| dy$$

for all $x \in \mathbb{R}^n$.

Lemma 3.2. [20] Let $b \in BMO(\mathbb{R}^n)$, $1 < s < \infty$. Then

$$M^{\sharp}([b,I^{\alpha}]f(x)) \leq C\|b\|_{BMO} \left[(M|I^{\alpha}f(x)|^{s})^{\frac{1}{s}} + (M^{s\alpha}|f(x)|^{s})^{\frac{1}{s}} \right],$$

where C > 0 is independed of f and x.

Lemma 3.3. [11] Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$ and $\omega \in A_{p(\cdot)}(\Omega)$. Then

$$||f\omega||_{L^{p(\cdot)}} \le C||\omega M^{\sharp}f||_{L^{p(\cdot)}}$$

with a constant C > 0 not depending on f.

Theorem 3.6. [2, Theorem 1.13] Let $b \in BMO(\mathbb{R}^n)$. Suppose that X is a Banach space of measurable functions defined on \mathbb{R}^n . Assume that M is bounded on X. Then the operator M_b is bounded on X, and the inequality

$$||M_b f||_X \le C||b||_*||f||_X$$

holds with constant C independent of f.

Corollary 3.2. Let $b \in BMO(\Omega)$, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$ and $\omega \in A_{p(\cdot)}(\Omega)$, then the operator M_b is bounded on $L^{p(\cdot)}_{\omega}(\mathbb{R}^n)$.

Theorem 3.7. [28] Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$, $\omega \in A_{p(\cdot)}(\Omega)$, $b \in BMO(\Omega)$ and the function $\varphi_1(x,r)$ and $\varphi_2(x,r)$ satisfy the condition

$$\sup_{t>r} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess inf}_{t< s<\infty} \varphi_1(x,s) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,s))}}{\|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}} \le C\varphi_2(x,r), \tag{3.7}$$

where C does not depend on $x \in \Omega$ and t. Then the operator M_b is bounded from the space $\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ to the space $\mathcal{M}^{p(\cdot),\varphi_2}_{\omega}(\Omega)$.

Theorem 3.8. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$ and $p \in \mathbb{P}^{log}_{\infty}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$. The following assertions are equivalent:

- (i) The operator $[b,I^{\alpha}]$ is bounded from $L^{p(\cdot)}_{\omega}(\Omega)$ to $L^{q(\cdot)}_{\omega}(\Omega)$.
- (ii) $b \in BMO(\Omega)$.

Proof. $(ii) \Rightarrow (i)$ Let $f \in L^{p(\cdot)}_{\omega}(\Omega)$ and $b \in BMO(\Omega)$. By the Lemma 3.3, we have

$$||[b,I^{\alpha}]f||_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim ||M^{\sharp}([b,I^{\alpha}]f)||_{L^{q(\cdot)}_{\omega}(\Omega)}$$

From Lemma 3.2, we have

$$||M^{\sharp}([b,I^{\alpha}]f)||_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim ||b||_{*} ||(M|I^{\alpha}f|^{s})^{\frac{1}{s}} + (M^{\alpha s}|f|^{s})^{\frac{1}{s}}||_{L^{q(\cdot)}_{\omega}(\Omega)}$$
$$\lesssim ||b||_{*} \left[||(M|I^{\alpha}f|^{s})^{\frac{1}{s}}||_{L^{q(\cdot)}_{\omega}(\Omega)} + ||(M^{\alpha s}|f|^{s})^{\frac{1}{s}}||_{L^{q(\cdot)}_{\omega}(\Omega)} \right].$$

By Theorem 2.1 and Corollary 2.1, we have

$$\left\| (M|I^{\alpha}f|^{s})^{\frac{1}{s}} \right\|_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim \left\| |I^{\alpha}f|^{s} \right\|_{L^{\frac{q(\cdot)}{s}}_{\omega^{sq(\cdot)}}(\Omega)}^{\frac{1}{s}} = \left\| I^{\alpha}f \right\|_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim \left\| f \right\|_{L^{p(\cdot)}_{\omega}(\Omega)}.$$

By Corollary 2.1, we have

$$\left\| (M^{\alpha s}|f|^s)^{\frac{1}{s}} \right\|_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim \|f\|_{L^{p(\cdot)}_{\omega}(\Omega)}.$$

Therefore

$$||[b, I^{\alpha}]f||_{L^{q(\cdot)}_{\omega}(\Omega)} \lesssim ||b||_* ||f||_{L^{p(\cdot)}_{\omega}(\Omega)}.$$

 $(i)\Rightarrow (ii)$ Now, let us prove the "only if" part. Let $[b,I^{lpha}]$ be bounded from $L^{p(\cdot)}_{\omega}(\Omega)$ to $L^{q(\cdot)}_{\omega}(\Omega)$, $1< p_+< rac{n}{lpha}$. Then

$$\begin{split} &|B(x,t)| \int_{\widetilde{B}(x,t)} |b(z) - b_{B(x,t)}| dz \\ &= \frac{1}{|B(x,t)|} \int_{\widetilde{B}(x,t)} \left| b(z) - \frac{1}{|B(x,t)|} \int_{\widetilde{B}(x,t)} b(y) dy \right| dz \\ &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{\widetilde{B}(x,t)} \frac{1}{|B(x,t)|^{1-\frac{\alpha}{n}}} \left| \int_{\widetilde{B}(x,t)} (b(z) - b(y)) \, dy \right| dz \\ &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{\widetilde{B}(x,t)} \left| \int_{\widetilde{B}(x,t)} (b(z) - b(y)) \, |x - y|^{\alpha - n} dy \right| dz \\ &\leq \frac{1}{|B(x,t)|^{1+\frac{\alpha}{n}}} \int_{\widetilde{B}(x,t)} \left| \left| b, I_{\alpha} \right| \chi_{B(x,t)} (z) \right| dz \\ &\leq Ct^{-n-\alpha} ||[b,I_{\alpha}] \chi_{B(x,t)}||_{L_{\omega}^{q(\cdot)}} ||\chi_{B(x,t)}||_{L_{\omega}^{q'(\cdot)}} \\ &\leq Ct^{-n-\alpha} ||\omega||_{L^{p(\cdot)}(B(x,t))} ||\omega^{-1}||_{L^{q'(\cdot)}(B(x,t))} \leq C. \end{split}$$

Hence we get

$$|B(x,t)|^{-1} \int_{\widetilde{B}(x,t)} |b(y) - b_{B(x,t)}| dy \le C.$$

This shows that $b \in BMO(\Omega)$.

The theorem has been proved.

Theorem 3.9. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}_{\infty}^{log}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$, $b \in BMO(\Omega)$. Then

$$||[b, I^{\alpha}]f||_{L^{q(\cdot)}_{\omega}(\widetilde{B}(x,t))} \le C||b||_{*}||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))}$$

$$\times \int_{t}^{\infty} \left(1 + \ln \frac{s}{t} \right) \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}, \tag{3.8}$$

where C does not depend on f, x and t.

Proof. We represent f as

$$f = f_1 + f_2, \quad f_1(y) = f(y)\chi_{\widetilde{B}(x,2t)}(y), \quad f_2(y) = f(y)\chi_{\Omega\setminus\widetilde{B}(x,2t)}(y), \quad t > 0,$$
(3.9)

and have

$$[b, I^{\alpha}]f(x) = [b, I^{\alpha}]f_1(x) + [b, I^{\alpha}]f_2(x).$$

By Theorem 3.8 we obtain

$$\begin{aligned} & \|[b, I^{\alpha}]f_1\|_{L^{q(\cdot)}_{\omega}(\widetilde{B}(x,t))} \le \|[b, I^{\alpha}]f_1\|_{L^{q(\cdot)}_{\omega}(\Omega)} \\ & \lesssim \|b\|_* \|f_1\|_{L^{p(\cdot)}_{\omega}(\Omega)} = \|b\|_* \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,2t))}. \end{aligned}$$

Then

$$||[b, I^{\alpha}]f_1||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \le C||b||_*||f||_{L^{p(\cdot)}(\widetilde{B}(x,2t))},$$

where the constant C is independent of f.

Taking into account that from the inequality (3.4) we have

$$||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,t))} \leq C||b||_{*}||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

and then

$$||[b, I^{\alpha}]f_{1}||_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \leq C||b||_{*}||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}.$$
(3.10)

When $|x-z| \le t$, $|z-y| \ge 2t$, we have $\frac{1}{2}|z-y| \le |x-y| \le \frac{3}{2}|z-y|$, and therefore

$$|[b, I^{\alpha}]f_2(x)| \leq \int_{\Omega \setminus \widetilde{B}(x,2t)} |b(y) - b(z)||z - y|^{\alpha - n}|f(y)|dy$$

$$\leq C \int_{\Omega \setminus \widetilde{B}(x,2t)} |b(y) - b(z)||x - y|^{\alpha - n}|f(y)|dy.$$

We obtain

$$\begin{split} & \int_{\Omega \setminus \widetilde{B}(x,2t)} |b(y) - b(z)| |x - y|^{\alpha - n} |f(y)| dy \\ & = \int_{\Omega \setminus \widetilde{B}(x,2t)} |b(y) - b(z)| |f(y)| \left(\int_{|x - y|}^{\infty} s^{\alpha - n - 1} ds \right) dy \\ & \leq C \int_{2t}^{\infty} s^{\alpha - n - 1} \left(\int_{\{y \in \Omega: 2t \leq |x - y| \leq s\}} |b(y) - b_{\widetilde{B}(x,t)}| |f(y)| dy \right) ds \\ & + C |b(z) - b_{\widetilde{B}(x,t)}| \int_{2t}^{\infty} s^{\alpha - n - 1} \left(\int_{\{y \in \Omega: 2t \leq |x - y| \leq s\}} |f(y)| dy \right) ds = V_1 + V_2. \end{split}$$

To estimate V_1 :

$$V_{1} = C \int_{2t}^{\infty} s^{\alpha - n - 1} \left(\int_{\{y \in \Omega: 2t \le |x - y| \le s\}} |b(y) - b_{\widetilde{B}(x,t)}| |f(y)| dy \right) ds$$

$$\leq C \int_{t}^{\infty} s^{\alpha - n - 1} ||b(\cdot) - b_{\widetilde{B}(x,s)}||_{L_{\omega}^{p'(\cdot)}(\widetilde{B}(x,s))} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ds$$

$$+ C \int_{t}^{\infty} s^{\alpha - n - 1} |b_{\widetilde{B}(x,t)} - b_{\widetilde{B}(x,s)}| \left(\int_{\widetilde{B}(x,s)} |f(y)| dy \right) ds$$

$$\leq C ||b||_{*} \int_{t}^{\infty} s^{\alpha - n - 1} ||\omega^{-1}||_{L^{p'(\cdot)}(\widetilde{B}(x,s))} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ds$$

$$+ C ||b||_{*} \int_{t}^{\infty} s^{\alpha - n - 1} \ln \frac{s}{t} ||\omega^{-1}||_{L^{p'(\cdot)}(\widetilde{B}(x,s))} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ds$$

$$\leq C ||b||_{*} \int_{t}^{\infty} \left(1 + \ln \frac{s}{t} \right) ||\omega||_{L^{p(\cdot)}(\widetilde{B}(x,s))}^{-1} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \frac{ds}{s}. \tag{3.11}$$

To estimate V_2 :

$$V_{2} = C|b(z) - b_{\widetilde{B}(x,t)}| \int_{2t}^{\infty} s^{\alpha - n - 1} \left(\int_{\{y \in \Omega: 2t \le |x - y| \le s\}} |f(y)| dy \right) ds$$

$$\leq C|B(x,t)|^{-1} \int_{\widetilde{B}(x,t)} |b(z) - b(y)| dy \int_{2t}^{\infty} ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$\leq CM_{b}\chi_{B(x,t)}(z) \int_{t}^{\infty} \left(1 + \ln \frac{s}{t} \right) ||f||_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} ||\omega||_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}, \tag{3.12}$$

where C does not depend on x, t. Then by Corollary 3.2 and (3.11), (3.12) we have

$$\|[b, I^{\alpha}]f_{2}\|_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \leq \|V_{1}\|_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} + \|V_{2}\|_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))}$$

$$\leq C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$+ C\|M_{b}\chi_{B(x,t)}\|_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$\leq C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$+ C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$\leq C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

$$\leq C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1 + \ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s} .$$

Hence

$$\begin{split} & \|[b,I^{\alpha}]f_{2}\|_{L_{\omega}^{q(\cdot)}(\widetilde{B}(x,t))} \\ \leq & C\|b\|_{*}\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{t}^{\infty} \left(1+\ln\frac{s}{t}\right) \|f\|_{L_{\omega}^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}, \end{split}$$

which together with (3.10) yields (3.8).

Theorem 3.10. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$, $b \in BMO(\Omega)$ and the functions $\varphi_1(x,t)$ and $\varphi_2(x,t)$ fulfill the condition

$$\int_{t}^{\infty} \left(1 + \ln \frac{s}{t}\right) \frac{\operatorname{ess inf}_{s < r < 1} \varphi_{1}(x, r) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} \le C\varphi_{2}(x, t). \tag{3.13}$$

Then the operators $[b,I^{\alpha}]$ is bounded from $\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ to $\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)$.

Proof. Let $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$, by condition (3.13) and Theorems 3.9, 3.3 with $v_2(r) = \varphi_2(x,r)^{-1}$, $v_1(r) = \varphi_1(x,r)^{-1} \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,r))}^{-1}$, $g(r) = \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,r))}$ and $w(r) = \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,r))}^{-1} r^{-1}$ we obtain

$$\begin{split} & \|[b,I^{\alpha}]f\|_{\mathcal{M}^{q(\cdot),\varphi_{2}}_{\omega}(\Omega)} \\ & \lesssim \|b\|_{*} \|\sup_{x \in \Omega, t > 0} \frac{1}{\varphi_{2}(x,t)} \int_{t}^{\infty} \left(1 + \ln \frac{s}{t}\right) \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s} \\ & \lesssim \|b\|_{*} \sup_{x \in \Omega, t > 0} \frac{1}{\varphi_{1}(x,t) \|\omega\|_{L^{p(\cdot)}(\widetilde{B}(x,t))}} \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,t))} = \|b\|_{*} \|\|f\|_{\mathcal{M}^{p(\cdot),\varphi_{1}}_{\omega}(\Omega)}. \end{split}$$

4 Riesz potential and its commutators in the spaces $V\mathcal{M}^{p(\cdot),\varphi}_{\omega}(\Omega)$

Theorem 4.1. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}^{log}_{\infty}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$ and the functions $\varphi_1(x,t)$ and $\varphi_2(x,t)$ fulfill the conditions

$$C_{\gamma_0} := \int_{\gamma_0}^1 \frac{\operatorname{ess inf}_{s < r < \infty} \varphi_1(x, r) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} < \infty \tag{4.1}$$

for every $\gamma_0 > 0$, and

$$\int_{t}^{\infty} \frac{\operatorname{ess inf}_{s < r < \infty} \varphi_{1}(x, r) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} \le C\varphi_{2}(x, t). \tag{4.2}$$

Then the operators I^{α} is bounded from $V\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ to $V\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)$.

Proof. The norm inequalities follow from Theorem 3.5, so we only have to prove that if

$$\lim_{r \to 0} \sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_1(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)} = 0,$$

then

$$\lim_{r \to 0} \sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_2(x, t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, t))}} \|I^{\alpha} f \chi_{\widetilde{B}(x, t)}\|_{L^{q(\cdot)}_{\omega}(\Omega)} = 0$$
 (4.3)

otherwise

To show that $\sup_{x\in\mathbb{R}^n}\frac{1}{\varphi_2(x,t)\|\omega\|_{L^q(\cdot)(\widetilde{B}(x,t))}}\|I^\alpha f\chi_{\widetilde{B}(x,t)}\|_{L^{q(\cdot)}_\omega(\Omega)}<\varepsilon \text{ for small } r\text{ , we split the right-hand side of (3.2):}$

$$\sup_{x \in \mathbb{R}^{n}} \frac{1}{\varphi_{2}(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|I^{\alpha} f \chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)} \le C_{0} \left(I_{1,\gamma_{0}}(x,t) + I_{2,\gamma_{0}}(x,t)\right), \tag{4.4}$$

where $\gamma_0>0$ will be chosen as shown below (we may take $\gamma_0<1$),

$$I_{1,\gamma_0}(x,t) := \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_t^{\gamma_0} \|f\|_{L^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

$$I_{2,\gamma_0}(x,t) := \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{\gamma_0}^{\infty} \|f\|_{L^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

and it is supposed that $t < \gamma_0$. Now we choose any fixed $\gamma_0 > 0$ such that

$$\sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_1(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)} < \frac{\varepsilon}{2CC_0}, \ \ \text{for all} \ \ 0 < t < \gamma_0,$$

where C and C_0 are constants from (4.2) and (4.4), which is possible since $f \in V\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$. Then

$$\sup_{x \in \mathbb{R}^n} CI_{1,\gamma_0}(x,t) < \frac{\varepsilon}{2}, \ 0 < t < \gamma_0,$$

by (4.2).

The estimation of the second term now may be made already by the choice of t sufficiently small thanks to the condition (4.1). We have

$$I_{2,\gamma_0}(x,t) \le C_{\gamma_0} \frac{\varphi_2(x,t)}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\|_{V\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)},$$

where C_{γ_0} is the constant from (4.1). Then, by (4.1) it suffices to choose r small enough such that

$$\frac{\varphi_2(x,t)}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} < \frac{\varepsilon}{2CC_{\gamma_0}\|f\|_{V\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)}},$$

which completes the proof of (4.3).

Theorem 4.2. Let $\Omega \subset \mathbb{R}^n$ be an open unbounded set, $0 < \alpha < n$, $p \in \mathbb{P}_{\infty}^{log}(\Omega)$, $p_+ < \frac{n}{\alpha}$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, $\omega \in A_{p(\cdot),q(\cdot)}(\Omega)$ and the functions $\varphi_1(x,t)$ and $\varphi_2(x,t)$ fulfill the conditions

$$C_{\gamma} := \int_{\gamma}^{\infty} \left(1 + \ln \frac{s}{t} \right) \frac{\operatorname{ess inf}_{s < r < \infty} \varphi_{1}(x, r) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} < \infty \tag{4.5}$$

for every γ , and

$$\int_{t}^{\infty} \left(1 + \ln \frac{s}{t}\right) \frac{\operatorname{ess inf}_{s < r < \infty} \varphi_{1}(x, r) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, r))}}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x, s))}} \frac{ds}{s} \le C\varphi_{2}(x, t). \tag{4.6}$$

Then the operators $[b,I^{\alpha}]$ is bounded from $V\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$ to $V\mathcal{M}^{q(\cdot),\varphi_2}_{\omega}(\Omega)$.

Proof. The norm inequalities follow from Theorem 3.10, so we only have to prove that

$$\lim_{r \to 0} \sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_1(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)} = 0 \Rightarrow$$

$$\lim_{r \to 0} \sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_2(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|[b,I^{\alpha}]f\chi_{\widetilde{B}(x,t)}\|_{L^{q(\cdot)}_{\omega}(\Omega)} = 0 \tag{4.7}$$

otherwise.

To show that $\sup_{x\in\mathbb{R}^n} \tfrac{1}{\varphi_2(x,t)\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|[b,I^\alpha]f\chi_{\widetilde{B}(x,t)}\|_{L^{q(\cdot)}_\omega(\Omega)} < \varepsilon \ \text{ for small } \ r \ , \text{ we split the right-hand side of (3.8):}$

$$\sup_{x \in \mathbb{R}^{n}} \frac{1}{\varphi_{2}(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|[b,I^{\alpha}]f\chi_{\widetilde{B}(x,t)}\|_{L^{q(\cdot)}_{\omega}(\Omega)} \le C_{0} \left(I_{1,\gamma}(x,r) + I_{2,\gamma}(x,r)\right), \tag{4.8}$$

where $\gamma > 0$ will be chosen as shown below (we may take $\gamma < 1$),

$$I_{1,\gamma}(x,t) := \|b\|_* \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_t^{\gamma} \left(1 + \ln \frac{s}{t}\right) \|f\|_{L^{p(\cdot)}_{\omega}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s},$$

$$I_{2,\gamma}(x,t) := \|b\|_* \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))} \int_{\gamma}^{\infty} \left(1 + \ln \frac{s}{t}\right) \|f\|_{L^{p(\cdot)}(\widetilde{B}(x,s))} \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,s))}^{-1} \frac{ds}{s}$$

and it is supposed that $\ t < \gamma$. Now we choose any fixed $\ \gamma > 0$ such that

$$\sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_1(x,t) \|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\chi_{\widetilde{B}(x,t)}\|_{L^{p(\cdot)}_{\omega}(\Omega)} < \frac{\varepsilon}{2CC_0 \|b\|_*}, \text{ for all } 0 < t < \gamma,$$

where C and C_0 are constants from (4.6) and (4.8), which is possible since $f \in V\mathcal{M}^{p(\cdot),\varphi_1}_{\omega}(\Omega)$. Then

$$\sup_{x \in \mathbb{R}^n} CI_{1,\gamma}(x,t) < \frac{\varepsilon}{2}, \ 0 < t < \gamma,$$

by (4.6).

The estimation of the second term now may be made already by the choice of r sufficiently small thanks to the condition (4.5). We have

$$I_{2,\gamma}(x,t) \le C_{\gamma} \|b\|_* \frac{\varphi_2(x,t)}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} \|f\|_{V\mathcal{M}_{\omega}^{q(\cdot),\varphi_2}(\Omega)},$$

where C_{γ} is the constant from (4.5). Then, by (4.5) it suffices to choose r small enough such that

$$\frac{\varphi_2(x,t)}{\|\omega\|_{L^{q(\cdot)}(\widetilde{B}(x,t))}} < \frac{\varepsilon}{2CC_{\gamma}\|b\|_*\|f\|_{V\mathcal{M}^{q(\cdot),\varphi_2}(\Omega)}},$$

which completes the proof of (4.6).

Acknowledgment. We thank the referee(s) for careful reading the paper and useful comments. The research of V.S. Guliyev was partially supported by the Ministry of Education and Science of the Russian Federation (the Agreement number: 02.a03.21.0008) and by the grant of 1st Azerbaijan-Russia Joint Grant Competition (Grant No. EIFBGM-4-RFTF-1/2017-21/01/1).

References

[1] D.R. Adams, A note on Riesz potentials, Duke Math. 42 (1975), 765-778.

- [2] M. Agcayazi, A. Gogatishvili, K Koca, R. Mustafayev, *A note on maximal commutators and commutators of maximal functions*, J. Math. Soc. Japan. **67**, 2 (2015), 581-593.
- [3] A. Akbulut, V.S. Guliyev, M.N. Omarova, *Marcinkiewicz integrals associated with Schrdinger operator and their commutators on vanishing generalized Morrey spaces*, Bound. Value Probl. 2017, Paper No. 121, 16 pp.
- [4] A. Almeida, J.J. Hasanov, S.G. Samko, *Maximal and potential operators in variable exponent Morrey spaces*, Georgian Math. J. **15**, 2 (2008), 1-15.
- [5] V.I. Burenkov, V.S. Guliyev, *Necessary and sufficient conditions for the bounded-ness of the Riesz potential in local Morrey-type spaces*, Potential Anal. **30**, 3 (2009), 211-249.
- [6] V. Burenkov, A. Gogatishvili, V.S. Guliyev, R. Mustafayev, *Boundedness of the fractional maximal operator in local Morrey-type spaces*, Complex Var. Elliptic Equ. **55**, 8-10 (2010), 739-758.
- [7] S. Chanillo, A note on commutators, Indiana Univ. Math. J. 31, 1 (1982) 7-16.
- [8] C. Capone, D. Cruz-Uribe, A. Fiorenza, *The fractional maximal operator and fractional integrals on variable* L_p *spaces*, Rev. Mat. Iberoam. **23** (2007), 743-770.
- [9] D. Cruz-Uribe, A. Fiorenza, J.M. Martell, C. Perez, *The boundedness of classical operators on variable* L^p *spaces*, Ann. Acad. Sci. Fenn. Math. **31** (2006), 239-264.
- [10] D. Cruz-Uribe, A. Fiorenza, *Variable Lebesgue spaces: Foundations and harmonic analysis*, Birkhauser/Springer, 2013. MR 3026953.
- [11] D. Cruz-Uribe, Li-An Daniel Wang, *Extrapolation and weighted norm inequalities in the variable Lebesgue spaces*, Trans. Amer. Math. Soc. **369**, 2 (2017), 1205-1235.
- [12] F. Chiarenza, M. Frasca, *Morrey spaces and Hardy–Littlewood maximal function*, Rend. Math. **7** (1987), 273-279.
- [13] L. Diening, Maximal functions on generalized Lebesgue spaces $L^{p(x)}$, Math. Inequal. Appl. 7, 2 (2004), 245-253.
- [14] L. Diening, P. Harjulehto, Hästö, M. Ružička. *Lebesgue and Sobolev spaces with variable exponents*, Springer-Verlag, Lecture Notes in Mathematics, vol. 2017, Berlin, 2011.
- [15] L. Diening, M. Rüźićka, Calderón-Zygmund operators on generalized Lebesgue spaces $L^{p(\cdot)}$ and problems related to fluid dynamics, J. Reine Angew. Math. **563** (2003), 197-220.

- [16] L. Diening, P. Hasto, A. Nekvinda, *Open problems in variable exponent Lebesgue and Sobolev spaces*, Function Spaces, Differential Operators and Nonlinear Analysis, Proceedings of the Conference held in Milovy, Bohemian-Moravian Uplands, May 28 June 2, 2004, Math. Inst. Acad. Sci. Czech Republick, Praha, 2005, 38-58.
- [17] F. Deringoz, V. Guliyev, S. Samko, *Boundedness of the maximal operator and its commutators on vanishing generalized Orlicz-Morrey spaces*, Ann. Acad. Sci. Fenn. Math. **40**, 2 (2015), 535-549.
- [18] A. Eroglu, Boundedness of fractional oscillatory integral operators and their commutators on generalized Morrey spaces, Bound. Value Probl. 2013, 2013:70, 12 pp.
- [19] A. Eroglu, V.S. Guliev, Dzh.V. Azizov, *Characterizations of fractional integral operators in generalized Morrey spaces on Carnot groups*, (Russian) Mat. Zametki **102**, 5 (2017), 789-804; translation in Math. Notes **102**, 5-6 (2017), 722-734.
- [20] G. Di Fazio, M. A. Ragusa, *Commutators and Morrey spaces*, Bollettino U.M.I. **7** 5-A (1991), 323-332.
- [21] V.S. Guliyev, Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces, J. Inequal. Appl. Art. ID 503948, (2009), 20 pp.
- [22] V.S. Guliyev, Generalized weighted Morrey spaces and higher order commutators of sublinear operators, Eurasian Math. J. 3, 3 (2012), 33-61.
- [23] V.S. Guliyev, Generalized local Morrey spaces and fractional integral operators with rough kernel, J. Math. Sci. (N. Y.) 193, 2 (2013), 211-227.
- [24] V.S.Guliyev, J.J. Hasanov, S.G. Samko, *Boundedness of the maximal, potential and singular operators in the generalized variable exponent Morrey spaces*, Math. Scand. **107** (2010), 285-304.
- [25] V.S. Guliyev, J.J. Hasanov, S.G. Samko, Boundedness of the maximal, potential and singular integral operators in the generalized variable exponent Morrey type spaces $\mathcal{M}^{p(\cdot),\theta(\cdot),\omega(\cdot)}(\Omega)$, J. Math. Sci. (N. Y.) **170**, 4 (2010), 423-443.
- [26] V.S. Guliev, J.J. Hasanov, S.G. Samko, *Maximal, potential and singular operators in the local "complementary" variable exponent Morrey type spaces*, J. Math. Anal. Appl. **401**, 1 (2013), 72-84.
- [27] V.S. Guliev, S.G. Samko, *Maximal, potential and singular operators in the generalized variable exponent Morrey spaces on unbounded sets*, J. Math. Sci. (N. Y.) **193**, 2 (2013), 228-248.

- [28] V.S. Guliyev, J.J. Hasanov, X.A. Badalov, *Maximal and singular integral operators and their commutators on generalized weighted Morrey spaces with variable exponent*, Math. Ineq. Appl. **21**, 1 (2018), 41-61.
- [29] P. Hästö, L. Diening, *Muckenhoupt weights in variable exponent spaces*, preprint, 2010.
- [30] J.J. Hasanov, *Hardy-Littlewood-Stein-Weiss inequality in the variable exponent Morrey spaces*, Pros. of Nat. Acad. Sci. of Azerb. **39** (48) (2013), 47-62.
- [31] V. Kokilashvili, S. Samko, On Sobolev theorem for the Riesz type potentials in Lebesgue spaces with variable exponent, Z. Anal. Anwend. 22, 4 (2003), 899-910.
- [32] V. Kokilashvili, S. Samko, Weighted boundedness of the maximal, singular and potential operators in variable exponent spaces, Analytic Methods of Analysis and Differential Equations, Cambridge Scientific Publishers, Eds: A.A.Kilbas and S.V.Rogosin, (2008), 139-164.
- [33] Y. Komori, S. Shirai, Weighted Morrey spaces and a singular integral operator, Math. Nachr. **282**, 2 (2009), 219-231.
- [34] O. Kovacik, J. Rakosnik, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, Czechoslovak Math. J. **41** (116), 4 (1991), 592-618.
- [35] T.S. Kopaliani, Infimal convolution and Muckenhoupt $A_{p(\cdot)}$ condition in variable L^p spaces, Arch. Math. **89**, 2 (2007), 185-192.
- [36] Kwok-Pun Ho, Singular integral operators, John-Nirenberg inequalities and TribelLizorkin type spaces on weighted Lebesgue spaces with variable exponents, Revista De La Union Matematica Argentina 57, 1 (2016), 85-101.
- [37] P. Long, H. Han, Characterizations of some operators on the vanishing generalized Morrey spaces with variable exponent, J. Math. Anal. Appl. 437 (2016), 419-430.
- [38] C.B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, Trans. Amer. Math. Soc. **43** (1938), 126-166.
- [39] T. Mizuhara, *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer Verlag, Tokyo (1991), 183-189.
- [40] Y. Mizuta, T. Shimomura, Sobolev embeddings for Riesz potentials of functions in Morrey spaces of variable exponent, J. Math. Japan **60** (2008), 583-602.
- [41] Y. Mizuta, T. Shimomura, Weighted Morrey spaces of variable exponent and Riesz potentials, Math. Nachr. 288, 8-9 (2015), 984-1002.

- [42] E. Nakai, *Hardy–Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*, Math. Nachr. **166** (1994), 95-103.
- [43] J. Peetre, On the theory of $\mathcal{L}_{p,\lambda}$ spaces, J. Funct. Anal. 4 (1969), 71-87.
- [44] M. A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces, J. Global Optim. 40, 1 (2008), 361-368.
- [45] M. A. Ragusa, A. Tachikawa, On interior regularity of minimizers of p(x) energy functionals, Nonlinear Anal. 93 (2013), 162-167.
- [46] N. Samko, *Maximal, potential and singular operators in vanishing generalized Morrey spaces*, J. Global Optim. **57**, 4 (2013), 1385-1399.
- [47] S. Samko, On a progress in the theory of Lebesgue spaces with variable exponent: maximal and singular operators, Integral Transforms Spec. Funct. **16**, 5-6 (2005), 461-482.
- [48] S.G. Samko, Differentiation and integration of variable order and the spaces $L^{p(x)}$, Ope-rator theory for complex and hypercomplex analysis (Mexico City, 1994), 203-219, Contemp. Math., 212, Amer. Math. Soc., Providence, RI, 1998.
- [49] I.I. Sharapudinov, *The topology of the space* $\mathcal{L}^{p(t)}([0, 1])$, Mat. Zametki **26**, 3-4 (1979), 613-632.
- [50] A. Scapellato, On some qualitative results for the solution to a Dirichlet problem in Local generalized Morrey spaces, Editor: Sivasundaram, S.; AIP Conference Proceedings Vol.1798 Article Number: UNSP 020138 (2017), doi: 10.1063/1.4972730
- [51] A. Scapellato, *Some properties of integral operators on generalized Morrey space*, Editors: Simos T.E., Tsitouras Ch.; AIP Conference Proceedings, Vol.1863, 21 Article number 510004 (2017), doi: 10.1063/1.4992662;
- [52] C. Vitanza, Functions with vanishing Morrey normand elliptic partial differential equations, in Proceedings of the Methods of Real Analysis and Partial Differential Equations, Springer, Capri, Italy, 1990, 147-150.