

A TOEPLITZ-LIKE OPERATOR WITH RATIONAL SYMBOL HAVING POLES ON THE UNIT CIRCLE III: THE ADJOINT

G.J. GROENEWALD, S. TER HORST, J. JAFTHA, AND A.C.M. RAN

ABSTRACT. This paper contains a further analysis of the Toeplitz-like operators T_ω on H^p with rational symbol ω having poles on the unit circle that were previously studied in [5, 6]. Here the adjoint operator T_ω^* is described. In the case where $p = 2$ and ω has poles only on the unit circle \mathbb{T} , a description is given for when T_ω^* is symmetric and when T_ω^* admits a selfadjoint extension. Also in the case where $p = 2$, ω has only poles on \mathbb{T} and in addition ω is proper, it is shown that T_ω^* coincides with the unbounded Toeplitz operator defined by Sarason in [12].

1. INTRODUCTION

In this paper we proceed with our study of unbounded Toeplitz-like operators on H^p with rational symbols that have poles on the unit circle \mathbb{T} which was initiated in [5]. Our previous work on such Toeplitz-like operators focused on their Fredholm properties (in [5]) and the various parts of their spectra (in [6]). Here we determine properties of the adjoint operator and conditions under which the operator is symmetric and when it has a selfadjoint extension.

Before we can define our Toeplitz-like operators, some notation has to be introduced. We write Rat for the space of rational complex functions, $\text{Rat}(\mathbb{T})$ for the subspace of Rat consisting of rational complex functions with poles only on the unit circle \mathbb{T} , and $\text{Rat}_0(\mathbb{T})$ for the subspace of strictly proper functions in $\text{Rat}(\mathbb{T})$. Now let $\omega \in \text{Rat}$, possibly with poles on \mathbb{T} . As in [5], we define the Toeplitz-like operator $T_\omega (H^p \rightarrow H^p)$, for $1 < p < \infty$, via

$$\text{Dom}(T_\omega) = \{g \in H^p \mid \omega g = f + \rho, \text{ with } f \in L^p, \rho \in \text{Rat}_0(\mathbb{T})\}, \quad T_\omega g = \mathbb{P}f. \quad (1.1)$$

Here \mathbb{P} is the Riesz projection of L^p onto H^p . The operator T_ω is densely defined and closed. In case $\omega \in \text{Rat}(\mathbb{T})$, explicit formulas for the domain, kernel, range, and a complement of the range were obtained in [6], as an extension of a result in [5] for the case where T_ω is Fredholm. We briefly recall these results in Section 2, as they will be frequently used throughout the paper.

In case ω has no poles on \mathbb{T} , in fact for any $\omega \in L^\infty$, the adjoint of the Toeplitz operator T_ω on H^p can be identified with the Toeplitz operator T_{ω^*} on $\overline{H^{p'}}$, with $1 < p' < \infty$ such that $1/p + 1/p' = 1$ and with ω^* defined as $\omega^*(z) = \overline{\omega(z)}$ on \mathbb{T} .

2010 *Mathematics Subject Classification.* Primary 47B35, 47A53; Secondary 47A68.

Key words and phrases. Toeplitz operators, unbounded operators, adjoint, symmetric operators.

This work is based on the research supported in part by the National Research Foundation of South Africa (Grant Number 90670 and 93406).

The identification of $(H^p)'$ and $H^{p'}$ goes via the usual pairing

$$\langle f, g \rangle_{p,p'} = \frac{1}{2\pi} \int_{\mathbb{T}} \overline{g(z)} f(z) dz \quad (f \in H^p, g \in H^{p'}).$$

In the sequel we use the same notation for the similarly defined pairing between L^p and $L^{p'}$ to identify $(L^p)'$ and $L^{p'}$, and in both cases the indices will often be omitted.

For the Toeplitz-like operators studied in this paper the situation is more complicated than for Toeplitz operators with L^∞ symbols. However, we do obtain that T_ω^* can be identified with the restriction of the Toeplitz-like operator T_{ω^*} on $H^{p'}$ to a dense subspace of its domain. Like for the operator T_ω , in case ω is in $\text{Rat}(\mathbb{T})$ we obtain a more explicit description of T_ω^* , which we present after introducing some further notation.

Throughout the paper \mathcal{P} denotes the space of complex polynomials and \mathcal{P}_k , for any non-negative integer k , denotes the subspace of \mathcal{P} of polynomials of degree at most k . The degree of a polynomial $r \in \mathcal{P}$ is denoted as $\deg(r)$. Given $r \in \mathcal{P}$ with $\deg(r) = k$, say $r(z) = r_0 + zr_1 + \dots + z^k r_k$, we define the polynomial $r^\#$ by

$$r^\#(z) = z^k \overline{r(1/\bar{z})} = \overline{r_0} z^k + \overline{r_1} z^{k-1} + \dots + \overline{r_k}.$$

The following theorem is our first main result.

Theorem 1.1. *Let $\omega = s/q \in \text{Rat}$ with $s, q \in \mathcal{P}$ co-prime and $1 < p < \infty$. Factor $s = s_- s_0 s_+$ and $q = q_- q_0 q_+$ with s_-, q_- having roots only inside \mathbb{T} , s_0, q_0 having roots only on \mathbb{T} , and s_+, q_+ having roots only outside \mathbb{T} . Set $m = \deg(q)$, $n = \deg(s)$, $m_\pm = \deg(q_\pm)$, $n_\pm = \deg(s_\pm)$, $m_0 = \deg(q_0)$, $n_0 = \deg(s_0)$ and let $1 < p' < \infty$ with $1/p + 1/p' = 1$. Then*

$$\text{Dom}(T_\omega^*) = (q_0)^\# H^{p'} \subset \text{Dom}(T_{\omega^*}) \quad \text{and} \quad T_\omega^* = T_{\omega^*}|_{(q_0)^\# H^{p'}}. \quad (1.2)$$

Furthermore, we have

$$\begin{aligned} \text{Ran}(T_\omega^*) &= T_{z^{m-n}(s_+)^\#/(q_+)^\#} Q_{n_0+n_- - m_0 - m_-}(s_0)^\# H^{p'}, \\ \text{Ker}(T_\omega^*) &= \left\{ \frac{(q_-)^\#(q_0)^\# r}{(s_-)^\#} \mid \deg(r) < n_- - m_- - m_0 \right\}. \end{aligned} \quad (1.3)$$

Here $Q_k = I_{H^{p'}} - P_{\mathcal{P}_{k-1}}$, with $P_{\mathcal{P}_{k-1}}$ the standard projection in $H^{p'}$ onto $\mathcal{P}_{k-1} \subset H^{p'}$ to be interpreted as 0 if $k \leq 0$, i.e., $Q_k = I_{H^{p'}}$ if $k \leq 0$. Thus, for $n_0 + n_- \leq m_0 + m_-$ we have $\text{Ran}(T_\omega^*) = T_{z^{m-n}/(q_+)^\#}(s_+ s_0)^\# H^{p'}$. Moreover,

$$\dim \text{Ker}(T_\omega^*) = \max \{0, \#\{\text{zeroes of } \omega \text{ inside } \mathbb{D}\} - \#\{\text{poles of } \omega \text{ in } \overline{\mathbb{D}}\}\},$$

where the multiplicities of the zeroes and poles are taken into account. Hence, $\dim \text{Ker}(T_\omega^*)$ is the maximum of 0 and $n_- - m_- - m_0$. In particular, T_ω^* is injective if and only if the number of poles of ω inside $\overline{\mathbb{D}}$ is greater than or equal to the number of zeroes of ω inside \mathbb{D} , multiplicities taken into account.

Before giving a proof of Theorem 1.1 in Section 4, we prove the specialization of this result for the case $\omega \in \text{Rat}(\mathbb{T})$ in Section 3. For this purpose we first provide a description of T_{ω^*} in Section 2.

The injectivity result, but not the description of $\text{Ker}(T_\omega^*)$, can also be derived from general theory and results on T_ω . Indeed, according to Theorem II.3.7 in [4], T_ω^* is injective if and only if T_ω has dense range, so that the claim follows from Proposition 2.4 in [6]. More can be obtained in this way, since H^p , $1 < p < \infty$,

is reflexive. By Theorem II.2.14 of [4] it follows that $T_{\omega}^{**} = T_{\omega}$, with the usual identifications of the dual spaces. Hence, applying the above to T_{ω}^* we find that T_{ω}^* has dense range if and only if T_{ω} is injective; see also Theorem II.4.10 in [4]. By Banach's Closed Range Theorem, cf., [14], T_{ω}^* has closed range if and only if T_{ω} has closed range. Again applying results from [6] now gives the following result.

Corollary 1.2. *Let $\omega \in \text{Rat}$ and $1 < p < \infty$. Then T_{ω}^* has closed range if and only if ω has no zeroes on \mathbb{T} , or equivalently, ω^* has no zeroes on \mathbb{T} . Moreover, T_{ω}^* has dense range if and only if*

$$\# \left\{ \begin{array}{l} \text{poles of } \omega \text{ inside } \overline{\mathbb{D}} \\ \text{multi. taken into account} \end{array} \right\} \leq \# \left\{ \begin{array}{l} \text{zeroes of } \omega \text{ inside } \overline{\mathbb{D}} \\ \text{multi. taken into account} \end{array} \right\}.$$

Beyond Section 4, and in the remainder of this introduction, we only consider the case $p = 2$ and $\omega \in \text{Rat}(\mathbb{T})$. By comparing the results on T_{ω} and T_{ω}^* it is obvious T_{ω} cannot be selfadjoint, except when ω has no poles on \mathbb{T} . In Section 5 we describe in terms of ω when T_{ω}^* is symmetric, in which case $T_{\omega}^* \subset T_{\omega}$, and whenever T_{ω}^* is symmetric we describe when T_{ω}^* admits a selfadjoint extension. The following theorem collects some of the main results of Section 5; it follows directly from Theorem 5.1, Corollaries 5.2 and 5.7, Propositions 5.4 and 5.9.

Theorem 1.3. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime. Consider T_{ω} on H^2 . Then*

$$T_{\omega}^* \text{ is symmetric} \iff \omega(\mathbb{T}) \subset \mathbb{R}.$$

In particular, if T_{ω}^ is symmetric, then $\deg(s) \leq \deg(q) \leq 2\deg(s)$. Furthermore, if T_{ω}^* is symmetric, then T_{ω}^* admits a selfadjoint extension if and only if the number of roots of $s - iq$ and $s + iq$ in \mathbb{D} , counting multiplicities, coincide. This happens in particular if $\omega(\mathbb{T}) \neq \mathbb{R}$, but cannot happen in case $\deg(q)$ is odd.*

Several other conditions for T_{ω}^* to be symmetric and/or have a selfadjoint extension are derived in Section 5.

In [12] Sarason introduced and studied an unbounded Toeplitz-like operator with symbol in the Smirnov class. In Section 6 we show that if $\omega \in \text{Rat}(\mathbb{T})$ is proper, then the adjoint operator T_{ω}^* is precisely a Toeplitz-like operator of the type studied by Sarason. Hence in this case our Toeplitz-like operator $T_{\omega} = T_{\omega}^{**}$ coincides with the adjoint of the Toeplitz-like operator considered in [12]. Based on ideas in [12], we also show that $H(\overline{\mathbb{D}})$, the space of functions analytic on a neighborhood of $\overline{\mathbb{D}}$, is contained in $\text{Dom}(T_{\omega})$ and in fact is a core of T_{ω} .

In the last section of [12], Sarason introduces a class of closed, densely defined Toeplitz-like operators on H^2 determined by algebraic properties, which was further investigated by Rosenfeld in [10, 11]. In particular, this class of Toeplitz-like operators contains the unbounded Toeplitz-like operator studied by Sarason and is closed under taking adjoints, and hence contains our Toeplitz-like operators with proper symbols in $\text{Rat}(\mathbb{T})$. In fact, we will show in Section 6 that T_{ω} is contained in the class of Toeplitz-like operators for any ω in Rat .

2. THE OPERATOR T_{ω^*} FOR $\omega \in \text{Rat}(\mathbb{T})$

In this section we recall some results from [5, 6] on the operator T_{ω} for $\omega \in \text{Rat}(\mathbb{T})$ that we will use in the sequel, and apply them to the operator T_{ω^*} . Hence, throughout this section let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ co-prime. We set $m = \deg(q)$ and $n = \deg(s)$. Furthermore, factor $s = s_- s_0 s_+$ with s_- , s_0 and s_+

polynomials having roots only inside, on, or outside \mathbb{T} , respectively. We then recall from Theorem 2.2 in [6] that

$$\begin{aligned} \text{Ker}(T_\omega) &= \{r/s_+ \mid \deg(r) < m - \deg(s_- s_0)\}; \\ \text{Dom}(T_\omega) &= qH^p + \mathcal{P}_{m-1}; \quad \text{Ran}(T_\omega) = sH^p + \tilde{\mathcal{P}}, \end{aligned} \quad (2.1)$$

where $\tilde{\mathcal{P}}$ is the subspace of \mathcal{P} given by

$$\tilde{\mathcal{P}} = \{r \in \mathcal{P} \mid rq = r_1 s + r_2 \text{ for } r_1, r_2 \in \mathcal{P}_{m-1}\} \subset \mathcal{P}_{n-1}. \quad (2.2)$$

Furthermore, $H^p = \overline{\text{Ran}(T_\omega)} + \tilde{\mathcal{Q}}$ forms a direct sum decomposition of H^p , where

$$\tilde{\mathcal{Q}} = \mathcal{P}_{k-1} \quad \text{with} \quad k = \max\{\deg(s_-) - m, 0\}, \quad (2.3)$$

following the convention $\mathcal{P}_{-1} := \{0\}$. Furthermore, the action of T_ω is as follows.

$$\begin{aligned} T_\omega g &= sh + \tilde{r} \quad (g = qh + r \in qH^p + \mathcal{P}_{m-1} = \text{Dom}(T_\omega)), \\ \text{where } \tilde{r} &\in \mathcal{P}_{n-1} \text{ is such that } rs = \tilde{r}q + r_2 \text{ for some } r_2 \in \mathcal{P}_{m-1}. \end{aligned}$$

We also recall from Lemma 5.3 in [5] that

$$T_{z^\kappa \omega} = T_{z^\kappa} T_\omega \quad \text{for any integer } \kappa \leq 0. \quad (2.4)$$

Recall that ω^* is defined as $\omega^*(z) = \overline{\omega(z)}$ on \mathbb{T} , i.e., $\omega^*(z) = \overline{s(z)}/\overline{q(z)}$. For $z \in \mathbb{T}$

$$\overline{q(z)} = \overline{q_0 + zq_1 + \cdots + z^m q_m} = \overline{q_0} + \overline{q_1} \frac{1}{z} + \cdots + \overline{q_m} \frac{1}{z^m} = \frac{1}{z^m} q^\sharp(z).$$

Hence $q^\sharp(z) = z^m \overline{q(z)}$, and likewise $s^\sharp(z) = z^n \overline{s(z)}$. Thus we have

$$\omega^*(z) = \frac{z^{m-n} s^\sharp(z)}{q^\sharp(z)} \quad \text{if } m \geq n \quad \text{and} \quad \omega^*(z) = \frac{s^\sharp(z)}{z^{n-m} q^\sharp(z)} \quad \text{if } m < n. \quad (2.5)$$

In fact, the formula $\omega^*(z) = z^{m-n} s^\sharp(z)/q^\sharp(z)$ holds in both cases, but is not always a representation as the ratio of two polynomials. Note in particular that $\omega^* \in \text{Rat}(\mathbb{T})$ in case ω is proper, while this need not be the case if ω is not proper. Thus, if ω is proper, the above formulas apply directly, while for the non-proper case, using (2.4) we can reduce certain questions to questions concerning the Toeplitz operator T_{s^\sharp/q^\sharp} with symbol s^\sharp/q^\sharp which is in $\text{Rat}(\mathbb{T})$.

A polynomial $r \neq 0$ is called self-inversive in case $r = \gamma r^\sharp$ for a constant $\gamma \in \mathbb{C}$, which necessarily is unimodular. In fact, γ is the ratio $r_0/\overline{r_n}$ with $r_0 = r(0)$ and r_n the leading coefficient of r . By a theorem of Cohn [2], a polynomial r has all its roots on \mathbb{T} if and only if r is self-inversive and its derivative has all its roots in the closed unit disc $\overline{\mathbb{D}}$. Hence, any polynomial with roots only on \mathbb{T} is self-inversive. In particular, $q = \gamma q^\sharp$ and $s_0 = \rho(s_0)^\sharp$ for unimodular constants γ and ρ .

More generally, in the transformation $r \rightarrow r^\sharp$, the nonzero roots of r (including multiplicity) transfer along the unit circle via the map $\alpha \mapsto 1/\overline{\alpha} = |\alpha|^{-2}\alpha$, while the degree decreases by the multiplicity of 0 as a root of r . Consequently, in the factorization $s^\sharp = (s_+)^\sharp (s_0)^\sharp (s_-)^\sharp$, the polynomials $(s_+)^\sharp$, $(s_0)^\sharp$ and $(s_-)^\sharp$ contain the roots of s^\sharp inside, on and outside \mathbb{T} , respectively, taking multiplicities into account. We write $(s_+)^\sharp$ rather than s_+^\sharp , etc., to avoid confusion with what one may interpret as $(s^\sharp)_+$.

We now apply the above to T_{ω^*} acting on $H^{p'}$, $1 < p' < \infty$, to fit better with the remainder of the paper.

Proposition 2.1. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ co-prime, $m = \deg(q)$ and $n = \deg(s)$. Factor $s = s_- s_0 s_+$ with s_- , s_0 and s_+ polynomials having roots only inside, on, or outside \mathbb{T} , respectively. Then for T_{ω^*} on $H^{p'}$, $1 < p' < \infty$, we have*

$$\text{Ker}(T_{\omega^*}) = \{r_0/(s_-)^{\sharp} \mid \deg(r_0) < \deg(s_-)\}, \quad \text{Dom}(T_{\omega^*}) = q^{\sharp} H^{p'} + \mathcal{P}_{m-1}. \quad (2.6)$$

Moreover, we have

$$\begin{aligned} \text{Ran}(T_{\omega^*}) &= z^{m-n} s^{\sharp} H^{p'} + \tilde{\mathcal{P}}_* \quad \text{if } m \geq n, \\ \text{Ran}(T_{\omega^*}) &= T_{z^{m-n}}(s^{\sharp} H^{p'} + \tilde{\mathcal{P}}_*) \quad \text{if } m < n, \end{aligned} \quad (2.7)$$

where for $m \geq n$ the subspace $\tilde{\mathcal{P}}_*$ is given by

$$\tilde{\mathcal{P}}_* = \{r \in \mathcal{P} \mid r q^{\sharp} = z^{m-n} r_1 s^{\sharp} + r_2 \text{ for } r_1, r_2 \in \mathcal{P}_{m-1}\} \subset \mathcal{P}_{m-n+\deg(s^{\sharp})-1}, \quad (2.8)$$

while for $m < n$ we have

$$\tilde{\mathcal{P}}_* = \{r \in \mathcal{P} \mid r q^{\sharp} = r_1 s^{\sharp} + r_2 \text{ for } r_1, r_2 \in \mathcal{P}_{m-1}\} \subset \mathcal{P}_{\deg(s^{\sharp})-1}. \quad (2.9)$$

Furthermore, $\text{Ran}(T_{\omega^*})$ is dense in $H^{p'}$.

Proof. We separate the cases $m \geq n$ and $m < n$.

For $m \geq n$, we have $\omega^* = \tilde{s}/\tilde{q} \in \text{Rat}(\mathbb{T})$ with $\tilde{s} = z^{m-n} s^{\sharp}$ and $\tilde{q} = q^{\sharp}$. Hence \tilde{s} factors as $\tilde{s} = (z^{m-n} (s_+)^{\sharp}) (s_0)^{\sharp} (s_-)^{\sharp}$, where the factors have all their roots inside, on, or outside \mathbb{T} , respectively. Also, $\deg(q^{\sharp}) = \deg(q)$ and $\deg((s_+)^{\sharp}) = \deg(s_+)$. So the formulas for $\text{Dom}(T_{\omega^*})$ and $\text{Ran}(T_{\omega^*})$ follow directly from (2.1), while the formula for $\text{Ker}(T_{\omega^*})$ follows because the bound on the degree of r_0 can be computed as

$$m - \deg(z^{m-n} (s_+)^{\sharp} (s_0)^{\sharp}) = n - \deg((s_+)^{\sharp} (s_0)^{\sharp}) = n - \deg(s_+ s_0) = \deg(s_-).$$

Finally, a complement of the closure of $\text{Ran}(T_{\omega^*})$ is given by \mathcal{P}_{k-1} with k the maximum of 0 and $\deg(z^{m-n} (s_+)^{\sharp}) - m = \deg((s_+)^{\sharp}) - n \leq 0$. Hence $\mathcal{P}_{-1} = \{0\}$. Thus T_{ω^*} has dense range, as claimed.

In case $m < n$, we have $T_{\omega^*} = T_{z^{m-n}} T_{s^{\sharp}/q^{\sharp}}$ and s^{\sharp}/q^{\sharp} is in $\text{Rat}(\mathbb{T})$. Applying the above results for T_{ω} to $T_{s^{\sharp}/q^{\sharp}}$ directly gives the formulas for $\text{Dom}(T_{\omega^*})$ and $\text{Ran}(T_{\omega^*})$.

To see that the formula for $\text{Ker}(T_{\omega^*})$ holds, we follow the argumentation of the proof of Lemma 4.1 in [5]. For $g \in \text{Dom}(T_{\omega^*}) = \text{Dom}(T_{s^{\sharp}/q^{\sharp}})$ to be in $\text{Ker}(T_{\omega^*})$ is equivalent to $T_{s^{\sharp}/q^{\sharp}} g \in \mathcal{P}_{n-m-1}$. In other words, by Lemma 3.2 in [5], to $s^{\sharp} g = q^{\sharp} \tilde{r} + r_1$ with $r_1 \in \mathcal{P}_{m-1}$ and $\tilde{r} \in \mathcal{P}_{n-m-1}$, since then $T_{s^{\sharp}/q^{\sharp}} g = \tilde{r}$. The latter happens precisely when $g = r/(s_-)^{\sharp}$ with $r \in \mathcal{P}_{\deg(s_-)-1}$. Indeed, in that case $\deg((s_+)^{\sharp} (s_0)^{\sharp} r) < n$ which in the equation $(s_+)^{\sharp} (s_0)^{\sharp} r = s^{\sharp} g = q^{\sharp} \tilde{r} + r_1$ corresponds to $\deg(\tilde{r}) < m-1$, as required. Finally, we note that a complement of $\overline{\text{Ran}(T_{s^{\sharp}/q^{\sharp}})}$ in $H^{p'}$ is given by \mathcal{P}_{k-1} with $k = \max\{0, \deg s_+^{\sharp} - m\} \leq n-m$. Let $f \in H^{p'}$ and write $z^{n-m} f = h + r \in \overline{\text{Ran}(T_{s^{\sharp}/q^{\sharp}})} + \mathcal{P}_{k-1}$. Then $f = T_{z^{m-n}} z^{n-m} f = T_{z^{m-n}}(h + r) = T_{z^{m-n}} h \in T_{z^{m-n}} \overline{\text{Ran}(T_{s^{\sharp}/q^{\sharp}})} \subset \overline{\text{Ran}(T_{z^{m-n}} T_{s^{\sharp}/q^{\sharp}})} = \overline{\text{Ran}(T_{\omega^*})}$. Thus also in this case $\text{Ran}(T_{\omega^*})$ is dense in $H^{p'}$. \square

We conclude this section with a lemma will be of use in the sequel.

Lemma 2.2. *Let $r_1, r_2 \in \mathcal{P}$. Set $n_i = \deg(r_i)$, for $i = 1, 2$, and $n = \deg(r_1 + r_2)$. Then*

$$(r_1 + r_2)^{\sharp} = z^{n-n_1} r_1^{\sharp} + z^{n-n_2} r_2^{\sharp}.$$

In case $n < \max\{n_1, n_2\}$, then $n_1 = n_2$ and 0 is a root of $r_1^\sharp + r_2^\sharp$ with multiplicity $n - n_1$, so that the left hand side in the above identity still is a polynomial without a root at 0.

Proof. By definition, for $z \in \mathbb{T}$ we have

$$\begin{aligned} (r_1 + r_2)^\sharp(z) &= z^n (\overline{r_1(1/\bar{z})} + \overline{r_2(1/\bar{z})}) = z^{n-n_1} z^{n_1} \overline{r_1(1/\bar{z})} + z^{n-n_2} z^{n_2} \overline{r_2(1/\bar{z})} \\ &= z^{n-n_1} r_1^\sharp(z) + z^{n-n_2} r_2^\sharp(z). \end{aligned} \quad \square$$

3. THE ADJOINT OF T_ω FOR $\omega \in \text{Rat}(\mathbb{T})$

In this section we prove the first main result, Theorem 1.1, for the special case that $\omega \in \text{Rat}(\mathbb{T})$. In this case, the result specializes to the following theorem, which we prove in this section.

Theorem 3.1. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime and $1 < p < \infty$. Set $m = \deg(q)$ and $n = \deg(s)$ and let $1 < p' < \infty$ with $1/p + 1/p' = 1$. Then*

$$\text{Dom}(T_\omega^*) = q^\sharp H^{p'} \subset \text{Dom}(T_\omega^*) \quad \text{and} \quad T_\omega^* = T_\omega^*|_{q^\sharp H^{p'}}. \quad (3.1)$$

In fact, for $g = q^\sharp v \in q^\sharp H^{p'}$ we have $T_\omega^ g = T_{z^{m-n} s^\sharp} v$. Moreover, factorize $s = s_- s_0 s_+$ with s_- , s_0 and s_+ polynomials having roots only inside, on, or outside \mathbb{T} , respectively. Then*

$$\text{Ran}(T_\omega^*) = T_{z^{m-n} s^\sharp} H^{p'} \quad \text{and} \quad \text{Ker}(T_\omega^*) = \left\{ \frac{q^\sharp r}{(s_-)^\sharp} \mid \deg(r) < \deg(s_-) - m \right\}. \quad (3.2)$$

In particular, we have

$$\dim \text{Ker}(T_\omega^*) = \max \{0, \# \{\text{zeroes of } \omega^* \text{ outside } \mathbb{T}\} - \# \{\text{poles of } \omega^* \text{ on } \mathbb{T}\}\},$$

where the multiplicities of the zeroes and poles are taken into account. Thus T_ω^ is injective if and only if ω has at least as many poles inside \mathbb{T} as zeroes inside \mathbb{T} unequal to 0, multiplicities taken into account.*

We first present some auxiliary lemmas. Throughout, let $1 < p, p' < \infty$ such that $1/p + 1/p' = 1$. We will consider T_ω as an operator with domain in H^p and T_ω^* as an operator with domain in $H^{p'}$.

Lemma 3.2. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime, $m = \deg(q)$ and $n = \deg(s)$. Then*

$$q^\sharp H^{p'} \subset \text{Dom}(T_\omega^*) \cap \text{Dom}(T_\omega^*) \quad \text{and} \quad T_\omega^*|_{q^\sharp H^{p'}} = T_\omega^*|_{q^\sharp H^{p'}}.$$

Moreover, for $g = q^\sharp v \in q^\sharp H^{p'}$, with $v \in H^{p'}$, we have $T_\omega^ g = T_{z^{m-n} s^\sharp} v$, and thus $T_\omega^*(q^\sharp H^{p'}) = T_{z^{m-n} s^\sharp} H^{p'}$.*

Proof. The inclusion $q^\sharp H^{p'} \subset \text{Dom}(T_\omega^*)$ follows from Proposition 2.1. Let g be in $q^\sharp H^{p'}$, say $g(z) = q^\sharp(z)v(z)$ for $v \in H^{p'}$. We show that for $f \in \text{Dom}(T_\omega)$ we have $\langle T_\omega f, g \rangle_{p,p'} = \langle f, T_\omega^* g \rangle_{p,p'}$. Let $f \in \text{Dom}(T_\omega)$ and $h = T_\omega f \in H^p$, i.e., $sf = qh + r$ for some $r \in \mathcal{P}_{m-1}$, by [5, Lemma 2.3]. Then

$$\begin{aligned} \langle T_\omega f, g \rangle_{p,p'} &= \langle h, q^\sharp v \rangle_{p,p'} = \langle h, z^m \bar{q} v \rangle_{p,p'} = \langle qh, z^m v \rangle_{p,p'} \\ &= \langle sf - r, z^m v \rangle_{p,p'} = \langle sf, z^m v \rangle_{p,p'} \quad (\text{because } \deg(r) < m, v \in H^{p'}) \\ &= \langle f, z^m \bar{s} v \rangle_{p,p'} = \langle f, z^{m-n} s^\sharp v \rangle_{p,p'} = \langle f, T_{z^{m-n} s^\sharp} v \rangle_{p,p'} \quad (\text{because } f \in H^p). \end{aligned}$$

It remains to show that $T_{\omega^*}g = T_{z^{m-n}s^\sharp}v$. If $m \geq n$, then $\omega^* = z^{m-n}s^\sharp/q^\sharp$ is in $\text{Rat}(\mathbb{T})$ and $\omega^*g = z^{m-n}s^\sharp v \in H^{p'}$, so that, $T_{\omega^*}g = z^{m-n}s^\sharp v = T_{z^{m-n}s^\sharp}v$, by Lemma 2.3 in [5]. In case $m < n$, we have $T_{\omega^*}g = T_{z^{m-n}s^\sharp/q^\sharp}g = T_{z^{m-n}s^\sharp}v$. \square

Lemma 3.3. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime, $m = \deg(q)$ and $n = \deg(s)$. Let $g \in \text{Dom}(T_\omega^*)$ and $k = T_\omega^*g \in H^{p'}$. Then for any $r \in \mathcal{P}_{n-1}$ and $r_1 \in \mathcal{P}_{m-1}$ so that*

$$sr_1 = qr + r_2 \text{ for some } r_2 \in \mathcal{P}_{m-1} \quad (3.3)$$

we have

$$\langle r_1, k \rangle_{p,p'} = \langle r, g \rangle_{p,p'}.$$

Moreover, we have

$$z^{m-n}s^\sharp g - q^\sharp k \in \mathcal{P}_{m-1} \text{ if } m \geq n \text{ and } s^\sharp g - z^{n-m}q^\sharp k \in \mathcal{P}_{n-1} \text{ if } m < n. \quad (3.4)$$

In particular, $\text{Dom}(T_\omega^*) \subset \text{Dom}(T_{\omega^*})$ and $T_\omega^* = T_{\omega^*}|_{\text{Dom}(T_\omega^*)}$.

Proof. Let $g \in \text{Dom}(T_\omega^*)$ and $k = T_\omega^*g$. Hence $\langle T_\omega f, g \rangle_{p,p'} = \langle f, k \rangle_{p,p'}$ for each $f \in \text{Dom}(T_\omega)$. Since $\omega \in \text{Rat}(\mathbb{T})$, we have $\text{Dom}(T_\omega) = qH^p + \mathcal{P}_{m-1}$. Let $f = qh + r_1 \in \text{Dom}(T_\omega)$, with $h \in H^p$ and $r_1 \in \mathcal{P}_{m-1}$. Then $T_\omega f = sh + r$ where $r \in \mathcal{P}_{n-1}$ is uniquely determined by (3.3). Thus

$$\langle sh, g \rangle + \langle r, g \rangle = \langle sh + r, g \rangle = \langle T_\omega f, g \rangle = \langle f, k \rangle = \langle qh + r_1, k \rangle = \langle qh, k \rangle + \langle r_1, k \rangle.$$

We obtain that

$$\langle sh, g \rangle - \langle qh, k \rangle = \langle r_1, k \rangle - \langle r, g \rangle.$$

However, in choosing $f \in \text{Dom}(T_\omega)$ we can choose $h \in H^p$ and $r_1 \in \mathcal{P}_{m-1}$ independently, and in particular set one or the other equal to zero, resulting in

$$\langle sh, g \rangle = \langle qh, k \rangle \quad (h \in H^p), \quad \langle r_1, k \rangle = \langle r, g \rangle \quad (r \in \mathcal{P}_{n-1}, r_1 \in \mathcal{P}_{m-1} \text{ as in (3.3)}).$$

The second identity proves the first claim of the lemma. From the first identity we obtain that

$$0 = \langle h, \overline{s}g - \overline{q}k \rangle_{p,p'} = \langle h, z^{-n}s^\sharp g - z^{-m}q^\sharp k \rangle_{p,p'} \quad (h \in H^p).$$

Thus $\mathbb{P}(z^{-n}s^\sharp g - z^{-m}q^\sharp k) = 0$. On the other hand, for $l = \max\{m, n\}$ we have

$$z^l(z^{-n}s^\sharp g - z^{-m}q^\sharp k) = z^{l-n}s^\sharp g - z^{l-m}q^\sharp k \in H^{p'}.$$

This can only occur if $z^{l-n}s^\sharp g - z^{l-m}q^\sharp k \in \mathcal{P}_{l-1}$, which proves the second claim.

To complete the proof, we show that $g \in \text{Dom}(T_{\omega^*})$ and $T_{\omega^*}g = k$. For $m \geq n$ we have $\omega^* \in \text{Rat}(\mathbb{T})$ and the first inclusion of (3.4) can be rewritten as

$$\omega^*g = \left(\frac{z^{m-n}s^\sharp}{q^\sharp} \right) g = k + \tilde{r}/q^\sharp, \quad \text{for some } \tilde{r} \in \mathcal{P}_{m-1}.$$

Since $\deg(q^\sharp) = \deg(q) = m$, it now follows that $g \in \text{Dom}(T_{\omega^*})$ and $T_{\omega^*}g = k$. In case $m < n$ we have $T_{\omega^*} = T_{z^{m-n}s^\sharp/q^\sharp}$ and $s^\sharp/q^\sharp \in \text{Rat}(\mathbb{T})$. Now the second inclusion of (3.4) gives

$$\left(\frac{s^\sharp}{q^\sharp} \right) g = z^{n-m}k + \tilde{r}/q^\sharp, \quad \text{for some } \tilde{r} \in \mathcal{P}_{n-1}.$$

Write $\tilde{r} = \tilde{r}_1 q^\sharp + \tilde{r}_2$ with $\tilde{r}_2 \in \mathcal{P}_{m-1}$. Then $\tilde{r}/q^\sharp = \tilde{r}_1 + \tilde{r}_2/q^\sharp$ and $\deg(\tilde{r}_1) < m - n$. Since $\tilde{r}_2/q^\sharp \in \text{Rat}_0(\mathbb{T})$ it follows that $g \in \text{Dom}(T_{s^\sharp/q^\sharp}) = \text{Dom}(T_{\omega^*})$ and $T_{s^\sharp/q^\sharp}g = z^{n-m}k + \tilde{r}_1$. But then $T_{\omega^*}g = T_{z^{m-n}s^\sharp/q^\sharp}g = T_{z^{m-n}}(z^{n-m}k + \tilde{r}_1) = k$. \square

A special case of the following result was proven as part of the proof of Theorem 2.2 in [6].

Lemma 3.4. *Let $r, \tilde{r} \in \mathcal{P}$ be co-prime. Then $rH^p \cap \tilde{r}H^p = r\tilde{r}H^p$.*

Proof. Let $\tilde{r}f = rg$ with $f, g \in H^p$. Then $f = r \cdot g/\tilde{r} \in H^p$, so we should show $\tilde{f} := g/\tilde{r} \in H^p$, i.e., \tilde{f} analytic on \mathbb{D} and $\int_{\mathbb{T}} |\tilde{f}(z)|^p dz < \infty$.

Since $g \in H^p$, the function \tilde{f} can only fail to be analytic at the roots of \tilde{r} inside \mathbb{D} . However, if this were the case, then $f = r\tilde{f}$ would also fail to be analytic in \mathbb{D} , since r and \tilde{r} are co-prime. Thus \tilde{f} is analytic on \mathbb{D} .

Divide \mathbb{T} as $\mathbb{T}_1 \cup \mathbb{T}_2$ with $\mathbb{T}_1 \cap \mathbb{T}_2 = \emptyset$ in such a way that \mathbb{T}_1 and \mathbb{T}_2 are both nonempty finite unions of line segments of \mathbb{T} so that the interior of \mathbb{T}_1 contains the roots of r and the interior of \mathbb{T}_2 the roots of \tilde{r} . Then $|\tilde{r}(z)| > N_1$ on \mathbb{T}_1 and $|r(z)| > N_2$ on \mathbb{T}_2 for some $N_1, N_2 > 0$. Note that $f = rf$ and $g = \tilde{r}\tilde{f}$. We then obtain

$$\int_{\mathbb{T}_2} |\tilde{f}(z)|^p dz = \int_{\mathbb{T}_2} |f(z)/r(z)|^p dz \leq N_2^{-p} \int_{\mathbb{T}_2} |f(z)|^p dz \leq (2\pi N_2^p)^{-1} \|f\|_{H^p}^p.$$

Using $g = \tilde{r}\tilde{f}$, one obtains similarly that $\int_{\mathbb{T}_1} |\tilde{f}(z)|^p dz \leq (2\pi N_1^p)^{-1} \|g\|_{H^p}^p$. Thus $\int_{\mathbb{T}} |\tilde{f}(z)|^p dz < \infty$. \square

Proof of Theorem 3.1. By Lemma 3.2, in order to prove (3.1), the formula for the action of T_ω^* on $q^\sharp H^{p'}$ and for the range of T_ω^* in (3.2), it remains to show that $\text{Dom}(T_\omega^*) \subset q^\sharp H^{p'}$.

View \mathcal{P} and \mathcal{P}_k , $k = 1, 2, \dots$, as subspaces of H^p or $H^{p'}$, write P_k for the projection onto \mathcal{P}_{k-1} and set $Q_k = I - P_k$. Also, the standard $k \times k$ compression of a Toeplitz operator T_ϕ on H^p (or $H^{p'}$) is denoted by $T_{\phi,k}$, i.e., $T_{\phi,k} = P_k T_\phi|_{\mathcal{P}_{k-1}}$. Now, the relation (3.3) between $r \in \mathcal{P}_{n-1}$ and $r_1 \in \mathcal{P}_{m-1}$ can be rewritten as

$$T_s r_1 - T_q r \in \mathcal{P}_{m-1},$$

or, equivalently, as

$$Q_m T_s P_m r_1 = Q_m T_s r_1 = Q_m T_q r = Q_m T_q P_n r. \quad (3.5)$$

We now consider the cases $m \geq n$ and $m < n$ separately.

First assume $m \geq n$. We can then decompose $Q_m T_s P_m$ and $Q_m T_q P_n$ as

$$\begin{aligned} Q_m T_s P_m &= \begin{bmatrix} 0 & T_{s^\sharp, n}^* T_{z^{m-n}}^* \\ 0 & 0 \end{bmatrix} : \mathcal{P}_{m-1} = \begin{bmatrix} \mathcal{P}_{m-n} \\ T_{z^{m-n}} \mathcal{P}_{n-1} \end{bmatrix} \rightarrow \begin{bmatrix} \mathcal{P}_{n-1} \\ T_z^n H^p \end{bmatrix}, \\ Q_m T_q P_n &= \begin{bmatrix} T_{q^\sharp, n}^* \\ 0 \end{bmatrix} : \mathcal{P}_{n-1} \rightarrow \begin{bmatrix} \mathcal{P}_{n-1} \\ T_{z^n} H^p \end{bmatrix}. \end{aligned}$$

Hence, in this case the identity in (3.5) can be write as

$$T_{s^\sharp, n}^* (T_{z^{m-n}}^* r_1) = T_{q^\sharp, n}^* r.$$

Since all Toeplitz matrices are upper triangular, we in fact have

$$T_{s^\sharp, m}^* T_{z^{m-n}, m}^* r_1 = T_{q^\sharp, m}^* r.$$

Note that $T_{q^\sharp, n}^*$ is invertible, because q has only roots on \mathbb{T} so that $q(0) \neq 0$. We obtain that for given $r_1 \in \mathcal{P}_{m-1}$, the polynomial $r \in \mathcal{P}_{n-1}$ that satisfies (3.3) is

uniquely determined by

$$r = (T_{q^\sharp, m}^*)^{-1} T_{s^\sharp, m}^* T_{z^{m-n}, m}^* r_1 = T_{s^\sharp, m}^* T_{z, m}^{*m-n} (T_{q^\sharp, m}^*)^{-1} r_1,$$

where the commutation of Toeplitz matrices can occur since they all have analytic symbols. Now take $r_1 \in \mathcal{P}_{m-1}$ arbitrary, and define r as above, so that (3.3) holds. Then, by Lemma 3.3, we have

$$\begin{aligned} \langle r_1, P_m k \rangle_{\mathcal{P}_{m-1}} &= \langle r_1, k \rangle_{p, p'} = \langle r, g \rangle_{p, p'} = \langle r, P_m g \rangle_{\mathcal{P}_{m-1}} \\ &= \langle T_{s^\sharp, m}^* T_{z, m}^{*m-n} (T_{q^\sharp, m}^*)^{-1} r_1, P_m g \rangle_{\mathcal{P}_{m-1}} \\ &= \langle r_1, (T_{q^\sharp, m}^*)^{-1} T_{z, m}^{*m-n} T_{s^\sharp, m}^* P_m g \rangle_{\mathcal{P}_{m-1}}. \end{aligned}$$

Since $r_1 \in \mathcal{P}_{m-1}$ is arbitrary, we obtain that $P_m k = (T_{q^\sharp, m}^*)^{-1} T_{z, m}^{*m-n} T_{s^\sharp, m}^* P_m g$, and thus

$$P_m T_{q^\sharp} k = T_{q^\sharp, m} P_m k = T_{z, m}^{*m-n} T_{s^\sharp, m}^* P_m g = P_m T_z^{*m-n} T_{s^\sharp} g.$$

This shows that $P_m q^\sharp k = P_m z^{m-n} s^\sharp g$. Together with the first inclusion in (3.4) we obtain that

$$q^\sharp k = z^{m-n} s^\sharp g.$$

Since q^\sharp and $z^{m-n} s^\sharp$ are co-prime, we can apply Lemma 3.4 to conclude $g \in q^\sharp H^{p'}$.

Next assume $m < n$. We can then write $\omega = \omega_0 + \omega_1$ uniquely with $\omega_0 \in \text{Rat}_0(\mathbb{T})$ and $\omega_1 \in \text{Rat}$ with no poles on \mathbb{T} , i.e. $\omega_1 \in L^\infty(\mathbb{T})$, see [5, Lemma 2.4]. In fact $\omega_1 \in \mathcal{P}$, since all poles of ω are on \mathbb{T} , and $\omega_0 = \tilde{s}/q$ with $\tilde{s} \in \mathcal{P}_{m-1}$. It now follows that $\text{Dom}(T_{\omega_0}^*) = q^\sharp H^{p'}$, and since T_{ω_1} is bounded, $\text{Dom}(T_\omega^*) = \text{Dom}(T_{\omega_0}^*) = q^\sharp H^{p'}$. Furthermore, $T_\omega^* = T_{\omega_0}^* + T_{\omega_1}^*|_{q^\sharp H^{p'}} = T_{\omega_0^*}^*|_{q^\sharp H^{p'}} + T_{\omega_1^*}^*|_{q^\sharp H^{p'}} = T_{\omega^*}^*|_{q^\sharp H^{p'}}$.

In the next part of the proof we prove the formula for $\text{Ker}(T_{\omega^*})$, without distinguishing between the proper and non-proper case. Let $g = q^\sharp v \in \text{Dom}(T_\omega^*)$ with $v \in H^{p'}$. Then $g \in \text{Ker}(T_\omega^*)$ if and only if $g \in \text{Ker}(T_{\omega^*})$, i.e., $g = q^\sharp v = r_1/(s_-)^\sharp$ for $r_1 \in \mathcal{P}_{\deg(s_-)-1}$, see Proposition 2.1. Thus $v = r_1/((s_-)^\sharp q^\sharp) \in \text{Rat} \cap H^{p'}$. Then $v \in H^{p'}$ implies $r_1 = q^\sharp r$, and $\deg(r) = \deg(r_1) - m < \deg(s_-) - m$. Hence $g = q^\sharp r/(s_-)^\sharp$ with $\deg(r) < \deg(s_-) - m$. That all such functions are in $\text{Ker}(T_\omega^*) = \text{Ker}(T_{\omega^*}) \cap q^\sharp H^{p'}$ follows directly from the formula for $\text{Ker}(T_{\omega^*})$ obtained in Proposition 2.1. The formula for the dimension of $\text{Ker}(T_\omega^*)$ follows directly and the condition for injectivity follows since $\deg(s_-)^\sharp$ is equal to the number on nonzero roots of s_- , counting multiplicity. \square

4. THE ADJOINT OF T_ω : GENERAL CASE

In the section we prove Theorem 1.1 in full generality. Hence let $\omega = s/q \in \text{Rat}$ with $s, q \in \mathcal{P}$ co-prime. As in Theorem 1.1, factor $s = s_- s_0 s_+$ and $q = q_- q_0 q_+$ with s_-, q_- having roots only inside \mathbb{T} , s_0, q_0 having roots only on \mathbb{T} , and s_+, q_+ having roots only outside \mathbb{T} . Set $m = \deg(q)$, $n = \deg(s)$, $m_\pm = \deg(q_\pm)$, $n_\pm = \deg(s_\pm)$, and $m_0 = \deg(q_0)$, $n_0 = \deg(s_0)$. By Lemma 5.1 in [5], and its proof, we can factor ω as $\omega = \omega_- (z^\kappa \omega_0) \omega_+$ with $\kappa = n_- - m_-$, $\omega_- = s_-/(z^\kappa q_-)$ having only poles and zeroes inside \mathbb{T} , $\omega_0 = s_0/q_0$ having only poles and zeroes on \mathbb{T} , and $\omega_+ = s_+/q_+$ having only poles and zeroes outside \mathbb{T} , and we have $T_\omega = T_{\omega_-} T_{z^\kappa \omega_0} T_{\omega_+}$. Moreover, T_{ω_-} and T_{ω_+} are bounded and boundedly invertible.

Note that $T_{\omega_-} T_{z^\kappa \omega_0}$ is closed and densely defined and $\text{Ran}(T_{\omega_+}) = H^p$, and thus by Corollary 1 in [13]

$$T_\omega^* = T_{\omega_+}^* (T_{\omega_-} T_{z^\kappa \omega_0})^*.$$

Furthermore, T_{ω_-} is bounded and $T_{z^\kappa \omega_0}$ is closed and densely defined. By Theorem 4 in [1] one has

$$(T_{\omega_-} T_{z^\kappa \omega_0})^* = T_{z^\kappa \omega_0}^* T_{\omega_-}^*.$$

Combining this and using that $T_{\omega_+}^* = T_{\omega_+}^*$ and $T_{\omega_-}^* = T_{\omega_-}^*$ we see that

$$T_\omega^* = T_{\omega_+}^* T_{z^\kappa \omega_0}^* T_{\omega_-}^* = T_{\omega_+}^* T_{z^\kappa \omega_0}^* T_{\omega_-}^* \quad \text{on } \text{Dom}(T_\omega^*).$$

Note that

$$\omega_-^* = \frac{(s_-)^\sharp}{(q_-)^\sharp}, \quad \omega_0^* = z^{m_0-n_0} \frac{(s_0)^\sharp}{(q_0)^\sharp}, \quad (z^\kappa \omega_0)^* = z^{m_0-n_0-\kappa} \frac{(s_0)^\sharp}{(q_0)^\sharp}, \quad \omega_+^* = z^{m_+-n_+} \frac{(s_+)^\sharp}{(q_+)^\sharp}.$$

By construction, ω_- and $1/\omega_-$ are both anti-analytic. Consequently, ω_-^* and $1/\omega_-^*$ are both analytic functions. This implies $T_{\omega_-}^*(q_0)^\sharp H^{p'} \subset (q_0)^\sharp H^{p'}$, and thus $T_{\omega_-}^*(q_0)^\sharp H^{p'} = (q_0)^\sharp H^{p'}$. Since $T_{\omega_+}^*$ is invertible, to see that $\text{Dom}(T_\omega^*) = (q_0)^\sharp H^{p'}$ it suffices to show $\text{Dom}(T_{z^\kappa \omega_0}^*) = (q_0)^\sharp H^{p'}$. For the case where $\kappa \geq 0$, so that $z^\kappa \omega_0 \in \text{Rat}(\mathbb{T})$, this follows directly from Theorem 3.1. For $\kappa < 0$, note that $T_{z^\kappa \omega_0} = T_{z^\kappa} T_{\omega_0}$, so that $T_{z^\kappa \omega_0}^* = T_{\omega_0}^* T_{z^\kappa}^* = T_{\omega_0}^* T_{z^{-\kappa}}$, again using Theorem 4 of [1]. Then $g \in \text{Dom}(T_{z^\kappa \omega_0}^*)$ holds if and only if $z^{-\kappa} g \in \text{Dom}(T_{\omega_0}^*) = (q_0)^\sharp H^{p'}$. By Lemma 3.4 this is the same as $g \in (q_0)^\sharp H^{p'}$, since $z^{-\kappa}$ and q_0^\sharp are co-prime. Thus in both cases we arrive at $\text{Dom}(T_\omega^*) = (q_0)^\sharp H^{p'}$. Moreover, we also find that $T_{z^\kappa \omega_0}^* = T_{(z^\kappa \omega_0)^*} |_{(q_0)^\sharp H^{p'}}$, so that

$$T_\omega^* = T_{\omega_+}^* T_{z^\kappa \omega_0}^* T_{\omega_-}^* = T_{\omega_+}^* T_{(z^\kappa \omega_0)^*} T_{\omega_-}^* |_{(q_0)^\sharp H^{p'}} = T_{\omega^*} |_{(q_0)^\sharp H^{p'}}.$$

Hence (1.2) holds.

Next we derive the formula for $\text{Ker}(T_\omega^*)$. For $\kappa \geq 0$ we have $g \in \text{Ker}(T_\omega^*)$ if and only if $T_{\omega_-}^* g \in \text{Ker}(T_{z^\kappa \omega_0}^*) = (q_0)^\sharp \mathcal{P}_{\kappa-m_0-1}$, where the last identity follows by applying Theorem 3.1 to $z^\kappa \omega_0$. Thus $g \in \text{Ker}(T_\omega^*)$ if and only if $((s_-)^\sharp / (q_-)^\sharp) g = (q_0)^\sharp r$, i.e., $g = (q_-)^\sharp (q_0)^\sharp r / (s_-)^\sharp$, for some $r \in \mathcal{P}_{\kappa-m_0-1}$, as claimed. For $\kappa < 0$ we have $g \in \text{Ker}(T_\omega^*)$ if and only if $z^{-\kappa} \omega_-^* g \in \text{Ker}(T_{\omega_0}^*)$. However, $\text{Ker}(T_{\omega_0}^*) = \{0\}$, by Theorem 3.1, so that $\text{Ker}(T_\omega^*) = \{0\}$, in line with the formula in (1.3). The formula for the dimension of $\text{Ker}(T_\omega^*)$ follows directly.

Now we turn to the formula for $\text{Ran}(T_\omega^*)$. Note that

$$\text{Ran}(T_\omega^*) = T_{\omega_+}^* \text{Ran}(T_{z^\kappa \omega_0}^* T_{\omega_-}^*) = T_{\omega_+}^* \text{Ran}(T_{z^\kappa \omega_0}^*). \quad (4.1)$$

We first show that $\text{Ran}(T_{z^\kappa \omega_0}^*) = T_{z^{m_0-n_0-\kappa}}(s_0)^\sharp H^{p'}$. Again, for the case $\kappa \geq 0$ this follows directly from Theorem 3.1. Assume $\kappa < 0$. Then $T_{z^\kappa \omega_0}^* = T_{\omega_0}^* T_{z^{-\kappa}}$. Hence,

$$\begin{aligned} \text{Ran}(T_{z^\kappa \omega_0}^*) &= T_{\omega_0}^* (z^{-\kappa} H^{p'} \cap \text{Dom}(T_{\omega_0})) = T_{\omega_0}^* (z^{-\kappa} H^{p'} \cap (q_0)^\sharp H^{p'}) \\ &= T_{\omega_0}^* z^{-\kappa} (q_0)^\sharp H^{p'}. \end{aligned}$$

The last identity follows by Lemma 3.4. Now the action of $T_{\omega_0}^*$, as described in Theorem 3.1, shows that $\text{Ran}(T_{z^\kappa \omega_0}^*) = T_{z^{m_0-n_0-\kappa}}(s_0)^\sharp H^{p'} = T_{z^{m_0-n_0-\kappa}}(s_0)^\sharp H^{p'}$. Since $1/q_+$ is analytic, $1/(q_+)^\sharp$ is anti-analytic, and therefore, independent of the sign of $m_+ - n_+$, we have

$$T_{\omega_+}^* = T_{1/(q_+)^\sharp} T_{z^{m_+-n_+}} T_{(s_+)^\sharp}.$$

Thus

$$\text{Ran}(T_\omega^*) = T_{1/(q_+)^\sharp} T_{z^{m_+-n_+}} T_{(s_+)^\sharp} T_{z^{m_0-n_0-\kappa}}(s_0)^\sharp H^{p'}.$$

Note that $T_{(s_+)^{\sharp}}$ and $T_{z^{m_0-n_0-\kappa}}$ need not commute, in case $m_0 - n_0 - \kappa < 0$. However, we do have $T_{(s_+)^{\sharp}} T_{z^{m_0-n_0-\kappa}} = T_{z^{m_0-n_0-\kappa}} T_{(s_+)^{\sharp}} Q_{\kappa+n_0-m_0}$. Moreover, since $(s_+)^{\sharp}$ is analytic, $T_{(s_+)^{\sharp}} Q_{\kappa+n_0-m_0} = Q_{\kappa+n_0-m_0} T_{(s_+)^{\sharp}} Q_{\kappa+n_0-m_0}$ and we have

$$T_{z^{m_0-n_0-\kappa}} T_{z^{m_0-n_0-\kappa}} Q_{\kappa+n_0-m_0} = T_{z^{m_0-n_0-\kappa}} Q_{\kappa+n_0-m_0} = T_{z^{m-n}} Q_{\kappa+n_0-m_0}.$$

Therefore, we have

$$\begin{aligned} \text{Ran}(T_{\omega}^*) &= T_{1/(q_+)^{\sharp}} T_{z^{m-n}} T_{(s_+)^{\sharp}} Q_{\kappa+n_0-m_0}(s_0)^{\sharp} H^{p'} \\ &= T_{z^{m-n}(s_+)^{\sharp}/(q_+)^{\sharp}} Q_{\kappa+n_0-m_0}(s_0)^{\sharp} H^{p'}, \end{aligned}$$

again using that $1/(q_+)^{\sharp}$ is anti-analytic and $(s_+)^{\sharp}$ is analytic. This gives the general formula for $\text{Ran}(T_{\omega}^*)$. In case $\kappa + n_0 - m_0 \leq 0$, we have $Q_{\kappa+n_0-m_0} = I$ and $T_{(s_+)^{\sharp}} Q_{\kappa+n_0-m_0}(s_0)^{\sharp} = (s_+ s_0)^{\sharp}$, as claimed.

5. SYMMETRIC OPERATORS AND SELFADJOINT EXTENSIONS

For $\omega \in \text{Rat}$, the second adjoint T_{ω}^{**} is well-defined and $T_{\omega}^{**} = T_{\omega}$, since T_{ω} is a closed, densely defined operator on a reflexive Banach space [8, Theorem III.5.24]. Now consider $\omega \in \text{Rat}(\mathbb{T})$ and $p = 2$. From Theorem 1.1 it is obvious that $T_{\omega} \neq T_{\omega}^*$, except in the degenerate case where q is constant, since $\text{Dom}(T_{\omega}) = qH^2 + \mathcal{P}_{\deg(q)-1}$ contains all polynomials while $\text{Dom}(T_{\omega}^*) = q^{\sharp}H^2$ only contains the polynomials that contain q^{\sharp} as a factor. Consequently, T_{ω} cannot be selfadjoint. In this section we consider the question when T_{ω}^* is symmetric, and, if this is the case, when does T_{ω}^* have a selfadjoint extension L . The first topic is addressed in the following theorem.

Theorem 5.1. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime. Set $n = \deg(s)$ and $m = \deg(q)$. Then the following are equivalent.*

- (1) T_{ω}^* is symmetric;
- (2) $\omega(\mathbb{T}) \subset \mathbb{R}$;
- (3) $\omega(z) = \tilde{\omega}(-i\frac{z+1}{z-1})$ with $\tilde{\omega}$ a real rational function with poles only on \mathbb{R} ;
- (4) the essential spectrum $\sigma_{\text{ess}}(T_{\omega})$ of T_{ω} is contained in \mathbb{R} ;
- (5) ω is proper, $s = z^{m-n}\tilde{s}$ with \tilde{s} self-inversive and $q_0\overline{s_n} = \overline{q_m}s_{m-n}$ holds, where $s(z) = \sum_{k=0}^n s_k z^k$ and $q(z) = \sum_{k=0}^m q_k z^k$.

Moreover, if T_{ω}^* is symmetric, then $T_{\omega}^* \subset T_{\omega}$.

Proof. We first prove the equivalence of (1) and (2), and that (1) implies $T_{\omega}^* \subset T_{\omega}$. Assume (2). Then, for $z \in \mathbb{T}$, not a root of q , we have $\omega^*(z) = \overline{\omega(z)} = \omega(z)$. Hence $\omega^* = \omega$. Since q has only roots on \mathbb{T} , we have $q = \gamma q^{\sharp}$ for a unimodular constant γ . Hence $qH^2 = q^{\sharp}H^2$. This shows $T_{\omega}^* = T_{\omega^*}|_{q^{\sharp}H^2} = T_{\omega}|_{qH^2} \subset T_{\omega}$. Since $(T_{\omega}^*)^* = T_{\omega}$, it follows that T_{ω}^* is symmetric and $T_{\omega}^* \subset T_{\omega}$. Conversely, assume (1). Then we still have $qH^2 = q^{\sharp}H^2$ and $T_{\omega}^* \subset (T_{\omega}^*)^* = T_{\omega}$. Hence $T_{\omega}^* = T_{\omega}|_{qH^2}$. In particular, we have $\omega^*q = T_{\omega^*}q = T_{\omega}^*q = T_{\omega}q = \omega q$. This implies $\omega = \omega^*$. Hence $\omega(z) = \overline{\omega(z)}$ for $z \in \mathbb{T}$, not a root of q . Thus $\omega(\mathbb{T}) \subset \mathbb{R}$.

That (2) and (3) are equivalent follows simply because in (3) ω is the composition of $\tilde{\omega}$ and the inverse Cayley transform, which maps the circle \mathbb{T} bijectively onto \mathbb{R} . The fact that $\tilde{\omega}$ is real rational, i.e., $\tilde{\omega} = \tilde{s}/\tilde{q}$ with \tilde{s} and \tilde{q} real polynomials, is equivalent to $\tilde{\omega}(\mathbb{R}) := \{\tilde{\omega}(t) : t \in \mathbb{R}, \tilde{q}(t) \neq 0\} \subset \mathbb{R}$. Also, the equivalence of (2) and (4) is a direct consequence of the fact that $\sigma_{\text{ess}}(T_{\omega}) = \omega(\mathbb{T})$, by [6, Theorem 1.1].

Finally, we prove (2) \Leftrightarrow (5). Since $q = \gamma q^\sharp$, we have

$$\omega^* = z^{m-n} \frac{s^\sharp}{q^\sharp} = z^{m-n} \gamma \frac{s^\sharp}{q}.$$

Thus, we have $\omega = \omega^*$ if and only if $z^{m-n} \gamma s^\sharp = s$. Hence (2) is equivalent to $z^{m-n} \gamma s^\sharp = s$. Now assume (2). Since $\deg(s^\sharp) \leq \deg(s)$, the identity $z^{m-n} \gamma s^\sharp = s$ can only occur if $m \geq n$, i.e., if ω is proper. The identity also shows that $s = z^{m-n} \tilde{s}$ for $\tilde{s} = \gamma s^\sharp$. On the other hand, $s^\sharp = (z^{m-n} \tilde{s})^\sharp = \tilde{s}^\sharp$. Thus $\tilde{s} = \gamma s^\sharp = \gamma \tilde{s}^\sharp$, which shows \tilde{s} is self-inversive, with constant γ . Note that $\gamma = q_0/\overline{q_m}$. Also, we have $s_0 = \dots = s_{m-n-1} = 0$ and $\tilde{s}(z) = \sum_{k=0}^{2n-m} s_{m-n+k} z^k$. Since \tilde{s} is self-inversive, $\tilde{s} = \delta \tilde{s}^\sharp$ with $\delta = s_{m-n}/\overline{s_n}$. But also $\delta = \gamma$, so $s_{m-n}/\overline{s_n} = q_0/\overline{q_m}$. Thus $q_0 \overline{s_n} = \overline{q_m} s_{m-n}$. Hence (5) holds. Conversely, assume (5). Reversing the above argument, it follows that $q_0 \overline{s_n} = \overline{q_m} s_{m-n}$ implies $\tilde{s} = \delta \tilde{s}^\sharp$ with $\delta = \gamma$. Thus $\gamma s^\sharp = \gamma \tilde{s}^\sharp = \tilde{s}$. This implies $s = z^{m-n} \tilde{s} = z^{m-n} \gamma s^\sharp$, and hence (2). \square

Corollary 5.2. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime. Assume T_ω^* is symmetric. Then $\deg(s) \leq \deg(q) \leq 2 \deg(s)$.*

Proof. By Theorem 5.1 condition (5) holds with $m = \deg(q)$ and $n = \deg(s)$. Since \tilde{s} is self-inversive, we have $\tilde{s}(0) \neq 0$. Consequently, 0 would be a non-removable singularity of $s = z^{m-n} \tilde{s}$ in case $m < n$, which gives a contradiction. Hence $m \geq n$. Furthermore, comparing the degrees on both sides of $s = z^{m-n} \tilde{s}$ yields, $n = m - n + \deg(\tilde{s}) \geq m - n$. Hence $m \leq 2n$. \square

When T_ω^* is symmetric, it need not be the case that T_ω^* has a selfadjoint extension. In Proposition 5.4 below we characterize when T_ω^* does have a selfadjoint extension. However, we first give a concrete example that shows this does not always happen.

Example 5.3. In [7] Helson considered the functions $\omega_k(z) = \left(-i \frac{z+1}{z-1}\right)^k$ for $k \in \mathbb{N}$. For all k we have $\omega_k(\mathbb{T}) \subset \mathbb{R}$, see Theorem 5.1 (3) above, hence $T_{\omega_k}^*$ is symmetric by Theorem 5.1. In fact, for k even $\omega_k(\mathbb{T}) = \mathbb{R}_+$, while for k odd we have $\omega_k(\mathbb{T}) = \mathbb{R}$. We show that $T_{\omega_k}^*$ does not have a selfadjoint extension for $k = 1$. In Example 5.8 we return to this example for general k .

For $k = 1$ we have $\omega(z) = \omega_1(z) = -i \frac{z+1}{z-1}$. Hence $\text{Dom}(T_\omega) = (z-1)H^2 + \mathbb{C}$ and $\text{Dom}(T_\omega^*) = (z-1)H^2$. Suppose T_ω^* has a selfadjoint extension L . Then $L = L^*$ and thus $T_\omega^* \subset L = L^* \subset T_\omega^{**} = T_\omega$. Since T_ω is not selfadjoint, the inclusions are strict. Hence $\text{Dom}(T_\omega^*) \subset \text{Dom}(L) \subset \text{Dom}(T_\omega)$, with strict inclusions. However, the complement of $\text{Dom}(T_\omega^*)$ in $\text{Dom}(T_\omega)$ is one-dimensional, hence not both inclusions can be strict. Thus T_ω does not admit a selfadjoint extension.

Proposition 5.4. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Then T_ω^* admits a selfadjoint extension if and only if the number of roots of $s - iq$ and $s + iq$ in \mathbb{D} , counting multiplicities, coincide.*

Proof. The operator T_ω^* is an adjoint, and hence closed, and by assumption symmetric. Following definition X.2.12 from [3] we define the deficiency subspaces of T_ω^* as the spaces

$$\mathcal{L}_+ = \text{Ker}(T_\omega^{**} - i) = (\text{Ran}(T_\omega^* + i))^\perp, \quad \mathcal{L}_- = \text{Ker}(T_\omega^{**} + i) = (\text{Ran}(T_\omega^* - i))^\perp,$$

and the deficiency indices as the integers $n_\pm = \dim \mathcal{L}_\pm$. Since $T_\omega^{**} = T_\omega$, we have

$$n_+ = \dim \text{Ker}(T_\omega - i) \quad \text{and} \quad n_- = \dim \text{Ker}(T_\omega + i).$$

Also, we have $T_\omega \pm i = T_{\omega \pm i}$. By item (b) of Theorem X.2.20 in [3], T_ω has a selfadjoint extension if and only if $n_+ = n_-$. Note that $\omega \pm i = (s \pm iq)/q$. We now apply Corollary 4.2 from [5] to $T_{\omega \pm i}$, to obtain that n_\pm is equal to the maximum of 0 and the difference of m and the number of roots of $s \pm iq$ in $\overline{\mathbb{D}}$, counting multiplicities. However, since T_ω^* is symmetric, ω is proper so the number of roots cannot exceed m . Note also that $\omega(\mathbb{T}) \subset \mathbb{R}$, so $s \pm iq$ cannot have roots on \mathbb{T} . It thus follows that T_ω^* has a selfadjoint extension if and only if the number of roots in \mathbb{D} of $s - iq$ and $s + iq$, counting multiplicities, coincide, as claimed. \square

Since T_ω^* is never selfadjoint for $\omega \in \text{Rat}(\mathbb{T})$ having at least one pole on \mathbb{T} , the formulas for n_\pm in the above proof along with item (a) of Theorem X.2.20 in [3] directly give the following corollary.

Corollary 5.5. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Then $s + iq$ or $s - iq$ must have a root in \mathbb{D} .*

Proposition 5.4 can be rephrased in terms of the index of the operators $T_{\omega \pm i}$.

Proposition 5.6. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Then $T_{\omega+i}$ and $T_{\omega-i}$ are both Fredholm and T_ω^* admits a selfadjoint extension if and only if the Fredholm indices of $T_{\omega+i}$ and $T_{\omega-i}$ coincide.*

Proof. This follows directly from Proposition 5.4 and Theorem 1.1 of [5] applied to $\omega + i$ and $\omega - i$, using that $\omega \pm i = (s \pm iq)/q$. \square

Corollary 5.7. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Assume $\omega(\mathbb{T}) \neq \mathbb{R}$. Then T_ω^* admits a selfadjoint extension.*

Proof. The Fredholm index of $T_{\omega-\lambda}$ is constant with respect to $\lambda \in \mathbb{C}$ on the connected components of \mathbb{C} separated by the essential spectrum of T_ω , which is equal to $\omega(\mathbb{T})$; see [6, Theorem 1.1]. Hence if $\omega(\mathbb{T}) \neq \mathbb{R}$, but $\omega(\mathbb{T}) \subset \mathbb{R}$ since T_ω^* is symmetric, then i and $-i$ are in the same connected component and thus $T_{\omega+i}$ and $T_{\omega-i}$ have the same index. The conclusion now follows from Proposition 5.6. \square

Example 5.8. We return to the functions $\omega_k(z) = \left(-i \frac{z+1}{z-1}\right)^k$ considered in Example 5.3. Since $\omega_k(\mathbb{T}) = \mathbb{R}_+$ for k even, we obtain directly from Corollary 5.7 that $T_{\omega_k}^*$ admits a selfadjoint extension in case k is even.

For odd values of k we have $\omega_k(\mathbb{T}) = \mathbb{R}$, and thus no conclusion can be drawn from Corollary 5.7. To deal with the odd case we resort to Proposition 5.4. Take $s(z) = (-i)^k(z+1)^k$ and $q = (z-1)^k$ and write k as $k = 2l + 1$. The polynomials $s \pm iq$ are given by

$$\begin{aligned} s(z) \pm iq(z) &= i \left((-1)^{l+1} (z+1)^{2l+1} \pm (z-1)^{2l+1} \right) \\ &= i \left((-1)^{l+1} \sum_{j=0}^{2l+1} \binom{2l+1}{j} z^j \pm \sum_{j=0}^{2l+1} \binom{2l+1}{j} z^j (-1)^{2l+1-j} \right) \\ &= i \sum_{j=0}^{2l+1} \binom{2l+1}{j} z^j \left((-1)^{l+1} \pm (-1)^{2l+1-j} \right) \\ &= i \sum_{j=0}^{2l+1} \binom{2l+1}{j} z^j \left((-1)^{l+1} \pm (-1)^{j-1} \right). \end{aligned}$$

For odd values of l one obtains:

$$\begin{aligned} s(z) - iq(z) &= -2i \left(\binom{2l+1}{0} + \cdots + \binom{2l+1}{2l-2} z^{2l-2} + \binom{2l+1}{2l} z^{2l} \right), \\ s(z) + iq(z) &= 2i \left(\binom{2l+1}{1} z + \cdots + \binom{2l+1}{2l-1} z^{2l-1} + \binom{2l+1}{2l+1} z^{2l+1} \right) \\ &= 2iz \left(\binom{2l+1}{2l} + \cdots + \binom{2l+1}{2} z^{2-2} + \binom{2l+1}{0} z^{2l} \right) \end{aligned}$$

Observe that $s + iq$ is of the form $izp_+(z^2)$ where p_+ is a real polynomial of degree $2l$ and that $s - iq$ is of the form $ip_-(z^2)$ where p_- is a real polynomial of degree $2l$. Because p_+ and p_- are real polynomials and the fact that z^2 is the variable rather than z itself, the nonzero roots of $zp_+(z^2)$ come either in pairs (z and $-z$) for real nonzero roots or in quadruples ($z, \bar{z}, -z, -\bar{z}$) for nonreal roots, while zero appears as a simple root. Similarly, the roots of $p_-(z^2)$ come in pairs (z and $-z$) or quadruples ($z, \bar{z}, -z, -\bar{z}$) and there is no root at zero. Hence $s + iq$ has an odd number of roots inside the unit disc, and $s - iq$ has an even number of roots inside the unit disc, so that the indices n_+ and n_- can never coincide. One further observes that $p_- = p_+^\sharp$. In a similar way, for even values of l the polynomial $s + iq$ will have an even number of roots inside the unit disc and $s - iq$ will have an odd number of roots inside the unit disc. Hence, in all cases where k is odd, T_ω^* does not have a selfadjoint extension.

We now present a proposition that rephrases the criteria of Proposition 5.4 in terms of the roots of $s + iq$ (or $s - iq$) only. The observation that $T_{\omega_k}^*$ in Example 5.8 has no selfadjoint extension follows as a special case. In general, T_ω^* cannot have a selfadjoint extension whenever $\deg(q)$ is odd for any $\omega \in \text{Rat}(\mathbb{T})$.

Proposition 5.9. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Set $l_\pm = m - \deg(s \pm iq)$ and define*

$$k_{\pm,1} = \# \left\{ \begin{array}{l} \text{zeroes of } \omega \pm i \text{ inside } \mathbb{T} \\ \text{multi. taken into account} \end{array} \right\}, \quad k_{\pm,2} = \# \left\{ \begin{array}{l} \text{zeroes of } \omega \pm i \text{ outside } \mathbb{T} \\ \text{multi. taken into account} \end{array} \right\},$$

Then

$$T_\omega^* \text{ has a selfadjoint extension} \iff l_+ + k_{+,2} = k_{+,1} \iff l_- + k_{-,2} = k_{-,1}.$$

In particular, if T_ω^* has a selfadjoint extension, then $\deg(q)$ must be even.

The basis for the proof of Proposition 5.9 lies in the following lemma, which clarifies the relation between $s + iq$ and $s - iq$ under the assumption that T_ω^* is symmetric.

Lemma 5.10. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ coprime, be such that T_ω^* is symmetric. Set $l_\pm = \deg(q) - \deg(s \pm iq)$ and let γ be the unimodular constant such that $q = \gamma q^\sharp$. Then*

$$s \pm iq = \gamma z^{l_\mp} (s \mp iq)^\sharp. \quad (5.1)$$

Moreover, we have $l_\pm = 0$ if and only if $\omega(0) = \pm i$. In particular, only one of l_+ and l_- can be nonzero.

Proof. Since T_ω^* is symmetric, by assumption, ω has the properties listed in Theorem 5.1. In particular, ω is proper, $m := \deg(q) \geq \deg(s) =: n$, and $s = z^{m-n} \bar{s}$ with

\tilde{s} self-inversive and the unimodular constants that establish the self-inversiveness of \tilde{s} and q coincide (equivalently, $q_0 \overline{s_n} = \overline{q_m} s_{m-n}$).

Note that $\deg(s \pm iq) \neq m$ occurs precisely when $\deg(s) = \deg(q)$ and the leading coefficients s_m and q_m of s and q , respectively, satisfy $s_m \pm iq_m = 0$, i.e., $s_m/q_m = \mp i$. Since $m = n$, the identity $q_0 \overline{s_n} = \overline{q_m} s_{m-n}$ shows $\omega(0) = s_0/q_0 = \overline{s_m}/\overline{q_m}$. Hence $\deg(s \pm iq) \neq m$ holds if and only if $\omega(0) = \mp i = \pm i$, as claimed.

We first prove (5.1) for the case $\omega(0) = 0$. So assume $\omega(0) = 0$, or equivalently, $s(0) = 0$. In this case $l_+ = l_- = 0$. Since $s = z^{m-n} \tilde{s}$ and $\tilde{s}(0) \neq 0$ (because \tilde{s} is self-inversive), we have $m > n$. Also note that $m - n$ is equal to the multiplicity of 0 as a root of s . We now employ Lemma 2.2, using that $\deg(s + iq) = m = \deg(iq)$, to obtain

$$\gamma(s \mp iq)^\sharp = z^{\deg(s+iq)-\deg(s)} \gamma s^\sharp \mp (-i) \gamma q^\sharp = z^{m-n} \gamma \tilde{s}^\sharp \pm iq = z^{m-n} \tilde{s} \pm iq = s \pm iq.$$

Hence (5.1) holds.

Now assume $\omega(0) \neq 0$, i.e., $s(0) \neq 0$. In that case $s = \tilde{s}$. Hence s is self-inversive with the same constant γ that establishes the self-inversiveness of q . This also yields $m = n$. Since s and q are self-inversive with the same constant γ , we have

$$\overline{s_{m-k}} q_k = \overline{q_{m-k}} s_{m-k} \gamma = \overline{q_{m-k}} s_k \quad \text{for } k = 0, \dots, m.$$

Hence for all k we have

$$\overline{s_{m-k}}(s_k + iq_k) = s_k(\overline{s_{m-k}} + i\overline{q_{m-k}}) \quad \text{and} \quad \overline{q_{m-k}}(s_k + iq_k) = q_k(\overline{s_{m-k}} + i\overline{q_{m-k}}).$$

In case $s_{m-k} = 0$ and $q_{m-k} = 0$, also $s_k = 0$ and $q_k = 0$, since $s_k = \gamma \overline{s_{m-k}}$ and $q_k = \gamma \overline{q_{m-k}}$, and thus $s_k + iq_k = 0 = \gamma(\overline{s_{m-k}} + i\overline{q_{m-k}})$. If either $s_{m-k} \neq 0$ or $q_{m-k} \neq 0$, divide the first identity by $\overline{s_{m-k}}$ or the second identity by $\overline{q_{m-k}}$ to arrive at $s_k + iq_k = \gamma(\overline{s_{m-k}} + i\overline{q_{m-k}})$. Hence

$$s_k + iq_k = \gamma(\overline{s_{m-k}} + i\overline{q_{m-k}}) \quad \text{for } k = 0, \dots, m. \quad (5.2)$$

In particular, $s_k + iq_k = 0$ if and only if $\overline{s_{m-k}} + i\overline{q_{m-k}} = 0$. It follows that 0 is a root of $s \pm iq$ with multiplicity l_\mp . Comparing coefficients, it follows that the identities in (5.1) correspond to the identities in (5.2). Hence (5.1) holds. \square

Proof of Proposition 5.9. Since T_ω^* is assumed to be symmetric, (5.1) holds. Together with the fact that the \sharp operator reflects roots over \mathbb{T} , this implies that the number of roots of $s \pm iq$ inside \mathbb{T} are equal to l_\pm plus the number of roots of $s \mp iq$ outside \mathbb{T} , counting multiplicities. In other words, we have

$$k_{+,1} = l_- + k_{-,2} \quad \text{and} \quad k_{-,1} = l_+ + k_{+,2}. \quad (5.3)$$

By Proposition 5.6, T_ω^* has a selfadjoint extension if and only if $s + iq$ and $s - iq$ have an equal number of roots inside \mathbb{T} , again counting multiplicities, equivalently, $k_{+,1} = k_{-,1}$. Given (5.3), it follows that $k_{+,1} = k_{-,1}$ is equivalent to $k_{+,1} = l_+ + k_{+,2}$, and likewise to $k_{-,1} = l_- + k_{-,2}$. This proves the two criteria for T_ω^* to have a selfadjoint extension.

By Lemma 5.10, either $l_+ = 0$ or $l_- = 0$. Say $l_+ = 0$. Since $s + iq$ cannot have roots on \mathbb{T} , we have $\deg(q) = \deg(s + iq) = k_{+,1} + k_{+,2}$. If T_ω^* admits a selfadjoint extension, then we have $k_{+,1} = l_+ + k_{+,2} = k_{+,2}$. Hence $\deg(q) = 2k_{+,1}$ is even. For $l_- = 0$ the arguments goes similarly. \square

Combining the fact that T_ω^* cannot have a selfadjoint extension in case $\omega = s/q \in \text{Rat}(\mathbb{T})$, s, q co-prime, and $\deg(q)$ odd with Corollary 5.7 immediately yields the following result.

Corollary 5.11. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ co-prime, be such that T_ω^* is symmetric and $\deg(q)$ is odd. Then $\omega(\mathbb{T}) = \mathbb{R}$.*

The next example shows that also with $\deg(q)$ even it can occur that T_ω^* does not admit a selfadjoint extension.

Example 5.12. Let $\omega = s/q$ with

$$s(z) = i(1 + az + z^2), \text{ for some } 0 \neq a \in \mathbb{R}, \text{ and } q(z) = 1 - z^2.$$

Then $m = n$ and

$$s^\sharp = -s, \quad q^\sharp = -q.$$

So T_ω^* is symmetric by Theorem 5.1 (5). Also, we have

$$(s + iq)(z) = i(2 + az) \quad \text{and} \quad (s - iq)(z) = iz(a + 2z).$$

Hence the number of roots of $s - iq$ inside \mathbb{D} is 1 if $|a| \geq 2$ and 2 if $0 \neq |a| < 2$, while the number of roots of $s + iq$ inside \mathbb{D} is 1 if $|a| > 2$ and 0 if $0 \neq |a| \leq 2$. Thus T_ω^* admits a selfadjoint extension if and only if $|a| > 2$.

6. COMPARISON WITH THE UNBOUNDED TOEPLITZ OPERATOR DEFINED BY SARASON

The Smirnov class N^+ consists of quotients $\frac{b}{a}$ with a and b H^∞ -functions such that the denominator a is an outer function. The function $\varphi = \frac{b}{a} \in N^+$ is said to be in *canonical form* if $a(0) > 0$ and $|a|^2 + |b|^2 = 1$ on \mathbb{T} . By Proposition 3.1 of [12], every function $\varphi \in N^+$ can be uniquely written in canonical form.

In [12], Sarason investigated an unbounded Toeplitz operator T_φ^{Sa} with symbol φ in N^+ , which is defined by

$$\text{Dom}(T_\varphi^{\text{Sa}}) = \{f \in H^2 : \varphi f \in H^2\}, \quad T_\varphi^{\text{Sa}} f = \varphi f \quad (f \in \text{Dom}(T_\varphi^{\text{Sa}})).$$

More generally, T_φ^{Sa} can be defined in this way for any holomorphic function φ on \mathbb{D} , but for T_φ^{Sa} to be densely defined, φ must be in N^+ ; see [12, Lemma 5.2].

Let $\varphi = \frac{b}{a} \in N^+$ be the canonical representation of φ . Then it is shown in Proposition 5.3 of [12] that $\text{Dom}(T_\varphi^{\text{Sa}}) = aH^2$. The adjoint of the operator T_φ^{Sa} is motivated by the action of the conjugate transpose of the matrix representation of T_φ^{Sa} , which is lower triangular. The domain of the adjoint operator is shown to contain the space $H(\overline{\mathbb{D}})$ of functions that are analytic on some neighborhood of the closed unit disc $\overline{\mathbb{D}}$, and the adjoint is equal to the closure of the operator on $H(\overline{\mathbb{D}})$; see [12, Lemmas 6.1 and 6.4].

Let $\omega = s/q \in \text{Rat}(\mathbb{T})$ with $s, q \in \mathcal{P}$ co-prime. Set $n = \deg(s)$ and $m = \deg(q)$. Assume ω is proper, i.e., $n \leq m$. Then $\omega^*(z) = z^{m-n}s^\sharp/q^\sharp \in \text{Rat}(\mathbb{T})$. Since q^\sharp has zeroes only on \mathbb{T} it is outer and thus $\omega^* \in N^+$. While in general T_ω and T_ω^{Sa} are different, the following proposition shows that T_ω coincides with $T_{\omega^*}^{\text{Sa}}$, and hence $T_\omega = T_\omega^{**} = T_{\omega^*}^{\text{Sa}}$. Without the properness assumption, ω^* is not in N^+ , because ω^* has a pole at 0, and hence $T_{\omega^*}^{\text{Sa}}$ is not defined.

Proposition 6.1. *Let $\tilde{\omega} = \tilde{s}/\tilde{q} \in \text{Rat}(\mathbb{T})$ with $\tilde{s}, \tilde{q} \in \mathcal{P}$ co-prime. Then $\text{Dom}(T_{\tilde{\omega}}^{\text{Sa}}) = \tilde{q}H^2$ and $T_{\tilde{\omega}}^{\text{Sa}} = T_{\tilde{\omega}}|_{\tilde{q}H^2}$. In particular, if $\omega \in \text{Rat}(\mathbb{T})$ is proper, then $T_\omega^* = T_{\omega^*}^{\text{Sa}}$.*

Proof. We first show $\text{Dom}(T_{\tilde{\omega}}^{\text{Sa}}) = \tilde{q}H^2$. Let $\tilde{\omega} = a/b$ be the canonical form of $\tilde{\omega}$. As noted above, $\text{Dom}(T_{\tilde{\omega}}^{\text{Sa}}) = aH^2$. By the Fejér-Riesz Theorem there is a polynomial r such that on \mathbb{T} we have $|r|^2 = |\tilde{s}|^2 + |\tilde{q}|^2$, r has no roots in

\mathbb{D} and $\arg(r(0)) = \arg(\tilde{q}(0))$. The latter is possible since $\tilde{q}(0) \neq 0$ and implies $\tilde{q}(0)/r(0) > 0$. Note that r also has no roots on \mathbb{T} , since \tilde{s} and \tilde{q} are co-prime. It follows that \tilde{q}/r and \tilde{s}/r are both H^∞ -functions, \tilde{q}/r is outer and $\tilde{q}(0)/r(0) > 0$. Hence $a = \tilde{q}/r$ and $b = \tilde{s}/r$, by the uniqueness of the canonical form. Also, since all the roots of r are outside \mathbb{T} , $r^{-1}H^2 = H^2$, so that $aH^2 = \tilde{q}H^2$.

Now let $f \in \text{Dom}(T_\omega^{\text{Sa}})$, say $f = \tilde{q}h$ with $h \in H^2$. Then $T_\omega^{\text{Sa}}f = \tilde{\omega}f = \tilde{s}h$. On the other hand, the fact that $\tilde{\omega}f = \tilde{s}h$ and $\tilde{s}h \in H^2$ shows $T_\omega f = \mathbb{P}\tilde{s}h = \tilde{s}h$. Hence $T_\omega^{\text{Sa}} = T_\omega|_{\tilde{q}H^2}$. \square

Next we employ some of the ideas from [12] to derive the following result. Recall that for a Hilbert space operator $T : \text{Dom}(T) \rightarrow \mathcal{H}$ a linear submanifold $\mathcal{D} \subset \text{Dom}(T)$ is called a *core* in case the graph $G(T|_{\mathcal{D}})$ of $T|_{\mathcal{D}}$ is dense in the graph $G(T)$ of T ; cf., page 166 in [8].

Theorem 6.2. *Let $\omega \in \text{Rat}(\mathbb{T})$. Then $H(\overline{\mathbb{D}})$ is contained in $\text{Dom}(T_\omega)$. If ω is proper, then $H(\overline{\mathbb{D}})$ is a core of T_ω .*

Proof of $H(\overline{\mathbb{D}}) \subset \text{Dom}(T_\omega)$. Write $\omega = \frac{s}{q} \in \text{Rat}_0(\mathbb{T})$ with $s, q \in \mathcal{P}$ coprime. Let $f \in H(\overline{\mathbb{D}})$. Then there exists a $R > 1$ such that f is still analytic on an open neighborhood of the closed disc with radius R . Set $\tilde{f}(z) = f(Rz)$, $\tilde{q}(z) = q(Rz)$ and $\tilde{s}(z) = s(Rz)$. Then $\tilde{f} \in H^2$ and \tilde{q} is a polynomial with no roots on \mathbb{T} and $\deg(q) = \deg(\tilde{q})$. By Theorem 3.1 in [5], $H^2 = \tilde{q}H^2 + \mathcal{P}_{\deg(q)-1}$. Thus $\tilde{s}\tilde{f} = \tilde{q}\tilde{h} + \tilde{r}$ for some $\tilde{h} \in H^2$ and $\tilde{r} \in \mathcal{P}$ with $\deg(\tilde{r}) < \deg(q)$. Now set $r(z) = \tilde{r}(z/R)$ and $h(z) = \tilde{h}(z/R)$. Then $r \in \mathcal{P}$ with $\deg(r) = \deg(\tilde{r}) < \deg(q)$ and $h \in H^2$, even $h \in H(\overline{\mathbb{D}})$. Also, we have $sf = qh + r$. Thus $f \in \text{Dom}(T_\omega)$. \square

Before proving the second claim of Theorem 6.2 it is useful to consider the value of T_ω when applied to the evaluation functional or reproducing kernel element $k_\lambda(z) = (1 - \bar{\lambda}z)^{-1}$, where $\lambda \in \mathbb{D}$. Note that $k_\lambda \in H(\overline{\mathbb{D}})$, hence $k_\lambda \in H^2$, and k_λ has the reproducing kernel property for H^2 :

$$\text{span}\{k_\lambda : \lambda \in \mathbb{D}\} \text{ dense in } H^2 \quad \text{and} \quad \langle h, k_\lambda \rangle = h(\lambda) \quad (h \in H^2, \lambda \in \mathbb{D}).$$

See [9] for a recent account of the theory of reproducing kernel Hilbert spaces and further references.

Lemma 6.3. *Let $\omega = s/q \in \text{Rat}(\mathbb{T})$, with $s, q \in \mathcal{P}$ co-prime, be proper. Then*

$$T_\omega k_\lambda = \overline{\omega^*(\lambda)} k_\lambda \quad (\lambda \in \mathbb{D}).$$

Proof. Suppose $g = T_\omega k_\lambda$ then $s(z)(1 - \bar{\lambda}z)^{-1} = q(z)g(z) + r(z)$, where $r \in \mathcal{P}_{m-1}$. Here $m = \deg(q)$. Hence $(1 - \bar{\lambda}z)g = (s + (1 - \bar{\lambda}z)r)/q$ is in $\text{Rat}(\mathbb{T})$ as well as in H^2 . This can only occur if $(1 - \bar{\lambda}z)g$ is a polynomial, i.e., $g = k_\lambda \tilde{r}$ for some $\tilde{r} \in \mathcal{P}$. Thus $s + (1 - \bar{\lambda}z)r = q\tilde{r}$. Since ω is proper, the degree of the left hand side is at most m . But then \tilde{r} is constant, say with value \tilde{c} . This shows $T_\omega k_\lambda = \tilde{c}k_\lambda$.

To determine \tilde{c} we evaluate the identity $s + (1 - \bar{\lambda}z)r = q\tilde{c}$ at $1/\bar{\lambda}$. This gives $s(1/\bar{\lambda}) = q(1/\bar{\lambda})\tilde{c}$. Note that

$$s^\sharp(\lambda) = \lambda^n \overline{s(1/\bar{\lambda})} \quad \text{and} \quad q^\sharp(\lambda) = \lambda^m \overline{q(1/\bar{\lambda})},$$

where $n = \deg(s)$. Hence

$$s(1/\bar{\lambda}) = \overline{\lambda^{-n} s^\sharp(\lambda)} \quad \text{and} \quad q(1/\bar{\lambda}) = \overline{\lambda^{-m} q^\sharp(\lambda)}.$$

This gives

$$\tilde{c} = \frac{\overline{\lambda^{-n} s^\#(\lambda)}}{\overline{\lambda^{-m} q^\#(\lambda)}} = \overline{\left(\frac{\lambda^{m-n} s^\#(\lambda)}{q^\#(\lambda)} \right)} = \overline{\omega^*(\lambda)}. \quad \square$$

Proof of Theorem 6.2. It remains to prove that $H(\overline{\mathbb{D}})$ is a core for T_ω in case ω is proper. So, assume ω is proper. We need to show that the graph of $T_\omega|_{H(\overline{\mathbb{D}})}$ is dense in the graph of T_ω . In other words, let $f, g \in H^2$ with (f, g) perpendicular to $G(T_\omega|_{H(\overline{\mathbb{D}})})$, then we need to show (f, g) is perpendicular to $G(T_\omega)$. Since $k_\lambda \in H(\overline{\mathbb{D}})$, for $\lambda \in \mathbb{D}$, we have

$$0 = \langle (f, g), (k_\lambda, T_\omega k_\lambda) \rangle = \langle f, k_\lambda \rangle + \langle g, \overline{\omega^*(\lambda)} k_\lambda \rangle = f(\lambda) + \omega^*(\lambda) g(\lambda) \quad (\lambda \in \mathbb{D}).$$

Hence $\omega^* g = -f$. In particular, $\omega^* g \in H^2$. Thus $g \in \text{Dom}(T_\omega^{\text{Sa}}) = \text{Dom}(T_\omega^*)$ and $T_\omega^* g = -f$, by Proposition 6.1. For any $h \in \text{Dom}(T_\omega)$ we have

$$\langle (f, g), (h, T_\omega h) \rangle = \langle (-T_\omega^* g, g), (h, T_\omega h) \rangle = -\langle T_\omega^* g, h \rangle + \langle g, T_\omega h \rangle = 0.$$

This proves our claim. \square

In Section 8 of [12], Sarason introduced the class of closed, densely defined operators T on H^2 which satisfy

- (1) $T_z \text{Dom}(T) \subset \text{Dom}(T)$;
- (2) $T_z^* T T_z = T$;
- (3) $f \in \text{Dom}(T)$, $f(0) = 0 \Rightarrow T_z^* f \in \text{Dom}(T)$.

This class of operators was further studied by Rosenfeld in [11], see also [10], in which he referred to such operators as Sarason-Toeplitz operators. The operators T_φ^{Sa} , for $\varphi \in N^+$, are Sarason-Toeplitz operators, and the class of operators is closed under taking adjoints, by Proposition 2.1 in [11]. Hence, by Proposition 6.1, T_ω is a Sarason-Toeplitz operator whenever $\omega \in \text{Rat}(\mathbb{T})$ is proper. We show that in fact T_ω is a Sarason-Toeplitz operator for any $\omega \in \text{Rat}$.

Proposition 6.4. *Let $\omega \in \text{Rat}$. Then T_ω on H^2 is a Sarason-Toeplitz operator.*

Proof. First consider $\omega \in \text{Rat}(\mathbb{T})$. That T_ω satisfies (1) and (2) was proved in [5, Lemma 2.3]. We claim that $T_z^* \text{Dom}(T_\omega) \subset \text{Dom}(T_\omega)$. Write $\omega = s/q$ with $s, q \in \mathcal{P}$ co-prime. Then $\text{Dom}(T_\omega) = qH^2 + \mathcal{P}_{\deg(q)-1}$. Let $f = qh + r \in \text{Dom}(T_\omega)$ with $h \in H^2$ and $r \in \mathcal{P}$, $\deg(r) < \deg(q)$. Then $T_z^* f = qT_z^* h + h(0)T_z^* q + T_z^* r$, which is in $qH^2 + \mathcal{P}_{\deg(q)-1} = \text{Dom}(T_\omega)$. Hence T_ω is a Sarason-Toeplitz operator in case $\omega \in \text{Rat}(\mathbb{T})$.

Now take $\omega \in \text{Rat}$ arbitrarily. By Lemma 5.1 in [5], see also Section 4 above, $\omega = \omega_- z^\kappa \omega_0 \omega_+$ with $\kappa \in \mathbb{Z}$, and ω_- , ω_0 and ω_+ in Rat with zeroes and poles only inside, on or outside \mathbb{T} , respectively. In particular, $\omega_0 \in \text{Rat}(\mathbb{T})$, ω_- and ω_-^{-1} are both anti-analytic, and ω_+ and ω_+^{-1} are both analytic. Also, $T_\omega = T_{\omega_-} T_{z^\kappa \omega_0} T_{\omega_+}$. Note that $z^\kappa \omega_0 \in \text{Rat}(\mathbb{T})$ in case $\kappa \geq 0$ and $T_{z^\kappa \omega_0} = T_{z^\kappa} T_{\omega_0}$ in case $\kappa < 0$ (by [5, Lemma 5.3]). In both cases it now easily follows that $T_{z^\kappa \omega_0}$ is a Sarason-Toeplitz operator. The claim for T_ω follows since $T_{\omega_+}^{\pm 1} T_z = T_z T_{\omega_+}^{\pm 1}$ and $T_{\omega_-}^{\pm 1} T_z^* = T_z^* T_{\omega_-}^{\pm 1}$. \square

In fact, by the same arguments one can show that T_ω on H^p , $1 < p < \infty$, satisfied (1)-(3) in case T_z^* is replaced by $T_{z^{-1}}$.

REFERENCES

- [1] J.A.W. van Casteren and S. Goldberg, The conjugate of the products of operators. *Studia Mathematica*. **38**, 1970, 125–130.
- [2] A. Cohn, Über die Anzahl der Wurzeln einer algebraischen Gleichung in einem Kreise, *Math. Z.* **14** (1922), 110–148.
- [3] J.B. Conway, *A Course in Functional Analysis*. 2nd Ed. Springer Verlag, 1990.
- [4] S. Goldberg, *Unbounded linear operators: Theory and applications*. McGraw-Hill, New York-Toronto-London, 1966.
- [5] G.J. Groenewald, S. ter Horst, J. Jaftha, A.C.M. Ran, A Toeplitz-like operator with rational symbol having poles on the unit circle I: Fredholm properties, *Oper. Theory Adv. Appl.*, to appear.
- [6] G.J. Groenewald, S. ter Horst, J. Jaftha, A.C.M. Ran, A Toeplitz-like operator with rational symbol having poles on the unit circle II: the spectrum, *Oper. Theory Adv. Appl.*, to appear.
- [7] H. Helson, Large analytic functions. In *Linear Operators in Function Spaces*, Oper. Theory Adv. Appl. **43**, 209 – 216, Birkhäuser, Basel, 1990.
- [8] T. Kato, *Perturbation theory for linear operators*, Reprint of the 1980 edition, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
- [9] V.I. Paulsen and M. Raghupathi, *An introduction to the theory of reproducing kernel Hilbert spaces*, Cambridge Studies in Advanced Mathematics **152**, Cambridge University Press, Cambridge, 2016.
- [10] J.A. Rosenfeld, *Classes of densely defined multiplication and toeplitz operators with applications to extensions of RKHS's*, Ph.D. Thesis, University of Florida, 2013.
- [11] J.A. Rosenfeld, The Sarason sub-symbol and the recovery of the symbol of densely defined Toeplitz operators over the Hardy space, *J. Math. Anal. Appl.* **440** (2016), 911–921.
- [12] D. Sarason, Unbounded Toeplitz operators, *Integral Equations and Operator Theory* **61** (2008), no. 2, 281 – 298.
- [13] M. Schechter, The conjugate of a product of operators. *Journal of Functional Analysis* **6**, 1970, 26 – 28.
- [14] K. Yosida, *Functional Analysis*, Fundamental Principles of Mathematical Sciences **123** (6th ed.), Springer-Verlag, Berlin-New York, 1980.

G.J. GROENEWALD, DEPARTMENT OF MATHEMATICS, UNIT FOR BMI, NORTH-WEST UNIVERSITY, POTCHEFSTROOM, 2531 SOUTH AFRICA
E-mail address: `Gilbert.Groenewald@nwu.ac.za`

S. TER HORST, DEPARTMENT OF MATHEMATICS, UNIT FOR BMI, NORTH-WEST UNIVERSITY, POTCHEFSTROOM, 2531 SOUTH AFRICA
E-mail address: `Sanne.TerHorst@nwu.ac.za`

J. JAFTHA, NUMERACY CENTRE, UNIVERSITY OF CAPE TOWN, RONDEBOSCH 7701; CAPE TOWN; SOUTH AFRICA
E-mail address: `Jacob.Jaftha@uct.ac.za`

A.C.M. RAN, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, VU UNIVERSITY AMSTERDAM, DE BOELELAAN 1081A, 1081 HV AMSTERDAM, THE NETHERLANDS AND UNIT FOR BMI, NORTH-WEST UNIVERSITY, POTCHEFSTROOM, SOUTH AFRICA
E-mail address: `a.c.m.ran@vu.nl`