

# Zeta-values from Euler to Weil-étale cohomology

Alexey Beshenov

Université de Bordeaux / Universiteit Leiden

Advised by Baptiste Morin and Bas Edixhoven

15 May 2017, Leiden



Universiteit  
Leiden



ALGANT  
Erasmus Mundus

2:3

# Outline

---

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .
- ▶ XIX century mathematics:  $\zeta(s)$  and  $\zeta_F(s)$ .

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .
- ▶ XIX century mathematics:  $\zeta(s)$  and  $\zeta_F(s)$ .
- ▶ XX century mathematics:  $\zeta_X(s)$ .

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .
- ▶ XIX century mathematics:  $\zeta(s)$  and  $\zeta_F(s)$ .
- ▶ XX century mathematics:  $\zeta_X(s)$ .
- ▶ Algebraic  $K$ -theory.

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .
- ▶ XIX century mathematics:  $\zeta(s)$  and  $\zeta_F(s)$ .
- ▶ XX century mathematics:  $\zeta_X(s)$ .
- ▶ Algebraic  $K$ -theory.
- ▶ Motivic cohomology.

# Outline

---

- ▶ XVIII century mathematics:  $\sum_{n \geq 1} \frac{1}{n^{2k}}$ .
- ▶ XIX century mathematics:  $\zeta(s)$  and  $\zeta_F(s)$ .
- ▶ XX century mathematics:  $\zeta_X(s)$ .
- ▶ Algebraic  $K$ -theory.
- ▶ Motivic cohomology.
- ▶ Weil-étale cohomology.



# Riemann zeta function before Riemann

# Riemann zeta function before Riemann

---

- **Pietro Mengoli**, 1644, the “Basel problem”:

$$\sum_{n \geq 1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \cdots = \text{?????}$$

# Riemann zeta function before Riemann

---

- ▶ **Pietro Mengoli**, 1644, the “Basel problem”:

$$\sum_{n \geq 1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \cdots = \text{?????}$$

- ▶ **Euler** (“*De summis serierum reciprocarum*”, 1740):  $\frac{\pi^2}{6}$ .

# Riemann zeta function before Riemann

---

- ▶ **Pietro Mengoli**, 1644, the “Basel problem”:

$$\sum_{n \geq 1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \cdots = \text{?????}$$

- ▶ **Euler** (“*De summis serierum reciprocarum*”, 1740):  $\frac{\pi^2}{6}$ .

- ▶ In general (ibid.),

$$\sum_{n \geq 1} \frac{1}{n^{2k}} = (-1)^{k+1} B_{2k} \frac{2^{2k-1}}{(2k)!} \pi^{2k}.$$

# Riemann zeta function before Riemann

---

- ▶ **Pietro Mengoli**, 1644, the “Basel problem”:

$$\sum_{n \geq 1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \dots = \text{?????}$$

- ▶ **Euler** (“*De summis serierum reciprocarum*”, 1740):  $\frac{\pi^2}{6}$ .

- ▶ In general (ibid.),

$$\sum_{n \geq 1} \frac{1}{n^{2k}} = (-1)^{k+1} B_{2k} \frac{2^{2k-1}}{(2k)!} \pi^{2k}.$$

- ▶ Bernoulli numbers:

$$B_0 = 1, B_1 = \frac{1}{2}, B_2 = \frac{1}{6}, B_3 = 0, B_4 = -\frac{1}{30}, B_5 = 0, \\ B_6 = \frac{1}{42}, B_7 = 0, B_8 = -\frac{1}{30}, B_9 = 0, B_{10} = \frac{5}{66}, \dots$$

(**Jacob Bernoulli**, “*Ars Conjectandi*”, 1713).

# Riemann zeta function before Riemann

---

- ▶ **Pietro Mengoli**, 1644, the “Basel problem”:

$$\sum_{n \geq 1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \dots = \text{?????}$$

- ▶ **Euler** (“*De summis serierum reciprocarum*”, 1740):  $\frac{\pi^2}{6}$ .

- ▶ In general (ibid.),

$$\sum_{n \geq 1} \frac{1}{n^{2k}} = (-1)^{k+1} B_{2k} \frac{2^{2k-1}}{(2k)!} \pi^{2k}.$$

- ▶ Bernoulli numbers:

$$B_0 = 1, B_1 = \frac{1}{2}, B_2 = \frac{1}{6}, B_3 = 0, B_4 = -\frac{1}{30}, B_5 = 0, \\ B_6 = \frac{1}{42}, B_7 = 0, B_8 = -\frac{1}{30}, B_9 = 0, B_{10} = \frac{5}{66}, \dots$$

(**Jacob Bernoulli**, “*Ars Conjectandi*”, 1713).

- ▶ Faulhaber’s formula:

$$\sum_{1 \leq i \leq n} i^k = \frac{1}{k+1} \sum_{0 \leq i \leq k} \binom{k+1}{i} B_i n^{k+1-i}$$

(**Johann Faulhaber**, “*Academia Algebrae*”, 1631;  
Bernoulli, 1713).

# Riemann zeta function

---

**Riemann**, *“Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse”* (1859):

# Riemann zeta function

---

**Riemann**, "*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*" (1859):

►  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad (\operatorname{Re} s > 1).$



# Riemann zeta function

---

**Riemann**, “*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*” (1859):

- ▶  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s}$  ( $\text{Re } s > 1$ ).
- ▶ **Euler** (“*Variæ observationes circa series infinitas*”, 1744):  
$$= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}}.$$

# Riemann zeta function

---

**Riemann**, “*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*” (1859):

- ▶  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s}$  ( $\operatorname{Re} s > 1$ ).
- ▶ **Euler** (“*Variæ observationes circa series infinitas*”, 1744):  
 $= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}}.$
- ▶ Meromorphic continuation to  $\mathbb{C}$  with one simple pole at  $s = 1$ .

# Riemann zeta function

---

**Riemann**, “*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*” (1859):

- ▶  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s}$  ( $\operatorname{Re} s > 1$ ).
- ▶ **Euler** (“*Variæ observationes circa series infinitas*”, 1744):  
 $= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}}$ .
- ▶ Meromorphic continuation to  $\mathbb{C}$  with one simple pole at  $s = 1$ .
- ▶ Functional equation

$$\zeta(1-s) = 2(2\pi)^{-s} \Gamma(s) \cos\left(\frac{\pi s}{2}\right) \zeta(s).$$

# Riemann zeta function

---

**Riemann**, “*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*” (1859):

- ▶  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s}$  ( $\text{Re } s > 1$ ).
- ▶ **Euler** (“*Variæ observationes circa series infinitas*”, 1744):  
 $= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}}$ .
- ▶ Meromorphic continuation to  $\mathbb{C}$  with one simple pole at  $s = 1$ .
- ▶ Functional equation

$$\zeta(1-s) = 2(2\pi)^{-s} \Gamma(s) \cos\left(\frac{\pi s}{2}\right) \zeta(s).$$

- ▶ (Trivial) simple zeros at  $s = -2, -4, -6, \dots$

# Riemann zeta function

---


**Riemann**, “*Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*” (1859):


- ▶  $\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s}$  ( $\operatorname{Re} s > 1$ ).
- ▶ **Euler** (“*Variæ observationes circa series infinitas*”, 1744):  
 $= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}}$ .
- ▶ Meromorphic continuation to  $\mathbb{C}$  with one simple pole at  $s = 1$ .
- ▶ Functional equation

$$\zeta(1-s) = 2(2\pi)^{-s} \Gamma(s) \cos\left(\frac{\pi s}{2}\right) \zeta(s).$$

- ▶ (Trivial) simple zeros at  $s = -2, -4, -6, \dots$
- ▶ Euler’s calculation  $\zeta(2k) = (-1)^{k+1} B_{2k} \frac{2^{2k-1}}{(2k)!} \pi^{2k}$  becomes

$$\zeta(-n) = -\frac{B_{n+1}}{n+1} \text{ for } n = 1, 2, 3, 4, \dots$$





# Riemann zeta function at positive odd integers



# Riemann zeta function at positive odd integers

- ▶ **Roger Apéry**, 1977:  $\zeta(3) = 1.2020569\dots$  is irrational.

# Riemann zeta function at positive odd integers

- ▶ **Roger Apéry**, 1977:  $\zeta(3) = 1.2020569\dots$  is irrational.
- ▶ **Tanguy Rivoal**, 2000: infinitely many irrationals  $\zeta(2k + 1)$ .

# Riemann zeta function at positive odd integers

- ▶ **Roger Apéry**, 1977:  $\zeta(3) = 1.2020569\dots$  is irrational.
- ▶ **Tanguy Rivoal**, 2000: infinitely many irrationals  $\zeta(2k + 1)$ .
- ▶ **Wadim Zudilin**, 2001: at least one irrational among  $\zeta(5)$ ,  $\zeta(7)$ ,  $\zeta(9)$ ,  $\zeta(11)$  (which one?)

# Riemann zeta function at positive odd integers

- ▶ **Roger Apéry**, 1977:  $\zeta(3) = 1.2020569\dots$  is irrational.
- ▶ **Tanguy Rivoal**, 2000: infinitely many irrationals  $\zeta(2k + 1)$ .
- ▶ **Wadim Zudilin**, 2001: at least one irrational among  $\zeta(5)$ ,  $\zeta(7)$ ,  $\zeta(9)$ ,  $\zeta(11)$  (which one?)
- ▶ Conjecture:  $\zeta(2k + 1)$  are transcendental, algebraically independent.

# Dedekind zeta function: definition

---

# Dedekind zeta function: definition

---

- ▶  $F/\mathbb{Q}$  — number field,  $\mathcal{O}_F :=$  the ring of integers.

# Dedekind zeta function: definition

---

- ▶  $F/\mathbb{Q}$  — number field,  $\mathcal{O}_F :=$  the ring of integers.
- ▶ **Dedekind**, appendix to Dirichlet's "Vorlesungen über Zahlentheorie" (1863):

$$\zeta_F(s) := \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_F \\ \mathfrak{a} \neq 0}} \frac{1}{N_{F/\mathbb{Q}}(\mathfrak{a})^s} = \prod_{\substack{\mathfrak{p} \subset \mathcal{O}_F \\ \text{prime}}} \frac{1}{1 - N_{F/\mathbb{Q}}(\mathfrak{p})^{-s}}. \quad (\operatorname{Re} s > 1)$$

# Dedekind zeta function: definition

---

- ▶  $F/\mathbb{Q}$  — number field,  $\mathcal{O}_F :=$  the ring of integers.
- ▶ **Dedekind**, appendix to Dirichlet's “Vorlesungen über Zahlentheorie” (1863):

$$\zeta_F(s) := \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_F \\ \mathfrak{a} \neq 0}} \frac{1}{N_{F/\mathbb{Q}}(\mathfrak{a})^s} = \prod_{\substack{\mathfrak{p} \subset \mathcal{O}_F \\ \text{prime}}} \frac{1}{1 - N_{F/\mathbb{Q}}(\mathfrak{p})^{-s}}. \quad (\operatorname{Re} s > 1)$$

- ▶ Note:  $\zeta_{\mathbb{Q}}(s) = \zeta(s)$ .



# Dedekind zeta function: functional equation

# Dedekind zeta function: functional equation

---

- **Hecke**, “Über die Zetafunktion beliebiger algebraischer Zahlkörper”, 1917: meromorphic continuation with simple pole at  $s = 1$ ; functional equation

$$\zeta_F(1-s) = |\Delta_F|^{s-1/2} \left(\cos \frac{\pi s}{2}\right)^{r_1+r_2} \left(\sin \frac{\pi s}{2}\right)^{r_2} (2(2\pi)^{-s} \Gamma(s))^d \zeta_F(s),$$

where

$r_1 :=$  real places,  $r_2 :=$  conjugate pairs of complex places;  
 $d := [F : \mathbb{Q}] = r_1 + 2r_2$  and  $\Delta_F :=$  discriminant.

# Dedekind zeta function: functional equation

---

- **Hecke**, “Über die Zetafunktion beliebiger algebraischer Zahlkörper”, 1917: meromorphic continuation with simple pole at  $s = 1$ ; functional equation

$$\zeta_F(1-s) = |\Delta_F|^{s-1/2} \left(\cos \frac{\pi s}{2}\right)^{r_1+r_2} \left(\sin \frac{\pi s}{2}\right)^{r_2} (2(2\pi)^{-s} \Gamma(s))^d \zeta_F(s),$$

where

$r_1 :=$  real places,  $r_2 :=$  conjugate pairs of complex places;  
 $d := [F : \mathbb{Q}] = r_1 + 2r_2$  and  $\Delta_F :=$  discriminant.

# Dedekind zeta function: functional equation

- **Hecke**, “Über die Zetafunktion beliebiger algebraischer Zahlkörper”, 1917: meromorphic continuation with simple pole at  $s = 1$ ; functional equation

$$\zeta_F(1-s) = |\Delta_F|^{s-1/2} \left(\cos \frac{\pi s}{2}\right)^{r_1+r_2} \left(\sin \frac{\pi s}{2}\right)^{r_2} (2(2\pi)^{-s} \Gamma(s))^d \zeta_F(s),$$

where

$r_1 :=$  real places,  $r_2 :=$  conjugate pairs of complex places;  
 $d := [F : \mathbb{Q}] = r_1 + 2r_2$  and  $\Delta_F :=$  discriminant.

- (Trivial) zeros:

$s:$	0	-1	-2	-3	-4	-5	...
order:	$r_1 + r_2 - 1$	$r_2$	$r_1 + r_2$	$r_2$	$r_1 + r_2$	$r_2$	...

# Class number formula

---

**Dirichlet**, *“Recherches sur diverses applications de l’analyse infinitésimale à la théorie des nombres”* (1839);

\* **Gauss**, *“Disquisitiones Arithmeticae”* (1801):

# Class number formula

---

**Dirichlet**, “*Recherches sur diverses applications de l’analyse infinitésimale à la théorie des nombres*” (1839);

\* **Gauss**, “*Disquisitiones Arithmeticae*” (1801):

- Pole at  $s = 1$ :

$$\lim_{s \rightarrow 1} (s - 1) \zeta_F(s) = \frac{2^{r_1} (2\pi)^{r_2} \# \text{Cl}(F)}{\# \mu_F \cdot \sqrt{|\Delta_F|}} R_F,$$

where  $\text{Cl}(F)$  — class group;  $\mu_F \subset \mathcal{O}_F^\times$  — roots of unity;  
 $R_F$  — Dirichlet regulator.

# Class number formula

---

**Dirichlet**, “*Recherches sur diverses applications de l’analyse infinitésimale à la théorie des nombres*” (1839);

\* **Gauss**, “*Disquisitiones Arithmeticae*” (1801):

- Pole at  $s = 1$ :

$$\lim_{s \rightarrow 1} (s - 1) \zeta_F(s) = \frac{2^{r_1} (2\pi)^{r_2} \# \text{Cl}(F)}{\# \mu_F \cdot \sqrt{|\Delta_F|}} R_F,$$

where  $\text{Cl}(F)$  — class group;  $\mu_F \subset \mathcal{O}_F^\times$  — roots of unity;  
 $R_F$  — Dirichlet regulator.

- Zero at  $s = 0$ :

$$\lim_{s \rightarrow 0} s^{-(r_1+r_2-1)} \zeta_F(s) = -\frac{\# \text{Cl}(F)}{\# \mu_F} R_F.$$

# Values of the Dedekind zeta function

---



# Values of the Dedekind zeta function

---

- ▶ If  $F$  is totally real ( $r_2 = 0$ ), then  $\zeta_F(-n) \neq 0$  for odd  $n$ .

# Values of the Dedekind zeta function

---

- ▶ If  $F$  is totally real ( $r_2 = 0$ ), then  $\zeta_F(-n) \neq 0$  for odd  $n$ .
- ▶ “Siegel–Klingen theorem”, 1961:  $\zeta_F(-n) \in \mathbb{Q}$ .

# Values of the Dedekind zeta function

---

- ▶ If  $F$  is totally real ( $r_2 = 0$ ), then  $\zeta_F(-n) \neq 0$  for odd  $n$ .
- ▶ “Siegel–Klingen theorem”, 1961:  $\zeta_F(-n) \in \mathbb{Q}$ .
- ▶ **Günter Harder**, “A Gauss–Bonnet formula for discrete arithmetically defined groups”, 1971:

$$\chi(\mathrm{Sp}_{2n}(\mathcal{O}_F)) = \frac{1}{2^{n(d-n)}} \prod_{1 \leq i \leq n} \zeta_F(1 - 2i).$$

# Values of the Dedekind zeta function

---

- ▶ If  $F$  is totally real ( $r_2 = 0$ ), then  $\zeta_F(-n) \neq 0$  for odd  $n$ .
- ▶ “Siegel–Klingen theorem”, 1961:  $\zeta_F(-n) \in \mathbb{Q}$ .
- ▶ **Günter Harder**, “A Gauss–Bonnet formula for discrete arithmetically defined groups”, 1971:

$$\chi(\mathrm{Sp}_{2n}(\mathcal{O}_F)) = \frac{1}{2^{n(d-n)}} \prod_{1 \leq i \leq n} \zeta_F(1 - 2i).$$

- ▶ Example:  $F = \mathbb{Q}$ ,  $n = 1$ ,  $\mathrm{Sp}_2 = \mathrm{SL}_2$ ,

$$\chi(\mathrm{SL}_2(\mathbb{Z})) = -\frac{1}{12} = -\frac{B_2}{2} = \zeta(-1)$$

(“orbifold Euler characteristic” of  $\mathcal{H}/\mathrm{SL}_2(\mathbb{Z})$ ).

# Zeta function of a scheme

---

# Zeta function of a scheme

---

- ▶  $X \rightarrow \operatorname{Spec} \mathbb{Z}$  — arithmetic scheme (separated, of finite type).

# Zeta function of a scheme

---

- ▶  $X \rightarrow \operatorname{Spec} \mathbb{Z}$  — arithmetic scheme (separated, of finite type).
- ▶  $\zeta_X(s) := \prod_{x \in X_0} \frac{1}{1 - N(x)^{-s}}$ .      ( $\operatorname{Re} s > \dim X$ ).  
 $X_0 :=$  closed points;  $N(x) := \#$ residue field at  $x$ .

# Zeta function of a scheme

---

- ▶  $X \rightarrow \operatorname{Spec} \mathbb{Z}$  — arithmetic scheme (separated, of finite type).
- ▶  $\zeta_X(s) := \prod_{x \in X_0} \frac{1}{1 - N(x)^{-s}}$ .      ( $\operatorname{Re} s > \dim X$ ).  
 $X_0 :=$  closed points;  $N(x) := \#$ residue field at  $x$ .
- ▶ Note:  $\zeta_{\operatorname{Spec} \mathbb{Z}}(s) = \zeta(s)$  and  $\zeta_{\operatorname{Spec} \mathcal{O}_F}(s) = \zeta_F(s)$ .



# Zeta function of a scheme

---

- ▶  $X \rightarrow \operatorname{Spec} \mathbb{Z}$  — arithmetic scheme (separated, of finite type).
- ▶  $\zeta_X(s) := \prod_{x \in X_0} \frac{1}{1 - N(x)^{-s}}$ .      ( $\operatorname{Re} s > \dim X$ ).  
 $X_0 :=$  closed points;  $N(x) := \#$ residue field at  $x$ .
- ▶ Note:  $\zeta_{\operatorname{Spec} \mathbb{Z}}(s) = \zeta(s)$  and  $\zeta_{\operatorname{Spec} \mathcal{O}_F}(s) = \zeta_F(s)$ .
- ▶ Conjecture (!): meromorphic continuation and a functional equation  $\zeta_X(s) \leftrightarrow \zeta_X(\dim X - s)$ .

# Zeta function of a scheme

---

- ▶  $X \rightarrow \operatorname{Spec} \mathbb{Z}$  — arithmetic scheme (separated, of finite type).
- ▶  $\zeta_X(s) := \prod_{x \in X_0} \frac{1}{1 - N(x)^{-s}}$ .      ( $\operatorname{Re} s > \dim X$ ).  
 $X_0 :=$  closed points;  $N(x) := \#$ residue field at  $x$ .
- ▶ Note:  $\zeta_{\operatorname{Spec} \mathbb{Z}}(s) = \zeta(s)$  and  $\zeta_{\operatorname{Spec} \mathcal{O}_F}(s) = \zeta_F(s)$ .
- ▶ Conjecture (!): meromorphic continuation and a functional equation  $\zeta_X(s) \leftrightarrow \zeta_X(\dim X - s)$ .
- ▶ Special values may be studied via  
 $K$ -theory  $K_n(X)$  or motivic cohomology  $H^i(X, \mathbb{Z}(n))$ .

# Algebraic K-theory

---

# Algebraic K-theory

---

- Input: an “exact category”  $\mathcal{C}$ .

Examples:  $\mathbf{VB}(X)$  and  $R\text{-}\mathbf{Proj}_{fg} \simeq \mathbf{VB}(\mathrm{Spec} R)$ .

# Algebraic K-theory

---

- ▶ Input: an “exact category”  $\mathcal{C}$ .  
Examples:  $\mathbf{VB}(X)$  and  $R\text{-}\mathbf{Proj}_{fg} \simeq \mathbf{VB}(\mathrm{Spec} R)$ .
- ▶ **Grothendieck**, 1957  
(work on [Grothendieck–Hirzebruch]–Riemann–Roch):

$$K_0(\mathcal{C}) := \frac{\mathbb{Z} \langle \text{isomorphism classes of objects of } \mathcal{C} \rangle}{[B] = [A] + [C] \text{ for each s.e.s. } 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0}.$$

# Algebraic K-theory

---

- Input: an “exact category”  $\mathcal{C}$ .

Examples:  $\mathbf{VB}(X)$  and  $R\text{-}\mathbf{Proj}_{fg} \simeq \mathbf{VB}(\mathrm{Spec} R)$ .

- **Grothendieck**, 1957

(work on [Grothendieck–Hirzebruch]–Riemann–Roch):

$$K_0(\mathcal{C}) := \frac{\mathbb{Z} \langle \text{isomorphism classes of objects of } \mathcal{C} \rangle}{[B] = [A] + [C] \text{ for each s.e.s. } 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0}.$$


- **Quillen**, 1973:

$$K_0(\mathcal{C}) \cong \pi_1(BQC, 0),$$

$$K_n(\mathcal{C}) := \pi_{n+1}(BQC, 0);$$

$Q$  — Quillen’s “Q-construction”,

$B$  — geometric realization of the nerve.



"I THINK YOU SHOULD BE MORE EXPLICIT HERE IN STEP TWO."

## Some of the few known calculations



# Some of the few known calculations

---

- Quillen, 1972:

$$K_0(\mathbb{F}_q) \cong \mathbb{Z},$$

$$K_{2n}(\mathbb{F}_q) = 0,$$

$$K_{2n-1}(\mathbb{F}_q) \cong \mathbb{Z}/(q^n - 1)\mathbb{Z}.$$

# Some of the few known calculations

---

- Quillen, 1972:

$$K_0(\mathbb{F}_q) \cong \mathbb{Z},$$

$$K_{2n}(\mathbb{F}_q) = 0,$$

$$K_{2n-1}(\mathbb{F}_q) \cong \mathbb{Z}/(q^n - 1)\mathbb{Z}.$$

- Note:  $\#K_{2n-1}(\mathbb{F}_q) = -\zeta_{\mathbb{F}_q}(-n)^{-1}$ .

# Some of the few known calculations

---

- ▶ **Quillen**, 1972:

$$K_0(\mathbb{F}_q) \cong \mathbb{Z},$$

$$K_{2n}(\mathbb{F}_q) = 0,$$

$$K_{2n-1}(\mathbb{F}_q) \cong \mathbb{Z}/(q^n - 1)\mathbb{Z}.$$

- ▶ Note:  $\#K_{2n-1}(\mathbb{F}_q) = -\zeta_{\mathbb{F}_q}(-n)^{-1}$ .
- ▶ **Quillen**, 1973:  $K_n(\mathcal{O}_F)$  are finitely generated.

# Some of the few known calculations

---

- **Quillen**, 1972:

$$\begin{aligned}K_0(\mathbb{F}_q) &\cong \mathbb{Z}, \\K_{2n}(\mathbb{F}_q) &= 0, \\K_{2n-1}(\mathbb{F}_q) &\cong \mathbb{Z}/(q^n - 1)\mathbb{Z}.\end{aligned}$$

- Note:  $\#K_{2n-1}(\mathbb{F}_q) = -\zeta_{\mathbb{F}_q}(-n)^{-1}$ .
- **Quillen**, 1973:  $K_n(\mathcal{O}_F)$  are finitely generated.
- **Armand Borel**, 1974:

$$\mathrm{rk} K_n(\mathcal{O}_F) = \begin{cases} 0, & n = 2k, \\ r_1 + r_2, & n = 4k + 1, \\ r_2, & n = 4k - 1. \end{cases} \quad (k > 0)$$

# Some of the few known calculations

---

- **Quillen**, 1972:

$$\begin{aligned}K_0(\mathbb{F}_q) &\cong \mathbb{Z}, \\K_{2n}(\mathbb{F}_q) &= 0, \\K_{2n-1}(\mathbb{F}_q) &\cong \mathbb{Z}/(q^n - 1)\mathbb{Z}.\end{aligned}$$

- Note:  $\#K_{2n-1}(\mathbb{F}_q) = -\zeta_{\mathbb{F}_q}(-n)^{-1}$ .
- **Quillen**, 1973:  $K_n(\mathcal{O}_F)$  are finitely generated.
- **Armand Borel**, 1974:

$$\mathrm{rk} K_n(\mathcal{O}_F) = \begin{cases} 0, & n = 2k, \\ r_1 + r_2, & n = 4k + 1, \\ r_2, & n = 4k - 1. \end{cases} \quad (k > 0)$$

- Note:  $\mathrm{rk} K_{2n+1}(\mathcal{O}_F) = \text{order of zero of } \zeta_F(s) \text{ at } s = -n$ .

# Torsion in the K-theory of $\mathbb{Z}$

---

# Torsion in the K-theory of $\mathbb{Z}$

---

- **Milnor**, 1971:  $K_2(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ .

# Torsion in the K-theory of $\mathbb{Z}$

---

- ▶ **Milnor**, 1971:  $K_2(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ .
- ▶ **Lee, Szczarba**, 1976:  $K_3(\mathbb{Z}) \cong \mathbb{Z}/48\mathbb{Z}$ .



# Torsion in the K-theory of $\mathbb{Z}$

---

- ▶ **Milnor**, 1971:  $K_2(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ .
- ▶ **Lee, Szczarba**, 1976:  $K_3(\mathbb{Z}) \cong \mathbb{Z}/48\mathbb{Z}$ .
- ▶ **Rognes**, 2000:  $K_4(\mathbb{Z}) = 0$ .

# Torsion in the K-theory of $\mathbb{Z}$

---

- ▶ **Milnor**, 1971:  $K_2(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ .
- ▶ **Lee, Szczarba**, 1976:  $K_3(\mathbb{Z}) \cong \mathbb{Z}/48\mathbb{Z}$ .
- ▶ **Rognes**, 2000:  $K_4(\mathbb{Z}) = 0$ .
- ▶ **Elbaz-Vincent, Gangl, Soulé**, 2002:  $K_5(\mathbb{Z}) \cong \mathbb{Z}$ .

# Torsion in the K-theory of $\mathbb{Z}$

- ▶ **Milnor**, 1971:  $K_2(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ .
- ▶ **Lee, Szczarba**, 1976:  $K_3(\mathbb{Z}) \cong \mathbb{Z}/48\mathbb{Z}$ .
- ▶ **Rognes**, 2000:  $K_4(\mathbb{Z}) = 0$ .
- ▶ **Elbaz-Vincent, Gangl, Soulé**, 2002:  $K_5(\mathbb{Z}) \cong \mathbb{Z}$ .
- ▶ Using the Bloch–Kato conjecture (**Voevodsky, Rost, ...**):

$n$ :	2	3	4	5
$K_n(\mathbb{Z})$ :	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/48\mathbb{Z}$	0	$\mathbb{Z}$
$n$ :	6	7	8	9
$K_n(\mathbb{Z})$ :	0	$\mathbb{Z}/240\mathbb{Z}$	(0?)	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
$n$ :	10	11	12	13
$K_n(\mathbb{Z})$ :	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/1008\mathbb{Z}$	(0?)	$\mathbb{Z}$
$n$ :	14	15	16	17
$K_n(\mathbb{Z})$ :	0	$\mathbb{Z}/480\mathbb{Z}$	(0?)	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
$n$ :	18	19	20	21
$K_n(\mathbb{Z})$ :	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/528\mathbb{Z}$	(0?)	$\mathbb{Z}$
$n$ :	22	23	24	25
$K_n(\mathbb{Z})$ :	$\mathbb{Z}/691\mathbb{Z}$	$\mathbb{Z}/65\,520\mathbb{Z}$	(0?)	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$

# Lichtenbaum's conjecture

---

# Lichtenbaum's conjecture

---

- Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .

# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre**, 1967):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .

# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre**, 1967):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .
- ▶ Dirichlet's unit theorem:  $\mathcal{O}_F^\times \cong \mathbb{Z}^{r_1+r_2-1} \oplus \mu_F$ .

# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre, 1967**):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .
- ▶ Dirichlet's unit theorem:  $\mathcal{O}_F^\times \cong \mathbb{Z}^{r_1+r_2-1} \oplus \mu_F$ .
- ▶ Class number formula:

$$\lim_{s \rightarrow 0} s^{-(r_1+r_2-1)} \zeta_F(s) = -\frac{\#K_0(\mathcal{O}_F)_{\text{tors}}}{\#K_1(\mathcal{O}_F)_{\text{tors}}} R_F.$$



# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre**, 1967):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .
- ▶ Dirichlet's unit theorem:  $\mathcal{O}_F^\times \cong \mathbb{Z}^{r_1+r_2-1} \oplus \mu_F$ .
- ▶ Class number formula:

$$\lim_{s \rightarrow 0} s^{-(r_1+r_2-1)} \zeta_F(s) = - \frac{\#K_0(\mathcal{O}_F)_{\text{tors}}}{\#K_1(\mathcal{O}_F)_{\text{tors}}} R_F.$$

- ▶ **Lichtenbaum**, 1973:

$$\lim_{s \rightarrow n} (n-s)^{-\mu_n} \zeta_F(-s) = \pm 2^? \frac{\#K_{2n}(\mathcal{O}_F)}{\#K_{2n+1}(\mathcal{O}_F)_{\text{tors}}} R_{F,n}.$$

$R_{F,n}$  — “higher regulators” (Borel, Beilinson).

# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre**, 1967):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .
- ▶ Dirichlet's unit theorem:  $\mathcal{O}_F^\times \cong \mathbb{Z}^{r_1+r_2-1} \oplus \mu_F$ .
- ▶ Class number formula:

$$\lim_{s \rightarrow 0} s^{-(r_1+r_2-1)} \zeta_F(s) = -\frac{\#K_0(\mathcal{O}_F)_{\text{tors}}}{\#K_1(\mathcal{O}_F)_{\text{tors}}} R_F.$$

- ▶ **Lichtenbaum**, 1973:

$$\lim_{s \rightarrow n} (n-s)^{-\mu_n} \zeta_F(-s) = \pm 2^? \frac{\#K_{2n}(\mathcal{O}_F)}{\#K_{2n+1}(\mathcal{O}_F)_{\text{tors}}} R_{F,n}.$$

$R_{F,n}$  — “higher regulators” (Borel, Beilinson).

- ▶ Example:  $\zeta(-1) = -\frac{B_2}{2} = -\frac{1}{12}$ ,  $\frac{\#K_2(\mathbb{Z})}{\#K_3(\mathbb{Z})} = \frac{\#\mathbb{Z}/2}{\#\mathbb{Z}/48} = \frac{1}{24}$ .

# Lichtenbaum's conjecture

---

- ▶ Easy:  $K_0(\mathcal{O}_F) \cong \text{Cl}(F) \oplus \mathbb{Z}$ .
- ▶ Not-so-easy (**Bass, Milnor, Serre, 1967**):  $K_1(\mathcal{O}_F) \cong \mathcal{O}_F^\times$ .
- ▶ Dirichlet's unit theorem:  $\mathcal{O}_F^\times \cong \mathbb{Z}^{r_1+r_2-1} \oplus \mu_F$ .
- ▶ Class number formula:

$$\lim_{s \rightarrow 0} s^{-(r_1+r_2-1)} \zeta_F(s) = -\frac{\#K_0(\mathcal{O}_F)_{\text{tors}}}{\#K_1(\mathcal{O}_F)_{\text{tors}}} R_F.$$

- ▶ **Lichtenbaum, 1973:**

$$\lim_{s \rightarrow n} (n-s)^{-\mu_n} \zeta_F(-s) = \pm 2^? \frac{\#K_{2n}(\mathcal{O}_F)}{\#K_{2n+1}(\mathcal{O}_F)_{\text{tors}}} R_{F,n}.$$

$R_{F,n}$  — “higher regulators” (Borel, Beilinson).

- ▶ Example:  $\zeta(-1) = -\frac{B_2}{2} = -\frac{1}{12}$ ,  $\frac{\#K_2(\mathbb{Z})}{\#K_3(\mathbb{Z})} = \frac{\#\mathbb{Z}/2}{\#\mathbb{Z}/48} = \frac{1}{24}$ .
- ▶ Example:  $\zeta(-11) = -\frac{B_{12}}{12} = \frac{691}{12 \cdot 2730} = \frac{691}{2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13}$ ,  
 $\frac{\#K_{22}(\mathbb{Z})}{\#K_{23}(\mathbb{Z})} = \frac{691}{65520} = \frac{691}{2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13}$ .

# Motivic cohomology

---

# Motivic cohomology

---

- ▶ **Bloch**, 1986: complexes of Zariski / étale sheaves  $\mathbb{Z}(n)$ .

# Motivic cohomology

---

- ▶ **Bloch**, 1986: complexes of Zariski / étale sheaves  $\mathbb{Z}(n)$ .
- ▶  $H^i(X_{\text{Zar}}, \mathbb{Z}(n))$ ,  $H^i(X_{\text{ét}}, \mathbb{Z}(n))$  — (hyper)cohomology groups.

# Motivic cohomology

---

- ▶ **Bloch**, 1986: complexes of Zariski / étale sheaves  $\mathbb{Z}(n)$ .
- ▶  $H^i(X_{\text{Zar}}, \mathbb{Z}(n))$ ,  $H^i(X_{\text{ét}}, \mathbb{Z}(n))$  — (hyper)cohomology groups.
- ▶ Finite generation, boundedness for arithmetic  $X$ : conjectures.

# Motivic cohomology

---

- ▶ **Bloch**, 1986: complexes of Zariski / étale sheaves  $\mathbb{Z}(n)$ .
- ▶  $H^i(X_{\text{Zar}}, \mathbb{Z}(n))$ ,  $H^i(X_{\text{ét}}, \mathbb{Z}(n))$  — (hyper)cohomology groups.
- ▶ Finite generation, boundedness for arithmetic  $X$ : conjectures.
- ▶ Possible motivation (no pun intended):

$$E_2^{pq} = H^p(X, \mathbb{Z}(q)) \implies K_{2q-p}(X),$$

similar to the Atiyah–Hirzebruch spectral sequence.



# Motivic cohomology

---

- ▶ **Bloch**, 1986: complexes of Zariski / étale sheaves  $\mathbb{Z}(n)$ .
- ▶  $H^i(X_{\text{Zar}}, \mathbb{Z}(n))$ ,  $H^i(X_{\text{ét}}, \mathbb{Z}(n))$  — (hyper)cohomology groups.
- ▶ Finite generation, boundedness for arithmetic  $X$ : conjectures.
- ▶ Possible motivation (no pun intended):

$$E_2^{pq} = H^p(X, \mathbb{Z}(q)) \implies K_{2q-p}(X),$$

similar to the Atiyah–Hirzebruch spectral sequence.

- ▶  $H^i(X_{\text{ét}}, \mathbb{Z}(n))$  might be better for studying the zeta-values.

# Weil-étale cohomology (Lichtenbaum, 2005)

## Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

# Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

1.  $H_{W,c}^i(X, \mathbb{Z}(0))$  are f.g., almost all zero.

# Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

1.  $H_{W,c}^i(X, \mathbb{Z}(0))$  are f.g., almost all zero.
2.  $H_{W,c}^i(X, \mathbb{Z}(0)) \otimes \mathbb{R} \xrightarrow{\cong} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$ .

# Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

1.  $H_{W,c}^i(X, \mathbb{Z}(0))$  are f.g., almost all zero.
2.  $H_{W,c}^i(X, \mathbb{Z}(0)) \otimes \mathbb{R} \xrightarrow{\cong} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$ .
3. For a canonical class  $\theta \in H_W^1(X, \widetilde{\mathbb{R}}(0))$

$$\dots \xrightarrow{\cup \theta} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup \theta} H_{W,c}^{i+1}(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup \theta} \dots$$

a bounded acyclic complex of f.d. vector spaces.

# Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

1.  $H_{W,c}^i(X, \mathbb{Z}(0))$  are f.g., almost all zero.
2.  $H_{W,c}^i(X, \mathbb{Z}(0)) \otimes \mathbb{R} \xrightarrow{\cong} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$ .
3. For a canonical class  $\theta \in H_W^1(X, \widetilde{\mathbb{R}}(0))$

$$\dots \xrightarrow{\cup \theta} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup \theta} H_{W,c}^{i+1}(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup \theta} \dots$$

a bounded acyclic complex of f.d. vector spaces.

4.  $\text{ord}_{s=0} \zeta(X, s) = \sum_{i \geq 0} (-1)^i \cdot i \cdot \text{rk}_{\mathbb{Z}} H_{W,c}^i(X, \mathbb{Z}(0))$ .

# Weil-étale cohomology (Lichtenbaum, 2005)

---

For an arithmetic scheme  $X$  there should (!) exist abelian groups  $H_{W,c}^i(X, \mathbb{Z}(0))$  and real vector spaces  $H_W^i(X, \widetilde{\mathbb{R}}(0))$ ,  $H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$  such that

1.  $H_{W,c}^i(X, \mathbb{Z}(0))$  are f.g., almost all zero.
2.  $H_{W,c}^i(X, \mathbb{Z}(0)) \otimes \mathbb{R} \xrightarrow{\cong} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0))$ .
3. For a canonical class  $\theta \in H_W^1(X, \widetilde{\mathbb{R}}(0))$

$$\dots \xrightarrow{\cup\theta} H_{W,c}^i(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup\theta} H_{W,c}^{i+1}(X, \widetilde{\mathbb{R}}(0)) \xrightarrow{\cup\theta} \dots$$

a bounded acyclic complex of f.d. vector spaces.

4.  $\text{ord}_{s=0} \zeta(X, s) = \sum_{i \geq 0} (-1)^i \cdot i \cdot \text{rk}_{\mathbb{Z}} H_{W,c}^i(X, \mathbb{Z}(0))$ .
5.  $\mathbb{Z} \cdot \lambda(\zeta^*(X, 0)^{-1}) = \bigotimes_{i \in \mathbb{Z}} \det_{\mathbb{Z}} H_{W,c}^i(X, \mathbb{Z}(0))^{(-1)^i}$ ,  
where  $\lambda: \mathbb{R} \xrightarrow{\cong} \left( \bigotimes_{i \in \mathbb{Z}} \det_{\mathbb{Z}} H_{W,c}^i(X, \mathbb{Z}(0))^{(-1)^i} \right) \otimes \mathbb{R}$ .



# Weil-étale cohomology

---

# Weil-étale cohomology

---

- **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.

# Weil-étale cohomology

---

- ▶ **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.
- ▶ **Lichtenbaum**, 2009:  $H_{W,c}^i(X, \mathbb{Z}(0))$  for  $X = \operatorname{Spec} \mathcal{O}_F$ .

# Weil-étale cohomology

---

- ▶ **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.
- ▶ **Lichtenbaum**, 2009:  $H_{W,c}^i(X, \mathbb{Z}(0))$  for  $X = \operatorname{Spec} \mathcal{O}_F$ .
- ▶ **Morin**, 2012:  $H_{W,c}^i(X, \mathbb{Z}(0))$   
for proper regular arithmetic schemes.

# Weil-étale cohomology

---

- ▶ **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.
- ▶ **Lichtenbaum**, 2009:  $H_{W,c}^i(X, \mathbb{Z}(0))$  for  $X = \operatorname{Spec} \mathcal{O}_F$ .
- ▶ **Morin**, 2012:  $H_{W,c}^i(X, \mathbb{Z}(0))$   
for proper regular arithmetic schemes.
- ▶ **Flach, Morin**, 2016:  $H_{W,c}^i(X, \mathbb{Z}(n))$  for  $n \in \mathbb{Z}$   
for proper regular arithmetic schemes.

# Weil-étale cohomology

---

- ▶ **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.
- ▶ **Lichtenbaum**, 2009:  $H_{W,c}^i(X, \mathbb{Z}(0))$  for  $X = \operatorname{Spec} \mathcal{O}_F$ .
- ▶ **Morin**, 2012:  $H_{W,c}^i(X, \mathbb{Z}(0))$   
for proper regular arithmetic schemes.
- ▶ **Flach, Morin**, 2016:  $H_{W,c}^i(X, \mathbb{Z}(n))$  for  $n \in \mathbb{Z}$   
for proper regular arithmetic schemes.
- ▶ My thesis, work in progress:  $H_{W,c}^i(X, \mathbb{Z}(n))$ 
  - ▶ for any arithmetic scheme (makes things harder)
  - ▶ and  $n = -1, -2, -3, -4, \dots$  (makes things easier)

# Weil-étale cohomology

---

- ▶ **Geisser**, 2004; **Lichtenbaum**, 2005: construction of  $H_{W,c}^i(X, \mathbb{Z}(n))$  for smooth varieties over finite fields.
- ▶ **Lichtenbaum**, 2009:  $H_{W,c}^i(X, \mathbb{Z}(0))$  for  $X = \operatorname{Spec} \mathcal{O}_F$ .
- ▶ **Morin**, 2012:  $H_{W,c}^i(X, \mathbb{Z}(0))$   
for proper regular arithmetic schemes.
- ▶ **Flach, Morin**, 2016:  $H_{W,c}^i(X, \mathbb{Z}(n))$  for  $n \in \mathbb{Z}$   
for proper regular arithmetic schemes.
- ▶ My thesis, work in progress:  $H_{W,c}^i(X, \mathbb{Z}(n))$ 
  - ▶ for any arithmetic scheme (makes things harder)
  - ▶ and  $n = -1, -2, -3, -4, \dots$  (makes things easier)
- ▶ Construction via the cycle complexes  $\mathbb{Z}(n)$ , following Flach and Morin (input: conjectures on finite generation and boundedness).

**Thank you!**