Pretend Clinical Care Pathway

Preoperative

Subheader here

Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs
Some example text from an itemized list.
Stop full liquids 8 hrs/midnight.
Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs
Some example text from an itemized list.
Stop full liquids 8 hrs/midnight.
Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Preoperative

Subheader here

Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs

Some example text from an itemized list.

Stop full liquids 8 hrs/midnight.

Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs

Some example text from an itemized list.

Stop full liquids 8 hrs/midnight.

Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Preoperative

Subheader here

Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs

Some example text from an itemized list.

Stop full liquids 8 hrs/midnight.

Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs Some example text from an itemized list. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs

Some example text from an itemized list.

Stop full liquids 8 hrs/midnight.

Stop clear liquids 4 hrs before surgery. More example text.