


## Preoperative



### Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



### Subheader here


- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



### Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

## Intraoperative



### Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.

- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here


- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.


## Postoperative



Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Version XYZ. Last updated Feb 31, 2023. Available online at <https://pathways.stanes.link/pretend>. Submit edits or comments there.