Preoperative



Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Intraoperative



Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.

Postoperative



Subheader here

Example of some paragraph text before a list. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs.

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

- Some example text from an itemized list. Stop full liquids 8 hrs/midnight.
- Stop clear liquids 4 hrs before surgery. More example text.



Subheader here

Example of some paragraph text. Consider allowing breast patients who arrive early to pre-op for lympho or mammography to drink water until 2 hrs. Stop full liquids 8 hrs/midnight. Stop clear liquids 4 hrs before surgery. More example text.