


Interactions, Measurements, and Theory of Radiation Ph.D. Qualifying Exam

1. **(15 Minutes)** A monoenergetic beam of photons is incident upon a 30° copper wedge ($Z=29$, $\rho = 8.96 \text{ g cm}^{-3}$). The exiting uncollided beam intensity is measured at 1.73 cm (denoted by I_1) and 5.2 cm (denoted by I_2) from the top of the wedge, respectively (as shown in the figure below). Given that $I_1/I_2 = 3.26$, what is the energy of the incident photons?


Mass Attenuation Coefficient for Cu		
Energy (MeV)	$\mu_{\text{en}}/\rho \text{ (cm}^2/\text{g)}$	$\mu/\rho \text{ (cm}^2/\text{g)}$
0.01	1.48E+02	2.16E+02
0.05	2.613	2.192
0.1	2.95E-01	4.58E-01
0.2	5.78E-01	1.56E-01
0.4	3.12E-02	9.41E-02
0.6	2.86E-02	7.63E-02
0.8	2.68E-02	6.61E-02
1	2.56E-02	5.90E-02



2. **(20 Minutes)** A neutron with a kinetic energy of 1.0 MeV traveling along the x-axis in the positive-x direction collides with a B-10 nucleus that is traveling along the x-axis in the negative-x direction (that is, directly towards the neutron). The B-10 nucleus has a kinetic energy of 0.1 MeV. The neutron is absorbed into the B-10 nucleus and an α -particle and Li-7 nucleus are emitted from the reaction. The α -particle has a kinetic energy of 1.5 MeV. The Li-7 nucleus following the reaction is left in an unexcited state. Determine the energy and direction (with respect to the original direction of the neutron) of the Li-7 nucleus. Note that the B-10 target nucleus was not at rest when it interacted with the neutron.
3. **(15 Minutes)** In this problem you will be asked to estimate the concentration of an activity in units of disintegrations per minute per unit mass (dpm/g) **and** the uncertainty of the measurement. It is not necessary for you to know the identity of the radionuclide or the half-life of the material to answer this question correctly.

A one gram sample is to be processed in a chemical laboratory. A series of blank chemical processes showed that the chemical recovery (also known as the chemical yield) was $70 \pm 5\%$. In addition, before the measurement, it was determined that the counting efficiency for this radionuclide was $20 \pm 2\%$. After processing the sample was counted in the same exact geometry using a measurement system with a precise timer. The measured count rate of the sample was 30 ± 5 counts per minute (cpm). Calculate the following:

- a. The activity concentration of the sample.
 - b. The associated uncertainty in the activity concentration.
4. **(15 minutes)** Consider a homogeneous slab of thickness $2X$ (cm) placed in a vacuum. The medium is purely absorbing with macroscopic cross section, Σ_a (cm^{-1}).
- a. A uniform isotropic volumetric source, Q_v ($\text{n/cm}^3\text{-s}$), is present in the slab. Using the integro-differential transport equation, compute the angular flux ($\text{n/cm}^2\text{-s-str}$) at an arbitrary point within the slab for $\mu > 0$.
 - b. An isotropic surface source, Q_s ($\text{n/cm}^2\text{-s}$), is placed at the left face of the slab. Using the integro-differential transport equation, compute the angular flux ($\text{n/cm}^2\text{-s-str}$) at an arbitrary point within the slab.
5. **(15 minutes)** Estimate the minimum mass of a neutron detector if it must produce one measureable energy deposition event per second when exposed to a fluence rate of $1 \text{ n/cm}^2\text{-s}$. Assume that the neutron kerma factor for the relevant material and detector configuration is $2 \times 10^{-11} \text{ Gy-cm}^2/\text{n}$, and that the energy deposited by a neutron interaction in this detector is 1 MeV .
6. **(15 minutes)** Consider neutron, proton and α -particle beams. Which nuclides would you select as target materials for producing ${}^{24}_{11}\text{Na}$ using each of these beams? To get credit, you must justify your answer.
7. **(10 minutes)** You are asked to construct a device that will provide minimum uninterrupted thermal power of 40 W for 40 years using the alpha decay of ${}^{238}\text{Pu}$. What is the initial mass of ${}^{238}\text{Pu}$ needed to meet the design criteria? (Data: The half-life of ${}^{238}\text{Pu}$ is 87.7 yr , and the alpha particle energy is $9 \times 10^{-13} \text{ J/decay}$).
8. **(15 minutes)** Consider gamma-ray spectroscopy using an intermediate-sized well-type HPGe detector where the source being counted is surrounded by the HPGe as shown below.



- a. Assume that the source is a relatively high activity ^{28}Al source sitting directly on the detector (^{28}Al emits a single 1778 keV photon per decay). Sketch what the measured energy spectrum would look like. Label and briefly explain pertinent features of your sketch.
- b. If the detector was instead extremely large what would you expect the measured energy spectrum to look like?