Ph.D. Qualifying Examination Interactions

- 1. (15 min.) A 6 MeV neutron is captured by Li-6. Derive the relevant conservation relations and then find the excitation energy of the compound nucleus. Note that the binding energy of the last neutron in Li-7 is 7.25 MeV.
- 2. (20 min.) An infinite slab of thickness t has a uniform surface source of monoenergetic photons on one face emitting $6S_a\eta^5/2\pi$ photons/cm²-sec-sr into the slab, where η is the cosine of the angle between $\hat{\Omega}$ and the inward normal, and S_a is constant. The slab has a linear attenuation coefficient of μ cm⁻¹ and is surrounded by vacuum. Using the Transport Equation find the angular flux density and the scalar flux density at the opposite face of the slab. Express your result for the scalar flux in terms of the Exponential Integrals. Recall:

$$\eta \frac{\partial \Psi}{\partial z} + \mu \Psi (z, \hat{\Omega}) = S(z, \hat{\Omega}),$$

$$E_n(x) \equiv \int_0^1 w^{n-2} e^{-x/w} dw.$$

3. (10 min.) Calculate the binding energy per nucleon for the isotope of hydrogen called tritium. Please state all your assumptions.


NOTE: The attached data table may be useful to you.

- 4. (10 min.) A thin tungsten foil with a mass of 0.1 grams is placed in an irradiation position in the TRIGA reactor at the Nuclear Science Center. The neutron fluence rate at this position is 2×10^{12} neutrons/cm² s. What is the activity of 187 W
 - (a) after 24 hours of irradiation and
 - (b) at saturation? Please state all your assumptions.

NOTE: A portion of the Chart of Nuclides is attached.

5. (10 min.) Estimate the count rate in a 10 cm³ hydrogen proportional counter operating at 7600 torr (10 atmospheres) when exposed to a monoenergetic 100 keV neutron field that produces a dose rate of 10 J/kg hr in water.

6. (20 min.) A positron (e⁺) of velocity $v_p = 2.5 \times 10^8$ m/s collides with a thin slab of ordinary matter at rest. A 0.600 MeV photon is observed to emerge from the block at an angle θ =30° relative to the path of the incident positron as in the following picture.



Assuming that in-flight annihilation with a "stationary" electron (e⁻) occurred in the block, indicate what other radiation or particle(s) were emitted and give their energies and their directions. (Explain your reasoning.)

- 7. (10 min.) A parallel beam of 1 GeV protons can be used for radiographic imaging of objects like rats, radiation detectors, and electronic instruments. The resulting images look very much like conventional x-rays, with dense areas of the object appearing light on a photographic film. Explain how these images are formed, that is, what physical interactions in the object and film are responsible for this "negative" image. Can 1 MeV protons be used to image changes in the thickness of a thin plastic film in the same way? The range of the 1 MeV proton is about twice the thickness of the plastic film. Explain why this will, or will not, work.
- 8. (15 min.) A NaI(Tl) detector inside a lead shield is exposed to Cs-137 gamma rays (662 keV) through a hole in the shield.

Describe each part of the detector-electronics system (a labeled sketch will do), and the physical processes that take place in each part of the detector-electronics system to produce a count in a particular channel on the MCA.

Sketch the expected spectrum that would be seen on the MCA, identify any features of the spectrum and name the interaction that has taken place to produce each feature.

9. (10 min.) Name two types of detectors that can be used to measure thermal neutron fluence rate in real time. Describe the advantages and disadvantages of each detector type.

Speed of light	•	, 801 . 93100000		
opera of right	J	5.77 192436 X 10" III/S		
Charge of electron	e	$1.60217733 \times 10^{-19}$ C		
Boltzmann constant	×	1.380658×10^{-23} J/K		
		$8.617385 \times 10^{-5} \text{ eV/K}$		
Faraday's constant		96485.309 C/mol		
Planck's constant	ų	$6.6260755 \times 10^{-34} \text{ J} \cdot \text{s}$		
		$4.1356692 \times 10^{-15} \text{ eV} \cdot \text{s}$		
Gravitational constant	S	$6.67259 \times 10^{-11} \text{ m}^3/\text{kg} \cdot \text{s}^2$		
Avogadro's number	×	$6.0221367 \times 10^{23} \text{ mol}^{-1}$		
Universal gas constant	`~	8.314510 J/mol·K		
Stefan-Boltzmann constant	ь	$5.6705 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$		
Rydberg constant	R	$10973731.534 \text{ m}^{-1}$		
Bohr radius	a_0	$0.529177249 \times 10^{-10}$ m		
Fine structure constant	, a	1/137.0359895		
Electron volt	eV	$1.60217733 \times 10^{-19} \text{ J}$		
Joule	—	$6.2415 \times 10^{-18} \text{ eV}$		
		10' ergs		
Erg		$10^{-7} \mathrm{J}$		
Unified mass unit		$1.66054 \times 10^{-27} \text{ kg}$	1.00000 u	931.502 MeV
Electron	ย	$9.1093897 \times 10^{-31}$ kg	$5.4857990 \times 10^{-4} \text{ u}$	0.5109990 MeV
Proton	d	$1.6726231 \times 10^{-27}$ kg	1.00727647 u	938.27231 MeV
Neutron	П	$1.6749286 \times 10^{-27} \text{ kg}$	1.00866490 u	939.56563 MeV
Deuteron	p	$3.3435856 \times 10^{-27} \text{ kg}$	2.01355320 u	1875.62883 MeV
Tritium (³ H)	Τ	$5.0073595 \times 10^{-27}$ kg	3.015500688 u	2808.94492 MeV
Alpha	σ	$6.6446618 \times 10^{-27} \text{ kg}$	4.001506178 u	3727.38025 MeV
Helium	He	$6.6464828 \times 10^{-27} \text{ kg}$	4.002603250 u	3728.43293 MeV
Comm. CONATA Durannanded Walnes of the Fundamental Physical Constants National Institute of Science and Tachnology Gaithershurg MD 1086 Internet add	Fundomontal	Physical Constants National Institute of So	sion on a Tachaclam Court	bbe terretal 1086 TM man

Source: CODATA Recommended Values of the Fundamental Physical Constants, National Institute of Science and Technology, Gaithersburg, MD, 1986. Internet address: physics.nist.gov/PhysRefData/codata86/codata86.html.

4	203	7_		
7.12 3.7 6.7 5.9 6.72.9	Hg203 4614 72792	123		
71203	Hg202 20.26 20.26	122		
T1202 12.23 d fe-vu 7439.6	Hg201 13.10	200.9702851	P(199 13.45 34.85 m 17.01 6-160, 73.01 75.3.0, 6, 20 493.7	121
10201 10364 1004.1333	Hg200	Au 199 3.14 d 3.14 d 6- 292, 25, 453 7 158 4, 208 2,	ਰੂ '	120
	Hg199	Au 198 2.17 d 2.6952 d 17 ii.e. - %2. / 2.169 / 2.	E1.372 Pt.197 1.590 b 18.3 b 17.40.7 17.10.7 17.7 19.10.7 19.1	119
	Hg198 H	F	196.966551 Pt196 25.3 04(0.04-0.6), 5	118
	Hg197	Au 196	94 P1195 94 P1195 95 17195 17185 64.29, 206 17181 06.20 174, 1781 06.20	Ir 94 1764 19.3 b 1784 19.3 b 174829, b-1331, gq.13E3
	Hg196 4.15 0 ₇ (120+3100), (39-410)	Au195	Pt194 329 329 4,0.10+1.1), (4+7) 193.9624630	192 IF193 73,834 19.33 627 19-47 34 19.33 17,744 4.4 1,744 4.4 1,744 19.4 1,744 19.4 1,744 19.4 1,744 19.4 1,744 19.4
TJ 204.3833 Thallium σ _a 3.4.13	Hg195	Au194 1.884 f. 1.49 7.338.2.293	P(193) 4.33 d 60 y (fr.135 s 1.7 y 12 to c	F 92 1.44 m 73,83 d F 51,53 17,194,1 1.316.5, 11,164,1 1.316.5, 11,164,1
81		:	Pt192 6.79 04(1+9), 110 04.40.2 mb 191.961047	1r191 37.3 54.14+ 260+660) 190.360591
	Hg 200.59 Mercury 0 ₄ ~374.8E1	Au 196.96655 Gold 5,98.7, 155Et	24 d 24 d 7538 9, 409.5. 358 9, E1019	
	80	79	Pt190 4.01 4.511 y 0.3.18 0.4.150, 70 0.4.4 mb	
			_ =	lridium G. 42E1, 28E2
			78	77

OS193	117		
OS192	116		
OS191 OS1 13.18 15.44 41 17.744 H- 141 (w); c. 17.14.90 Gr 20.5 G. 20.5 E. 314 101 00	115		
OS190 264 264 24(%+4), (22+8) 24 40.22 mb		W188 69.4 d 6-349 7207(a), 227.1, 636	114
OS189 14.1 14.1 24.(26.mb+20), 70 20.(40.01.mb	Re188 18.6 m 16.94 b 172.4 c β-2.110 β 1.15 (c. 17.130) 1.51 c 17.130 β		113
OS188 13.3 4.5.160 0.2.< CO mb	7		112
	Re186 Re18. 285.9 3.7184 62.4 17.90 91. 17.90 1.	W 85	Ta184 LT.b 7440.331. 9209 E2.87
	Ke185 37.40 57. (03+#12), 17E2 184.952955	W184 30.642 0 ₇ (.002+1.8), 14	Ta183 5.14 5.62 7.46.1.3540, 107.9. 1110
OS185 93.64 rest 6, 874 8. 880 4, 717 4,			Ta182 155 m 11434 1718 m 11434 1718 m 1718 1718 m 1718 109
	Re 186.207 Rhenium σ ₂ 89.82E1	W 183.84 Tungsten 0 ₃ -18.2.35E1	
9/	75	74	T3 T3181 180.9479 99.88 Tantalum σ _α < (μ1.70) σ _α 2066Ε1 (180.94799) 108
			73