



## Hot Topic

## KRAS: From undruggable to a druggable Cancer Target

Dipesh Uprety, Alex A. Adjei\*

Department of Oncology, Mayo Clinic, Rochester, MN, USA



## ARTICLE INFO

## Keywords:

K-RAS  
Kras  
G12C  
Cancers  
Druggable  
Undruggable

## ABSTRACT

RAS is the most frequently mutated oncogene in human cancers, with mutations in about 30% of all cancers. RAS exists in three different isoforms (*K-RAS*, *H-RAS* and *N-RAS*) with high sequence homology. *K-RAS* is the most commonly mutated RAS isoform. The Ras protein is a membrane bound protein with inherent GTPase activity and is activated by numerous extracellular stimuli, cycling between an inactive (GDP-bound) and active (GTP-bound) form. When bound to GTP, it is switched “on” and activates intracellular signaling pathways, critical for cell proliferation and angiogenesis. Mutated RAS is constitutively activated and persistently turned “on” thereby enhancing downstream signaling and leading to tumorigenesis. Various attempts to inhibit Kras in the past were unsuccessful. Recently, several small molecules (AMG510, MRTX849, JNJ-74699157, and LY3499446) have been developed to specifically target *K-RAS G12C*. Additionally, various other approaches including, SHP2, SOS1 and eIF4 inhibition, have been utilized to abrogate tumor growth in *K-RAS* mutant cells, resulting in a renewed interest in this pathway. In this review article, we provide an overview on the role of *K-RAS* in tumorigenesis, past approaches to inhibiting Kras, and current and future prospects for targeting Kras.

## Introduction

Ras is a membrane-bound protein that possesses inherent GTPase (guanosine triphosphatases) activity and is expressed in all humans [1]. When activated, it can “switch on” downstream pathways, which ultimately turn on genes that are involved in various physiological processes, including cell growth, differentiation, and survival. It was initially identified in the 1960s by Harvey and Kirsten as a retroviral oncogene when sarcomas were induced in rodents from a murine leukemogenic virus preparation; hence it's named- Kirsten rat sarcoma 2 viral oncogene homolog [2,3]. In the early 1980s, a mutated *K-RAS* oncogene was identified in a tumor biopsy of a 66-year-old male with squamous cell lung carcinoma [4]. This mutation was not identified in patient's white cells and in normal bronchial and parenchymal tissue, demonstrating the significance of somatic mutations in tumorigenesis. Subsequently, it was found that somatic *K-RAS* mutations are present in approximately 30% of all human cancers, commonly in lung, pancreas, colorectal, and cholangiocarcinoma [5–12]. In this review, we discuss Kras signaling, it's role in tumorigenesis and why this target has been considered “undruggable” historically. We also outline some strategies for targeting *K-RAS* mutant cancers by discussing promising new agents against Kras, including specific G12C inhibitors, SHP2 inhibitors, SOS1 inhibitors, and conclude by summarizing ongoing trials.

## Ras family members

*Ras structure and function*

There are two copies of *K-RAS*, namely *K-RAS1* and *K-RAS2* [13]. *K-RAS1* and *K-RAS2* are located on chromosome 6p11-12 and 12p11.1-12.1 respectively [14,15]. *K-RAS1* is a pseudo-gene [13]. Activating *K-RAS2* mutations have been identified in various human cancers. *K-RAS2* is simply referred to as *K-RAS*. The *K-RAS* gene consists of 6 exons spread over 35 kb of genomic DNA [16]. The structure of the *K-RAS* gene is depicted in Fig. 1. *K-RAS* is alternatively spliced to form *K-RAS 4A* and *K-RAS 4B*. The term *K-RAS* is generally used to indicate *K-RAS 4B* [17]. The Ras protein includes three closely related 21-kDa isoforms, H (Harvey rat sarcoma virus oncogene), N (human neuroblastoma) and K-ras (Kirsten rat sarcoma virus oncogene) [1]. Ras has three major domains: the G-domain, the C-terminal and the C-terminal CAAX box [18,19]. The G-domain, containing switch I and switch II loops, is a highly conserved domain and is responsible for GDP-GTP exchange [6,18]. The C-terminal including the CAAX box demonstrates a significant variation between RAS family members, and is required for post-translational modification [6,20]. Ras proteins bind with GDP (guanosine diphosphate) and GTP (guanosine triphosphate) with great affinity [21]. They act as “molecular switches” and cycle between the GDP-bound (inactive) and GTP-bound (active) forms. In the active

\* Corresponding author at: Mayo Clinic, 200 First St. SW, Rochester, MN, 55905.

E-mail address: [Adjei.alex@mayo.edu](mailto:Adjei.alex@mayo.edu) (A.A. Adjei).

<https://doi.org/10.1016/j.ctrv.2020.102070>

Received 18 June 2020; Received in revised form 4 July 2020; Accepted 6 July 2020

0305-7372/ © 2020 Elsevier Ltd. All rights reserved.




Fig. 1. Structure of *K-RAS* gene with associated mutations and their protein domains. Reproduced with permission from Ramakrishnan V et al. Effects of *KRAS* Gene Mutations in Gynecological Malignancies. Investigations in Gynecology Research & Womens Health.




Fig. 2. Simplified scheme of Mitogen Activation Protein Kinase activation and signaling cascade.

state, they transmit signals from the cell membrane to the nucleus, leading to activation of transcription factors which lead to the regulation of cell growth and differentiation (Fig. 2) [22].

#### Ras signaling

Ras signaling begins when a ligand binds to an upstream receptor,

such as a tyrosine kinase receptor. Almost all of the receptor tyrosine kinases are monomers [23]. A well-known pathway involves the interaction of epidermal growth factor (EGF) to its receptor (EGFR) [24]. Binding of EGF to EGFR induces dimerization of the receptor, followed by auto-phosphorylation [19,25]. The phosphorylated receptor binds to an adaptor protein Grb2 (Growth factor receptor-bound protein 2). This complex recruits son of sevenless (SOS) to the plasma membrane [23].

**Table 1**  
Prevalence of *K-RAS* mutations in human cancers.

Tumor Type	Frequency of <i>K-RAS</i> mutation
Pancreatic adenocarcinoma (61,62)	≥ 80%
Colorectal cancer (11,55–57)	30–50%
Non-small cell lung cancer (49)	
Adenocarcinoma	30%
Squamous cell carcinoma	5%
Cholangiocarcinoma (63)	
Extrahepatic	45–54%
Intrahepatic	10–15%
Multiple myeloma (64,65)	21–33%
Uterine endometrial (66, 88)	
Endometrioid	21–26%
Serous	2%
Gastric cancer (67–69)	3–13%
Testicular cancer (70–72)	9–16%
Cervical adenocarcinoma (73,74)	7–18%
Diffuse Large B-cell lymphoma (75)	1.6%
Esophageal (66, 76–78)	
Adenocarcinoma	3.8–17%
Squamous cell carcinoma	0–12%
Breast Cancer (79,80)	8–13%
Acute Myeloid Leukemia (81–83)	5–23%
Chronic Lymphocytic Leukemia (84,85)	0.7–7%
Bladder Cancer (86)	4%
Cutaneous malignant melanoma (87)	< 1%

Once recruited to the plasma membrane, SOS is capable of displacing GDP from Ras, allowing Ras-GTP interaction. Ras can also regulate SOS activity suggesting that the pathway could be bidirectional [26–28]. The binding of GTP to Ras induces changes in switch I and switch II loops of the G-domain, thereby activating Ras.

Hydrolysis of GTP to GDP inactivates Ras. Ras inherently has low GTPase activity. The intrinsic GTPase activity is stimulated further by GTPase Activating Proteins (GAPs) such as p120-GAP and NF1 (neurofibromin) [29–31]. This keeps Ras in the inactive form and prevents its persistent activation. In addition to p120 and NF1, numerous other Ras GTPases have been identified [32–34]. GAP represents a notable class of tumor suppressor genes. Normally, Ras signaling is transient. Mechanistically, inactivation of the Ras GAPs will persistently activate Ras and its effectors leading to malignant transformation. The most extensively studied tumor suppressor gene is NF1-GAP. Germline mutation of the NF1 gene predisposes to variety of tumors, including gliomas, neurofibromas, pheochromocytoma and leukemia [35–37]. Additionally, recent studies have demonstrated a high frequency of somatic NF1 mutations in a variety of sporadic tumors, including lung adenocarcinoma, leukemia, ovarian, multiple myeloma, glioblastoma and melanoma [38–40].

There are a number of effector molecules that an activated Ras can act upon including Raf, PI3K [6]. The Raf family is the best characterized Ras effector and the one with the strongest role in human cancer. Raf (Rapidly Accelerated Fibrosarcoma) protein is a serine/threonine kinase initially isolated in avian retrovirus and murine sarcoma virus [41]. It consists of three subtypes, A-raf, B-raf and Raf-1 (C-raf). Binding of GTP to Ras promotes recruitment of Raf to the cell membrane, dimerization of Raf and phosphorylation. Additionally, many factors that are not completely understood are involved in the proper activation of Raf [5,42]. Activated Raf phosphorylates MEK (Mitogen Activated Protein Kinase), which in turn, phosphorylates ERK (extracellular-signal-regulated kinase). *B-RAF* is frequently mutated in human cancers, including melanoma, thyroid malignancy and hairy cell leukemia [43–45]. When compared with *A-RAF* and *C-RAF*, *B-RAF* has a higher basal kinase activity and is easily activated by RAS [46,47].

The second best characterized Ras effector is Phosphoinositide 3'-kinase (*PI3-K*), which is activated by numerous mechanisms. One of the mechanisms involves binding of extracellular growth factor to its receptor tyrosine kinase, leading to dimerization of the receptor monomer

followed by auto-phosphorylation. Insulin Receptor Substrate-1 (IRS-1) then binds to the catalytic site of the phosphorylated dimer. Once bound to the dimer, IRS-1 serves as a binding and an activation site for PI3-K. A totally different mechanism of PI3-K activation involves direct binding of PI3-K with GTP-bound Ras. The activated PI3-K then migrates to the inner aspect of cell membrane leading to phosphorylation of phosphatidylinositol bisphosphate (PIP2) to phosphatidylinositol (3,4,5) trisphosphate (PIP3), which then activates a protein kinase AKT [48]. This ultimately activates mTOR. mTOR then activates the translation factor S6K. By binding to the larger ribosomal subunit, S6K induces translation of mRNA into protein. All the essential steps of the RAS-RAF and PI3-K signaling pathway are illustrated in Fig. 2.

## *K-RAS* mutations in human tumors

### *K-RAS* mutation subtypes

Various mutant forms of *K-RAS* are now recognized and are divided into three broad categories based on the mutated codon: G12 (mutation at codon 12), G13 (mutation at codon 13), and Q61 (mutation at codon 61).

The prevalence of *K-RAS* mutations in non-small cell lung cancer (NSCLC) is about 30% in adenocarcinoma and 5% in squamous cell carcinoma [49]. About 97% of *K-RAS* mutations in NSCLC occurs at exons 2 and 3 (G12, G13, and Q61) [50]. Also, they usually do not exist concomitantly with other sensitizing mutations, such as *EGFR*, *B-RAF* and *ALK* rearrangement [51]. *G12C* is the most common mutation subtype, accounting for about 40% of all *K-RAS* mutations followed by *G12V* [52–54].

In colorectal cancer, *K-RAS* mutations occur in about 30–50% of cases [11,55–57]. *G12D* and *G12V* are the two most common mutation subtypes [58,59]. Additionally, *K-RAS* mutations have also been identified in colorectal adenoma [60]. The prevalence of *K-RAS* mutations in pancreatic carcinoma is the highest with various studies showing the prevalence rate well above 80% with *G12D* being the most common subtype [61,62]. In cholangiocarcinoma, the prevalence of *K-RAS* mutation varies from 10% to 15% for intrahepatic cholangiocarcinoma and from 45% to 54% for extrahepatic cholangiocarcinoma [63]. *K-RAS* mutations are also found in various hematological malignancies (multiple myeloma, acute myeloid leukemia, and diffuse large B-cell lymphoma), other gastrointestinal malignancies (esophageal adenocarcinoma, gastric cancer), uterine carcinoma, and cervical adenocarcinoma [64–88]. The frequency of *K-RAS* mutations in various tumor types is summarized in Table 1.

### Predictive and prognostic value of *K-RAS* mutations

#### Prognostic value

The prognostic value of *K-RAS* mutations in various tumor types remains unclear. In NSCLC, *K-RAS* mutant NSCLC patients were considered to have a worse prognosis [89]. However, various studies have demonstrated conflicting results [10,90–92]. Mascaux and colleagues performed a systematic review of 5216 stage I-IV patients in forty-three studies from 1990 to 2003 [10]. The study demonstrated a worse survival outcome in patients with *K-RAS* mutations or p21 expression compared to those without these aberrations (HR, 1.35 [95% CI 1.16–1.56]). Moreover, the study revealed no significant impact of *K-RAS* mutations on survival for squamous histology and for the stage I and stage I-III cohorts. On the contrary, a pooled analysis utilizing the Lung Adjuvant Cisplatin Evaluation (LACE) database of 3,533 patients with stage I-III disease, demonstrated no difference in overall survival in patients with *K-RAS* mutant versus *K-RAS* wild-type NSCLC [90]. A more recent study by Pan and colleagues utilizing 41 studies from 2005 to 2015 with 13,103 patients, showed worse overall survival (HR, 1.56 [95% CI 1.39–1.76]) and disease free survival (HR, 1.57 [95% CI 1.17–2.09]) with *K-RAS* mutation in patients with early-stage resected

NSCLC [92].

In colon cancer, *K-RAS* mutations may confer poor prognosis but the data are not consistent for localized disease. While many studies demonstrated a negative impact of *K-RAS* mutations on survival [93–97], including those with localized disease [93,95,97], Roth and colleagues demonstrated that *K-RAS* mutations did not affect survival in stage II or III colon cancer [98]. Furthermore, the RASCAL-II study demonstrated that out of the 12 possible mutations on codon 12 or 13 of the *K-RAS* gene, only one mutation on codon 12, *G12V* (glycine to valine), was associated with inferior survival [99].

In pancreatic cancer, studies have demonstrated conflicting results on the prognostic value of *K-RAS* mutations on survival [100–102]. A study conducted by Bournet and colleagues involving 219 patients with locally advanced or metastatic pancreatic adenocarcinoma, demonstrated no difference in survival between *K-RAS* mutant and *K-RAS* wild-type tumors [103]. The study however showed that the *G12D* (glycine to aspartic acid) mutation had worse prognosis when compared with other mutation subtypes and *K-RAS* wild type. Additionally, coexistence of *CDKN2* aberrations and *K-RAS* mutation appeared to confer the worst prognosis [104].

In summary, the prognosis conferred by *K-RAS* mutations may differ based on the specific mutation and tumor type. This may, in part, explain the conflicting results from different studies. More studies examining mutation subtypes are needed to assess the real prognostic value of *K-RAS* mutations in tumors.

#### Predictive value

The predictive value of *K-RAS* mutations in NSCLC has been evaluated in multiple trials [105,106]. These trials have demonstrated similar response rates between the *K-RAS* mutant and *K-RAS* wild-type NSCLC. However, data suggest that *K-RAS* mutations may be a negative predictor of response to EGFR tyrosine kinase inhibitors in the minority of patients with concomitant *K-RAS* and sensitizing *EGFR* mutations [107–109]. The situation with immune checkpoint inhibitors is more complex. While there are conflicting individual studies on the outcome of *K-RAS* mutant NSCLC patients treated with PD-1/PD-L1 inhibitors, a recent meta-analysis of three prospective studies (CheckMate 057, POPLAR and OAK trial), demonstrated that patients with *K-RAS* mutant NSCLC had a superior survival compared to *K-RAS* wild type patients [110]. However, subset analyses suggest that patients with concomitant *STK11/LKB1* gene alterations may be less responsive to PD-1/PD-L1 inhibitors [111].

For colorectal cancer, *K-RAS* mutations are a major predictor of lack of response to therapy with monoclonal antibodies targeting EGFR (panitumumab, cetuximab) [112–115]. In addition, patients with *K-RAS G13D* mutations have an inferior response to chemotherapy as compared to those with other *K-RAS* mutation subtypes or those with *K-RAS* wild-type tumors [116,117]. There is a lack of data on the predictive value of *K-RAS* mutations on response to therapy as majority of pancreatic cancers harbor *K-RAS* mutations.

#### Rationale for targeting Kras in cancer therapy

Three factors support the hypothesis that Kras is a valid therapeutic target. First, *K-RAS* plays a distinct role in tumorigenesis. A study by Janseen and colleagues, in their transgenic mouse model, demonstrated that the transfection of oncogenic *K-RAS V12G* in epithelial cells of the large and small intestine led to the development of intestinal lesions including, invasive adenocarcinomas indicating a clear role of *K-RAS* in tumorigenesis [118]. Additionally, various mouse models have demonstrated the formation of frank tumors with the activation of oncogenic *K-RAS* [119,120]. Second, *K-RAS* mutant cancer cells are *K-RAS* dependent. Preclinical abrogation of mutant *K-RAS* inhibits tumor growth. A study by Collins and colleagues in a mouse model demonstrated that both primary and metastatic pancreatic adenocarcinoma lesions rely on constant Kras activity [121]. This notion is further

supported by preclinical studies in different tumor types [122,123]. Third, *K-RAS* mutant cancers represent about 30% of all human cancers as previously mentioned. Taken together, these factors make a compelling argument for targeting Kras for cancer therapy.

#### Historical approaches to Kras targeting for cancer therapy

The most successful approach to inhibiting oncogenic kinases has been the development of inhibitors that compete with ATP binding to the kinase domain. Kras utilizes GTP rather than ATP as a phosphate donor for signaling. Because of the tight binding of GTP (a thousand-fold tighter than ATP) to Kras, this approach has not been feasible, based on current technology.

Thus, a number of approaches have been utilized, as outlined below

##### Direct inhibition of Kras

Direct inhibition of Kras has been a goal of cancer drug development for several decades. SCH-53239 was the first small non-nucleotide molecule which was designed to prevent GDP to GTP nucleotide transition by binding with Ras protein and thereby preventing Ras activation [124]. Additionally, a water-soluble analog SCH-54292, which was able to bind to the switch II region of the Ras molecule, was developed [125]. However, the development of these compounds was dropped because of lack of potency.

##### Inhibition of RAS protein expression

###### Antisense oligonucleotides

Antisense oligonucleotides bind to their complementary mRNAs at a specific strand and thereby inhibit mRNA translation and ultimately the protein synthesis [126]. With this approach, a study by Gray and colleagues demonstrated a 90% reduction in Ras protein expression after targeting the 5′-flanking region of *H-RAS* in NIH-3T3 cells transformed by the *H-RAS* oncogene [127]. ISIS 2503 is an oligonucleotide that targets the 5′-untranslated region of human *H-RAS* mRNA and thereby reduces mRNA expression [128]. Cunningham *et al* conducted a phase I study utilizing this compound in 23 patients with solid tumors [129]. The compound was well tolerated. None of the patients achieved an objective response and only four patients (17%) had disease stability for 2 months or more. To evaluate the clinical activity of single agent ISIS 2503, a phase II study was conducted in sixteen patients with refractory colorectal cancer [130]. None of the patients achieved an objective response. Only one patient achieved disease stability after two cycles of treatment. Additionally, phase I and II studies were also conducted with this compound in combination with gemcitabine. In a phase I study of 27 patients with advanced cancer, the combination of ISIS 2503 and gemcitabine was well tolerated; partial response was noted in a single patient and disease stability in 5 patients [128]. This combination was further evaluated in a phase II study in 48 patients with unresectable or metastatic pancreatic adenocarcinoma [131]. At a median follow-up of 12.6 months, the study reported a 6-month survival rate of 57.5%, median survival of 6.6 months and a response rate of 10.4%. Further development of this agent was discontinued because of minimal efficacy.

##### Inhibition of K-RAS processing

###### Farnesyltransferase inhibitors

Prenylation is a post-translational addition of either a farnesyl (farnesylation) or geranylgeranyl (geranylgeranylation) moiety to the carboxyl terminus of Ras proteins that help in membrane localization. This is a rate limiting step in the post-translational modification of Kras [132]. Farnesyltransferase inhibitors were expected to block Ras farnesylation, thus preventing membrane localization and inhibiting Ras-mediated cellular proliferation. Various trials, utilizing a variety of



farnesyltransferase inhibitors, including tipifarnib (R115777), and lonafarnib (SCH 66336) were conducted in different tumor types [133–136]. A phase II trial of tipifarnib was conducted by Adjei *et al* in forty-four patients with stage IIIB or stage IV NSCLC [133]. There were no objective responses, the disease stability rate was 16%, median time to progression was 2.7 months and median OS was 7.7 months. Additionally, various phase II studies of single agent tipifarnib failed to show any objective responses in various solid tumors [136,137]. Also, a phase III study of tipifarnib failed to improve survival over best supportive care in advanced colorectal cancer [138]. A phase III study of tipifarnib in patients with advanced pancreatic adenocarcinoma randomized 688 patients to receive gemcitabine and tipifarnib or gemcitabine and placebo [139]. The study demonstrated no survival benefit with the addition of tipifarnib (median OS was 193 days and 182 days in the combination arm and gemcitabine arm respectively). In a phase III Intergroup study, 144 AML patients in remission were randomized to either receive tipifarnib or observation [140]. The study demonstrated no improvement in DFS with tipifarnib maintenance therapy.

Likewise, phase II trials of lonafarnib conducted in metastatic colorectal cancer and urothelial cancer showed no objective responses [134,135]. Additionally, a combination of lonafarnib and paclitaxel in a phase II trial of 33 patients with advanced NSCLC demonstrated a partial response of 10% and a disease stability rate of 38% [141]. Based on the results of this trial, a phase III trial of lonafarnib in combination with carboplatin-paclitaxel versus carboplatin-paclitaxel and placebo was initiated in patients with NSCLC (NCT00050336). But the study was terminated because of inadequate activity at interim analysis. These studies failed to demonstrate any benefit likely because Kras can be alternatively prenylated through geranylgeranylation [142].

Hras on the other hand is dependent solely on farnesylation for post translational modification, and the farnesyltransferase inhibitors will be expected to show activity. In support of this hypothesis, tipifarnib has recently been shown to have activity in patients with *H-RAS* mutant head and neck cancer, in a phase II study which demonstrated an overall response rate (ORR) of 56% and a median duration of response of 8.3 months [143].

#### Geranylgeranyltransferase inhibitors

Based on data indicating that Kras may be prenylated through geranylgeranylation, geranylgeranyltransferase inhibitors were evaluated in the clinic. GGTI-2418, a geranylgeranyl transferase inhibitor, was utilized in a phase I study, in 14 patients with advanced solid tumors [144]. The drug was well tolerated and no dose limiting toxicity was noted. However, no objective response was noted and the development of this class of agents was abandoned because of lack of efficacy.

#### Targeting downstream RAS effectors

##### RAF kinase inhibitors

Because of the multiple unsuccessful attempts at direct inhibition, subsequent approaches focused on the inhibition of downstream signaling of Kras.

Raf is the first protein that is phosphorylated by activated Ras in the mitogen-activated protein kinase pathway. Sorafenib (BAY 43-9006) was the first compound initially developed to specifically target Raf. It is currently approved for numerous cancers including hepatocellular carcinoma, gastrointestinal stromal tumor, renal cell carcinoma and thyroid cancer [145–147]. It is, however, not a specific or potent Raf kinase inhibitor and its antitumor activity is due to inhibition of several other receptor tyrosine kinases [148]. For instance, in pancreatic cancer, where the prevalence of *K-RAS* mutation is high, sorafenib when used in combination with chemotherapy did not demonstrate any significant clinical activity [149,150]. Subsequently, various potent inhibitors of B-Raf (dabrafenib, vemurafenib and encorafenib) were introduced and are now approved for numerous tumor types with a *B-*

*RAF* mutation (particularly *B-RAF V600E*), including melanoma, NSCLC, anaplastic thyroid and colon cancer [151–153]. Mechanistically, these agents should be effective in *K-RAS* mutant cancers as Raf is downstream of Ras, and these agents are specific inhibitors of B-Raf unlike sorafenib. However, B-raf inhibition paradoxically activates ERK signaling in wild-type *B-RAF* cells [154]. In *K-RAS* mutant cells, B-raf inhibition activates upstream proteins leading to ERK activation through an alternative pathway. One of the mechanisms by which this happens is through C-Raf activation [155]. In support of these data, the C-Raf activation cascade has been noted in *B-RAF* wild type cells and not in *B-RAF* mutant cells [156]. Currently, there are no active clinical trials utilizing single-agent *B-RAF* inhibitors in *K-RAS* mutant solid tumors.

##### MEK inhibitors

*MEK* inhibitors when used in conjunction with *B-RAF* inhibitors are superior to *B-RAF* inhibitors alone in *B-RAF* mutant cancers. Currently, three *MEK* inhibitors, trametinib, cobimetinib and binimetinib, are approved in combination with *B-RAF* inhibitors for patients with *B-RAF* mutant melanoma, NSCLC and colon cancer (along with cetuximab) [157–159].

Single agent *MEK* inhibition has shown disappointing results in various tumor types. A phase II study, of an oral *MEK* inhibitor (CI-1040), demonstrated minimal efficacy with no complete and partial responses among 67 patients with NSCLC, breast, colon and pancreatic cancer [160]. In another phase II trial, 84 patients with NSCLC, who had received one or two prior lines of treatment, were randomized to either receive pemetrexed or selumetinib (AZD6244, a *MEK* inhibitor) [161]. The study demonstrated no significant difference in progression-free survival between the two arms.

Inhibition of *MEK* induces PI3K activation leading to activation of EGFR [162]. Based on these data, various trials have utilized *MEK* inhibitors in combination with EGFR inhibitors. Phase II studies of selumetinib combined with erlotinib failed to show activity in previously treated NSCLC [163] and pancreatic carcinoma [164].

The combination of *MEK* inhibitors and chemotherapy has also been evaluated for *K-RAS* mutant NSCLC. A number of phase II studies were performed [165–168] culminating in the phase III SELECT-1 trial which randomized patients with *K-RAS* mutant advanced NSCLC with disease progression after first-line treatment to either selumetinib plus docetaxel or placebo plus docetaxel [169]. Median PFS was 3.9 months with selumetinib plus docetaxel therapy versus 2.8 months in placebo plus docetaxel (HR, 0.93 [95% CI 0.77–1.12];  $P = 0.44$ ). In summary, *MEK* inhibition either alone or in combination is not an effective therapy for *K-RAS* mutant cancers.

##### ERK inhibitors

Since ERK is the final downstream kinase of the MAP Kinase pathway, it has been hypothesized that *ERK* inhibition may be effective in *K-RAS* mutant tumors. This hypothesis is supported by preclinical data [170]. Various *ERK* inhibitors, including, LY3214496, BVD-523, MK-8353 and KO-947, are in early phase of clinical development either alone or in combination (NCT02857270, NCT-01781429, NCT02972034, NCT03745989, NCT03051035). The dose escalation portion of a phase I trial of LY3214496 in patients with *K-RAS*, *N-RAS* or *B-RAF* mutant advanced or metastatic cancer has been reported [171]. No concerning toxicities were noted. The study is now in the second phase, where LY3214496 is utilized either alone or in combination with abemaciclib or nab-paclitaxel plus gemcitabine in various tumor types (NCT02857270).

##### CDK4/6 inhibitors

Abemaciclib is a cyclin-dependent kinase (CDK 4/6) inhibitor currently approved in combination with hormonal therapy for patients with advanced or metastatic hormone receptor positive and HER-

**Table 2**

Completed phase II or phase III studies of downstream Kras signaling inhibition in tumors with high prevalence of K-RAS mutations.

Study	Tumor type	Study design/ Phase	Sample Size	Intervention	ORR (%)	Median PFS (months)	Median OS (months)
GISCAD study <i>Cascinuet al</i> (151)	Pancreatic adenocarcinoma	Open-label Randomized Phase II	114	Gemcitabine/cisplatin plus sorafenib versus Gemcitabine/ cisplatin	3.4 vs. 3.6	4.3 vs. 4.5	7.5 vs. 8.3
<i>Kindler et al</i> (152)	Pancreatic adenocarcinoma	Open-label Single arm Phase II	17	Gemcitabine plus sorafenib	0	3.2	4.0
<i>Rinehart et al</i> (162)	CRC, NSCLC, breast, pancreatic cancer	Open-label Single arm Phase II	67	CI-1040	0	4.4*	NRe
<i>Hainsworth et al</i> (163)	NSCLC	Open-label Randomized Phase II	84	Selumetinib (AZD6244) versus pemetrexed	5 vs. 5	2.2 vs. 3.0	NRe
<i>Carter et al</i> (165)	NSCLC	Randomized Phase II	79	K-RAS wild-type: erlotinib versus erlotinib plus selumetinib K-RAS mutant: selumetinib versus selumetinib plus erlotinib	5 vs. 12 0 vs. 10	2.4 vs. 2.1 4.0 vs. 2.3	6.3 vs. 12.9 10.5 vs. 21.8
<i>Blumenschein et al</i> (176)	NSCLC K-RAS, NRAS, BRAF, or MEK1 mutant	Open-label Randomized Phase II	134	Trametinib versus docetaxel	12 vs. 12	2.8 vs. 2.6	8 vs. NR
SELECT-1 <i>Janne et al</i> (171)	NSCLC K-RAS mutant	Randomized Double Blind Phase III	510	Selumetinib plus docetaxel versus placebo plus docetaxel	20.1 vs. 13.7	3.9 vs. 2.8	8.7 vs. 7.9
<i>Ko et al</i> (166)	PDAC	Non-randomized Single arm Phase II	46	Selumetinib plus erlotinib	0	1.9	7.3
<i>Infante et al</i> (170)	Pancreatic adenocarcinoma	Double-blind Randomized Phase II	160	Trametinib plus gemcitabine versus placebo plus gemcitabine	22 vs. 18	3.8 vs. 3.5	8.4 vs. 6.7
JUNIPER <i>Goldman et al</i> (175)	NSCLC K-RAS mutant	Open-label Randomized Phase III	453	Abemaciclib versus erlotinib	8.9 vs. 2.7	3.6 vs. 1.9	7.4 vs. 7.8

ORR, Objective Response Rate; PFS, Progression Free Survival; OS, Overall Survival; CRC, colorectal carcinoma; NSCLC, non-small cell lung cancer; NRe, not reported; \* median duration of stable disease; NR, Not Reached; PDAC, pancreatic ductal adenocarcinoma.

negative breast cancer. In a mouse tumor model that recapitulates human NSCLC, *Puyol* and colleagues demonstrated that CDK4 inhibition can induce selective death of K-RAS mutant cancer cells [172]. In a phase III open-label trial, patients with stage IV, K-RAS mutant NSCLC after having disease progression on platinum-doublet were randomized to abemaciclib or erlotinib. ORR was 8.9% in the abemaciclib arm versus 2.7% in the erlotinib arm ( $P = 0.01$ ) [173]. Likewise, median PFS was 3.6 months with abemaciclib versus 1.9 months with erlotinib (HR, 0.58; 95% CI 0.47–0.72). However, despite having a better response rate and PFS, abemaciclib did not improve overall survival (median OS with abemaciclib was 7.4 months versus 7.8 months with erlotinib).

The results of the above mentioned studies are summarized in Table 2 [149,150,160,161,163,164,168,169,173,174]. In addition, various ongoing trials that have targeted downstream Kras effectors or utilized indirect Kras inhibition approaches are summarized in table 3.

### Covalent Kras G12C inhibitors

Kras has been considered “undruggable” despite decades of extensive attempts to develop an effective anti-Ras therapy, as described above. Recent studies have identified small molecules that can selectively target and inactivate the K-RAS G12C mutant variant [175–180]. K-RAS G12C results from a missense mutation (glycine-to-cysteine substitution) at codon 12. This leads to impairment of GAP mediated hydrolysis of GTP to GDP, thereby locking the Kras protein in a hyperexcitable state.

K-RAS G12C is the most common mutant variant in NSCLC accounting for about 40% of all K-RAS mutant tumors and about 13% of all lung adenocarcinoma [52,54,181]. Additionally, it is present in about 3% of colorectal cancer cases and a small subset of patients with pancreatic, endometrial and urothelial cancers [66,182]. The frequency

of different K-RAS mutations in various tumor types is shown in Fig. 3 [183].

Ostrem and colleagues developed a series of compounds that could target the mutant Kras G12C protein by covalently binding to the mutant cysteine residue [175]. With this approach, they were able to selectively target the mutant cells and spare the normal ones. Additionally, they found that these inhibitors were binding to a new allosteric binding pocket, the switch-II pocket (S-IIP). This pocket extends from the mutant cysteine residue into a pocket comprising mainly of the switch II region. By targeting this specific pocket, these compounds displace glycine 60 towards the switch I region leading to conformational disruption of GTP bound Ras and thereby preventing further downstream signaling. However, the initial lead compound developed by Ostrem and colleagues (compound 12) had poor pharmacologic properties [176,177].

Consequently, ARS853, which had more than 600-fold improved engagement with Kras G12C over compound 12, was discovered [176,177]. Two studies, by *Lito et al* and *Patricelli et al*, demonstrated reduction in GTP-bound Kras levels in K-RAS G12C mutant cancer cell lines after treatment with ARS853 [176,177]. Additionally, they also showed that ARS833 bound preferentially to the GDP-bound state of Kras G12C. Since this compound was selectively inhibiting GDP-bound Kras, there was a significant concern about its efficacy in vivo. Consequently, *Janes* and colleagues identified a compound, ARS-1260, which selectively targets the switch II pocket and also inhibits Kras in the GTP-bound state [178]. They demonstrated that ARS-1260 covalently inhibits Kras G12C activity in vitro, and exhibited antitumor activity in subcutaneous xenograft models bearing K-RAS G12C but not G12V mutations.

Subsequently, *Canon* and colleagues demonstrated the activity of AMG 510 in K-RAS G12C mouse xenografts [179]. Similar preclinical studies of another Kras G12C inhibitor, MRTX849 have been published

**Table 3**  
Ongoing trials targeting Kras downstream signaling in *K-RAS* mutant solid tumors.

Site of action	Study/Clinical Trial Identifier	Tumor Type	Mutation profile	Phase/Study Design	Intervention	Recruitment status	Estimated Enrollment
PI3K	NCT04073680	Solid tumor	PIK3CA or K-RAS	Ib/II Open-label	Serabelisib and Canagliflozin	Not yet recruiting	60
RAF	NCT02974725	NSCLC, melanoma	K-RAS or BRAF (NSCLC), N-RAS (melanoma)	Ib Open-label	LXH254 with LTT462 or trametinib or ribociclib	Recruiting	195
RAF	NCT04249843	Solid tumor	B-RAF, K-RAS, N-RAS	Ia/Ib Open-label	BGB-3245	Recruiting	69
RAF MEK	NCT03681483	NSCLC	K-RAS	I Open-label	RO5126766	Recruiting	31
MEK	NCT03704688	NSCLC	K-RAS	I Open-label	Trametinib and ponatinib	Recruiting	37
MEK	NCT04132505	Pancreatic adenocarcinoma	K-RAS	I Open-label	Binimetinib and hydroxychloroquine	Recruiting	39
MEK	NCT03299088	NSCLC	K-RAS	Ib Open-label	Trametinib and pembrolizumab	Recruiting	42
MEK	NCT02613650	Colorectal	H-RAS, N-RAS or K-RAS	Ib Open-label	MEK162 and mFOLFIRI	Recruiting	30
MEK	NCT03981614	Colorectal	K-RAS or N-RAS	II Randomized Open-label	Binimetinib plus palbociclib versus TAS-102 (trifluridine and tipiracil)	Recruiting	112
MEK	SELECT-1 NCT01933932	NSCLC	K-RAS	III Randomized Double-blind	Selumetinib plus docetaxel versus docetaxel plus placebo	Active, not recruiting	510
MEK	M14LTK NCT02230553	NSCLC	K-RAS mutant and PIK3CA wild-type	I/II Open-label	Trametinib and lapatinib	Recruiting	30
MEK	NCT03990077	NSCLC	K-RAS	I Open-label	HL-085 and docetaxel	Not yet recruiting	27
MEK	M14AFS NCT02450656	NSCLC	K-RAS mutant and PIK3CA wild-type	I/II Randomized Open-label	Selumetinib plus afatinib versus docetaxel	Recruiting	320
MEK	MEKiAUTO NCT04214418	Pancreatic adenocarcinoma colorectal	K-RAS	I/II Open-label	Cobimetinib, hydroxychloroquine and atezolizumab	Not yet recruiting	175
MEK	NCT03170206	NSCLC	K-RAS	I/II Open-label	Binimetinib and palbociclib	Recruiting	72
MEK	NCT02079740	Solid tumor	K-RAS or N-RAS	Ib/II Open-label	Trametinib and navitoclax	Recruiting	130
MEK	NCT03637491	Solid tumor	K-RAS or N-RAS	Ib/II Open-label Randomized	Binimetinib and avelumab or binimetinib, avelumab and talazoparib or binimetinib and talazoparib	Recruiting	122
ERK	NCT02857270	Melanoma, NSCLC, colorectal, pancreatic adenocarcinoma	BRAF or RAS NSCLC, BRAF or NRAS melanoma, BRAF colorectal	I Open-label	LY3214996 alone and in combination with abemaciclib, nab-paclitaxel and gemcitabine or cetuximab and encorafenib	Recruiting	272
ERK	NCT04145297	GI malignancy	KRAS, NRAS, HRAS, BRAF non-V600, MEK, ERK	I Open-label	Ulixertinib (BVD-523) and hydroxychloroquine	Recruiting	12
ERK	NCT03051035	Solid tumor	BRAF, KRAS, NRAS or HRAS in non-squamous histology	I Open-label	KO-947 alone	Active but not recruiting	100
ERK	NCT03415126	Solid tumor	BRAF mutant melanoma, N-RAS or H-RAS mutant solid tumors, K-RAS mutant CRC, K-RAS mutant NSCLC	I Open-label	ASN007	Active but not recruiting	49
mTOR	NCT03520842	Non-squamous NSCLC	K-RAS	II Open-label	Regorafenib and methotrexate	Recruiting	18

NSCLC, non-small cell lung cancer.

[180].

A phase I study of AMG 510 was presented at the European Society for Medical Oncology (ESMO) annual meeting and at the World Lung Cancer Conference in 2019, by Govindan and colleagues [184]. A total of 76 patients with *K-RAS G12C* mutant solid tumors were enrolled in the study. There was no dose limiting toxicity. Most of the patients (34.2%) had grade 1 or grade 2 treatment-related adverse events. 6 patients had grade 3 adverse events, which included anemia and diarrhea. The recommended phase II dose was 960 mg once daily. Among NSCLC cohort (n = 23), the ORR was 48% and disease control rate (DCR) was 96%. In the colorectal cancer cohort (n = 29), the ORR was

3% and the DCR was 79%. There are two other phase I trials (NCT03600883, NCT04185883), which are actively recruiting patients with *K-RAS G12C* mutant solid malignancies. These trials will also assess the safety and feasibility of various therapeutic agents in combination with AMG510, including a PD-1 inhibitor, MEK inhibitor, a SHP2 allosteric inhibitor, and a pan-ErbB tyrosine kinase inhibitor. Additionally, a phase III trial is scheduled to start accrual this summer in patients with previously treated locally advanced and unresectable or metastatic *K-RAS G12C* mutant NSCLC with randomization to AMG510 or docetaxel (NCT04303780). The clinical activity of AMG 510 in colorectal cancer is minimal to modest compared to the activity in NSCLC,




Fig. 3. K-RAS mutation frequency in different tumor types.

suggesting that signaling networks in colorectal cancer are different from NSCLC. As an example, Braf inhibition in *BRAF V600E* mutation is much more effective in NSCLC compared to colorectal cancer, where bypass signaling in EGFR abrogates the effect of Braf inhibition. Thus, concurrent inhibition of EGFR is needed to achieve impressive responses after Braf inhibition. This same mechanism seems to be present in *K-RAS G12C* mutant colorectal cancer, as demonstrated by Amodio and colleagues [185].

Studies with another Kras G12C inhibitor are also ongoing. A phase 1/2 multiple expansion study of MRTX849 is currently accruing patients (NCT03785249). In this trial, patients with advanced, unresectable or metastatic solid tumors with a *K-RAS G12C* mutation will be enrolled to access the safety, pharmacokinetics, tolerability and clinical activity of MRTX849. This trial will also evaluate the safety of the combination of MRTX849 with other therapeutic agents, including, a PD-1 inhibitor in patients with NSCLC and cetuximab in patients with colorectal cancer. Another phase 1/2 study will be opening in the near future utilizing a combination of MRTX849 and TNO155 in patients with KRAS G12C mutant cancer (NCT04330664). TNO155 is a SHP2 inhibitor and will be discussed in detail below. Two other *K-RAS G12C* inhibitors, ARS-3248/JNJ-74699157, and LY3499446 are under investigation (NCT04006301, NCT04165031). Table 4 summarizes all the active trials in *K-RAS* mutant solid tumors which utilize novel direct inhibitors of Kras.

## PAN K-RAS inhibitors

### SOS1 inhibitors

BI-3406 is an orally bioavailable drug designed to inhibit the son of sevenless 1 (SOS1) protein. Hofmann and colleagues have demonstrated that this Boehringer-Ingelheim drug only inhibits SOS1, and not SOS2 [186]. They further demonstrated that in *K-RAS*-mutant cancer, including G12 and G13. By inhibiting SOS1, BI-3406 reduced GTP-KRAS levels thereby restricting tumor cell proliferation. BI 1701963, which is a BI-3406 analog, is in phase I trials, either alone or in combination with Trametinib in patients with *K-RAS* mutant solid tumors (NCT04111458).

## SHP2 inhibitors

SHP2, a protein tyrosine phosphatase (PTPN11), relays stimulatory signals from various membrane receptor tyrosine kinases to the MAPK kinase signaling pathway [187]. Chen and colleagues initially developed SHP099, a selective and orally bioavailable allosteric inhibitor of SHP2, and demonstrated its antitumor activity in receptor tyrosine kinase-driven cancers in patient derived tumor xenograft models [188]. Later, Mainardi and colleagues demonstrated an importance of SHP2 inhibition in controlling *K-RAS* mutant tumor growth by MEK inhibition [189]. They demonstrated that MEK inhibition can reduce phosphorylated ERK in cell lines of three tumor types (NSCLC, pancreatic cancer and colon cancer). Furthermore, they found that ERK levels slowly started rising along with a rise in SHP2 levels suggesting the activation of a feedback loop involving receptor tyrosine kinase. Additionally, when they simultaneously blocked SHP2 and MEK, they found a strong synergy between a SHP2 inhibitor and a MEK inhibitor in all three cell lines, and the strongest effect was observed in NSCLC cell lines. In addition, they demonstrated that the PTPN11-knockout cells demonstrated lower baseline RAS-GTP levels and had an increased sensitivity to MEK inhibitor. Based on these preclinical data, it is reasonable to utilize SHP inhibitor in *K-RAS* mutant tumor. There are two SHP 2 inhibitors, namely, RMC 4630 and TNO155, in early phase of development, (NCT03634982, NCT03989115, NCT04000529, NCT03114319, NCT04330664).

## Transcription regulator eIF4 inhibitors

Protein synthesis is catalyzed by eukaryotic translational initiation factor 4 (eIF4) which is responsible for recruitment of the 5'-untranslated segment of the mRNA to the ribosomal subunit [190]. eIF4A, a component of eIF4 complex, is an ATPase/RNA helicases and its specific role in this process is mRNA unwinding to facilitate ribosome binding [190]. Since this protein complex is an essential component of the translation initiation of multiple oncogenic pathways, including *K-RAS*, targeting this protein in *K-RAS* mutant cancer cases can potentially control tumor growth and proliferation. eFT226 is a first in class selective inhibitor of eIF4A. Thompson and colleagues have demonstrated antitumor activity of this compound in a preclinical study in B-



**Table 4**  
Ongoing clinical trials of novel Kras inhibitors in K-RAS mutant solid tumors.

Site of action	Study/Clinical Trial Identifier	Tumor Type	Mutation profile	Phase/Study Design	Intervention	Recruitment status	Estimated Enrollment
K-RAS G12C	CodeBreak 101 NCT04185883	Solid tumor	K-RAS G12C	Ib	AMG 510 alone and in combination with MEK, PD-1, SHP2 or ErbB inhibitor	Recruiting	250
K-RAS G12C	CodeBreak 200 NCT04303780	NSCLC	K-RAS G12C	III Randomized	AMG 510 plus Docetaxel versus Docetaxel	Not yet recruiting	650
K-RAS G12C	CodeBreak 100 NCT04303780	Solid tumor	K-RAS G12C	Open-label I/II	AMG 510 alone and in combination with PD-1 inhibitor	Recruiting	533
K-RAS G12C	NCT03785249	Solid tumor	K-RAS G12C	Open-label I/II	MRTX849 alone	Recruiting	200
K-RAS G12C	NCT04330664	Solid tumor	K-RAS G12C	Open-label I/II	MRTX849 in combination with TNO155	Recruiting	148
K-RAS G12C	NCT04165031	Solid tumor	K-RAS G12C	Open-label I/II	LY3499446 alone or in combination with abemaciclib, cetuximab, erlotinib, docetaxel	Recruiting	230
K-RAS G12C	NCT04006301	Solid Tumor	K-RAS G12C	I	JNJ-74699157 alone	Recruiting	140
SOS1	NCT04111458	Solid tumor	K-RAS	Open-label I	BI 1,701,963 alone and in combination with trametinib	Recruiting	140
SHP2	NCT03634982	Solid tumor	RTK, K-RAS G12, BRAF class 3, NF1 LOF	Open-label I	RMC-4630 alone	Recruiting	240
SHP2	NCT03989115	Solid tumor	K-RAS, BRAF class 3, NF1 LOF	Open-label Ib/II	RMC-4630 in combination with cobimetinib	Recruiting	144
SHP2	NCT03114319	Solid tumor	EGFR NSCLC, K-RAS G12C NSCLC, CRC, Esophageal SCC, HN SCC, RAS/RAF wild-type other solid tumor	Open-label I	TNO155 alone	Recruiting	135
SHP2	NCT04000529	NSCLC, CRC, HNSCC	K-RAS G12C NSCLC, K-RAS wild-type NSCLC, K-RAS codon 12, 13, or 61 for CRC (dose escalation)	Ib Open-label	TNO155 in combination with spartalizumab or ribociclib	Recruiting	126
mRNA based vaccine	NCT03948763	NSCLC, CRC, pancreatic adenocarcinoma	K-RAS G12D, G12V, G13D, G12C	I	mRNA-5671/V941 alone and in combination with pembrolizumab	Recruiting	100
eIF4	NCT04092673	Solid tumor	HER2, ERBB3, FGFR1, FGFR2, K-RAS	Open-label I/II Open-label	eFT226 (Zotatifin) alone	Recruiting	45

CRC, colorectal carcinoma; NSCLC, non-small cell lung cancer.




Fig. 4. Various drugs targeting at different levels on Kras pathway.

cell lymphoma where there is a PI3K/AKT/mTOR pathway aberrancy [191]. Additionally, Thompson and colleagues have demonstrated in vivo tumor growth inhibition in solid tumor xenograft models with FGFR1/2 or HER2 amplifications, including NSCLC, breast and colorectal cancers [192]. A phase I trial of eFT 226 (zotatifin) is currently recruiting patients with HER2, ERBB3, FGFR1, FGFR2 and K-RAS mutant solid tumors (NCT 04092673). Various drugs targeting at different levels on Kras pathway is depicted in Fig. 4.

### Other agents in late preclinical development

Recently, mRNA-based vaccination is being utilized to investigate specific immune responses against cancer cells. Mutanome is a distinct set of somatic mutations unique to an individual's tumor. As majority of these mutations are unique to each individual, Sahin *et al* investigated the concept of individualized mutanome vaccines by implementing an RNA-based neo-epitope approach in patients with stage III or IV melanoma [193]. After identifying non-synonymous mutations in 13 patients, an RNA vaccine was engineered encoding 10 selected mutations per patient, which was then injected intranodally. Following vaccination, all patients developed T-cell responses and two of the five patients with metastatic disease achieved an objective response. This study unlocked a novel path for a more personalized treatment and has drawn significant attention. A phase I trial of mRNA-5671/V941 (that encodes antigen for K-RAS G12D, G12V, G12C and G13D) as monotherapy and in combination with pembrolizumab in patients with solid tumors with four prevalent K-RAS mutations is currently underway (NCT03948763). The mRNA-5671/V941 vaccine is intended to target majority of the K-RAS mutations that occur in solid tumors.

Additionally, a novel short inhibitory peptide, KRpep-2d, is recently identified using a T7 phage display technique. KRpep-2d is a 19-mer cyclic peptide which is able to non-covalently and selectively inhibit Kras G12D activity with high potency [194,195]. It acts as an allosteric inhibitor by binding near the switch II pocket [196]. This molecule is still in its infancy but is likely to enter clinical trial in near future for K-RAS G12D mutant tumors.

### Conclusion

While K-RAS has been seen as an attractive target for cancer

therapy, all approaches taken to inhibit K-RAS either directly or indirectly through inhibiting post translational modification or downstream signaling to date have been ineffective. Advances in genomics and molecular biology, however, have for the first time suggested that direct inhibition of K-RAS may be possible. An identification of a targetable binding pocket (S-IIP) in K-RAS G12C recently resulted in a renewed interest in targeting K-RAS via G12C inhibition. The covalent Kras G12C inhibitors have provided the first clinical evidence of the ability to inhibit a class of K-RAS mutant tumors. This initial success has rekindled interest in Kras inhibition, with a number of other approaches including SHP2, SOS1 and eIF4 inhibition being tested in the clinic. These newer approaches, if successful would abrogate the activity of all mutant isoforms of K-RAS. Thus, we are for the first time at the cusp of successfully drugging this hitherto undruggable target.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### References

- [1] Barbacid M. ras genes. *Annu Rev Biochem* 1987;56:779–827.
- [2] Harvey JJ. An unidentified virus which causes the rapid production of tumours in mice. *Nature* 1964;204:1104–5.
- [3] Kirsten WH, Mayer LA. Morphologic responses to a murine erythroblastosis virus. *J Natl Cancer Inst* 1967;39(2):311–35.
- [4] Santos E, Martin-Zanca D, Reddy EP, Pierotti MA, Della Porta G, Barbacid M. Malignant activation of a K-ras oncogene in lung carcinoma but not in normal tissue of the same patient. *Science* 1984;223(4637):661–4.
- [5] Fernandez-Medarde A, Santos E. Ras in cancer and developmental diseases. *Genes Cancer* 2011;2(3):344–58.
- [6] Friday BB, Adjei AA. K-ras as a target for cancer therapy. *BBA* 2005;1756(2):127–44.
- [7] Wang JY, Lian ST, Chen YF, Yang YC, Chen LT, Lee KT, *et al*. Unique K-ras mutational pattern in pancreatic adenocarcinoma from Taiwanese patients. *Cancer Lett* 2002;180(2):153–8.
- [8] Shibata D, Almoguera C, Forrester K, Dunitz J, Martin SE, Cosgrove MM, *et al*. Detection of c-k-ras mutations in fine needle aspirates from human pancreatic adenocarcinomas. *Cancer Res* 1990;50(4):1279–83.
- [9] Pao W, Girard N. New driver mutations in non-small-cell lung cancer. *Lancet Oncol* 2011;12(2):175–80.
- [10] Mascaux C, Iannino N, Martin B, Paesmans M, Berghmans T, Dusart M, *et al*. The

- role of RAS oncogene in survival of patients with lung cancer: a systematic review of the literature with meta-analysis. *Br J Cancer* 2005;92(1):131–9.
- [11] Peeters M, Kafatos G, Taylor A, Gastanaga VM, Oliner KS, Hechmati G, Terwey J-H, van Krieken JH. Prevalence of RAS mutations and individual variation patterns among patients with metastatic colorectal cancer: a pooled analysis of randomised controlled trials. *Eur J Cancer* 2015;51(13):1704–13. <https://doi.org/10.1016/j.ejca.2015.05.017>.
  - [12] Munoz-Maldonado C, Zimmer Y, Medova M. A comparative analysis of individual RAS mutations in cancer biology. *Front Oncol* 2019;9:1088.
  - [13] Jancik S, Drabek J, Radzioch D, Hajdich M. Clinical relevance of KRAS in human cancers. *J Biomed Biotechnol* 2010;2010:150960.
  - [14] McBride OW, Swan DC, Tronick SR, Gol R, Klimanis D, Moore DE, et al. Regional chromosomal localization of N-ras, K-ras-1, K-ras-2 and myb oncogenes in human cells. *Nucleic Acids Res* 1983;11(23):8221–36.
  - [15] Popescu NC, Amsbaugh SC, DiPaolo JA, Tronick SR, Aaronson SA, Swan DC. Chromosomal localization of three human ras genes by in situ molecular hybridization. *Somat Cell Mol Genet* 1985;11(2):149–55.
  - [16] Sameer AS. Colorectal cancer: molecular mutations and polymorphisms. *Front Oncol* 2013;3:114.
  - [17] Chen Z, Otto JC, Berge MO, Young SG, Casey PJ. The C-terminal polylysine region and methylation of K-Ras are critical for the interaction between K-Ras and microtubules. *J Biol Chem* 2000;275(52):41251–7.
  - [18] Vogler O, Barcelo JM, Ribas C, Escriba PV. Membrane interactions of G proteins and other related proteins. *BBA* 2008;1778(7–8):1640–52.
  - [19] Lemmon MA, Schlessinger J. Regulation of signal transduction and signal diversity by receptor oligomerization. *Trends Biochem Sci* 1994;19(11):459–63.
  - [20] Gao J, Liao J, Yang GY. CAAX-box protein, prenylation process and carcinogenesis. *Am J Transl Res* 2009;1(3):312–25.
  - [21] Lowy DR, Willumsen BM. Function and regulation of ras. *Annu Rev Biochem* 1993;62:851–91.
  - [22] Adjei AA. Blocking oncogenic Ras signaling for cancer therapy. *J Natl Cancer Inst* 2001;93(14):1062–74.
  - [23] Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 2000;103(2):211–25.
  - [24] Liebmann C. Regulation of MAP kinase activity by peptide receptor signalling pathway: paradigms of multiplicity. *Cell Signal* 2001;13(11):777–85.
  - [25] Heldin CH. Dimerization of cell surface receptors in signal transduction. *Cell* 1995;80(2):213–23.
  - [26] Iversen L, Tu HL, Lin WC, Christensen SM, Abel SM, Iwig J, et al. Molecular kinetics. Ras activation by SOS: allosteric regulation by altered fluctuation dynamics. *Science* 2014;345(6192):50–4.
  - [27] Margarit SM, Sondermann H, Hall BE, Nagar B, Hoelz A, Pirruccello M, et al. Structural evidence for feedback activation by Ras.GTP of the Ras-specific nucleotide exchange factor SOS. *Cell* 2003;112(5):685–95.
  - [28] Sondermann H, Soisson SM, Boykevich S, Yang SS, Bar-Sagi D, Kuriyan J. Structural analysis of autoinhibition in the Ras activator Son of sevenless. *Cell* 2004;119(3):393–405.
  - [29] Campbell PM, Der CJ. Oncogenic Ras and its role in tumor cell invasion and metastasis. *Semin Cancer Biol* 2004;14(2):105–14.
  - [30] Drugan JK, Rogers-Graham K, Gilmer T, Campbell S, Clark GJ. The Ras/p120 GTPase-activating protein (GAP) interaction is regulated by the p120 GAP pleckstrin homology domain. *Jo Boil Chem* 2000;275(45):35021–7.
  - [31] Paminsinlatham P, Hadj-Slimane R, Lepelletier Y, Allain B, Toccafondi M, Garbay C, et al. p120-Ras GTPase activating protein (RasGAP): a multi-interacting protein in downstream signaling. *Biochimie* 2009;91(3):320–8.
  - [32] Maekawa M, Li S, Iwamatsu A, Morishita T, Yokota K, Imai Y, et al. A novel mammalian Ras GTPase-activating protein which has phospholipid-binding and Btk homology regions. *Mol Cell Biol* 1994;14(10):6879–85.
  - [33] Jin H, Wang X, Ying J, Wong AH, Cui Y, Srivastava G, et al. Epigenetic silencing of a Ca(2+)-regulated Ras GTPase-activating protein RASAL defines a new mechanism of Ras activation in human cancers. In: *Proceedings of the national academy of sciences of the United States of America*. 2007;104(30):12353–8.
  - [34] Kim JH, Liao D, Lau LF, Hagan RL. SynGAP: a synaptic RasGAP that associates with the PSD-95/SAP90 protein family. *Neuron* 1998;20(4):683–91.
  - [35] Ledbetter DH, Rich DC, O'Connell P, Leppert M, Carey JC. Precise localization of NF1 to 17q11.2 by balanced translocation. *Am J Hum Genet* 1989;44(1):20–4.
  - [36] Tucker T, Riccardi VM, Sutcliffe M, Vielkind J, Wechsler J, Wolkenstein P, et al. Different patterns of mast cells distinguish diffuse from encapsulated neurofibromas in patients with neurofibromatosis 1. *J Histochem Cytochem: Off J Histochem Soc* 2011;59(6):584–90.
  - [37] Sorensen SA, Mulvihill JJ, Nielsen A. Long-term follow-up of von Recklinghausen neurofibromatosis. Survival and malignant neoplasms. *New Engl J Med* 1986;314(16):1010–5.
  - [38] Ding L, Getz G, Wheeler DA, Mardis ER, McLellan MD, Cibulskis K, et al. Somatic mutations affect key pathways in lung adenocarcinoma. *Nature* 2008;455(7216):1069–75.
  - [39] Boudry-Labis E, Roche-Lestienne C, Nibourel O, Boissel N, Terre C, Perot C, et al. Neurofibromatosis-1 gene deletions and mutations in de novo adult acute myeloid leukemia. *Am J Hematol* 2013;88(4):306–11.
  - [40] Philpott C, Tovell H, Frayling IM, Cooper DN, Upadhyaya M. The NF1 somatic mutational landscape in sporadic human cancers. *Human Genomics* 2017;11(1):13.
  - [41] Zebisch A, Troppmair J. Back to the roots: the remarkable RAF oncogene story. *Cell Mole Life Sci: CMLS* 2006;63(11):1314–30.
  - [42] Stokoe D, Macdonald SG, Cadwalader K, Symons M, Hancock JF. Activation of Raf as a result of recruitment to the plasma membrane. *Sci (New York, NY)* 1994;264(5164):1463–7.
  - [43] Long GV, Menzies AM, Nagrial AM, Haydu LE, Hamilton AL, Mann GJ, et al. Prognostic and clinicopathologic associations of oncogenic BRAF in metastatic melanoma. *J Clin Oncol: Off J Am Soc Clin Oncol* 2011;29(10):1239–46.
  - [44] Kimura ET, Nikiforova MN, Zhu Z, Knauf JA, Nikiforov YE, Fagin JA. High prevalence of BRAF mutations in thyroid cancer: genetic evidence for constitutive activation of the RET/PTC-RAS-BRAF signaling pathway in papillary thyroid carcinoma. *Cancer Res* 2003;63(7):1454–7.
  - [45] Tiacci E, Trifonov V, Schiavoni G, Holmes A, Kern W, Martelli MP, et al. BRAF mutations in hairy-cell leukemia. *New Engl J Med* 2011;364(24):2305–15.
  - [46] Wellbrock C, Karasides M, Marais R. The RAF proteins take centre stage. *Nat Rev Mol Cell Biol* 2004;5(11):875–85.
  - [47] Emuss V, Garnett M, Mason C, Marais R. Mutations of C-RAF are rare in human cancer because C-RAF has a low basal kinase activity compared with B-RAF. *Cancer Res* 2005;65(21):9719–26.
  - [48] Castellano E, Downward J. RAS interaction with PI3K: more than just another effector pathway. *Genes Cancer* 2011;2(3):261–74.
  - [49] Roberts PJ, Stinchcombe TE, Der CJ, Socinski MA. Personalized medicine in non-small-cell lung cancer: is KRAS a useful marker in selecting patients for epidermal growth factor receptor targeted therapy? *J Clin Oncol: Off J Am Soc Clin Oncol* 2010;28(31):4769–77.
  - [50] Wood K, Hensing T, Malik R, Sargia R. Prognostic and predictive value in KRAS in non-small-cell lung cancer: a review. *JAMA Oncol* 2016;2(6):805–12.
  - [51] Gainor JF, Varghese AM, Ou SH, Kabraji S, Awad MM, Katayama R, et al. ALK rearrangements are mutually exclusive with mutations in EGFR or KRAS: an analysis of 1,683 patients with non-small cell lung cancer. *Clin Cancer Res: Off J Am Assoc Cancer Res* 2013;19(15):4273–81.
  - [52] Scheffler M, Ihle MA, Hein R, Merkelbach-Bruse S, Scheel AH, Siemanowski J, et al. K-ras mutation subtypes in NSCLC and associated co-occurring mutations in other oncogenic pathways. *J Thoracic Oncol: Off Publ Int Assoc Study Lung Cancer* 2019;14(4):606–16.
  - [53] Karachaliou N, Mayo C, Costa C, Magri I, Gimenez-Capitan A, Molina-Vila MA, et al. KRAS mutations in lung cancer. *Clin Lung Cancer* 2013;14(3):205–14.
  - [54] Biernacka A, Tsongalis PD, Peterson JD, de Abreu FB, Black CC, Gutmann EJ, et al. The potential utility of re-mining results of somatic mutation testing: KRAS status in lung adenocarcinoma. *Cancer Genet* 2016;209(5):195–8.
  - [55] Kafatos G, Niepel D, Lowe K, Jenkins-Anderson S, Westhead H, Garawin T, et al. RAS mutation prevalence among patients with metastatic colorectal cancer: a meta-analysis of real-world data. *Biomarkers Med* 2017;11(9):751–60.
  - [56] Watanabe T, Yoshino T, Uetake H, Yamazaki K, Ishiguro M, Kurokawa T, et al. KRAS mutational status in Japanese patients with colorectal cancer: results from a nationwide, multicenter, cross-sectional study. *Jpn J Clin Oncol* 2013;43(7):706–12.
  - [57] Andreyev HJ, Norman AR, Cunningham D, Oates JR, Clarke PA. Kirsten ras mutations in patients with colorectal cancer: the multicenter “RASCAL” study. *J Natl Cancer Inst* 1998;90(9):675–84.
  - [58] Hayama T, Hashiguchi Y, Okamoto K, Okada Y, Ono K, Shimada R, et al. G12V and G12C mutations in the gene KRAS are associated with a poorer prognosis in primary colorectal cancer. *Int J Colorectal Dis* 2019;34(8):1491–6.
  - [59] Hershkovitz D, Simon E, Bick T, Prinz E, Noy S, Sabo E, et al. Adenoma and carcinoma components in colonic tumors show discordance for KRAS mutation. *Hum Pathol* 2014;45(9):1866–71.
  - [60] Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, Leppert M, et al. Genetic alterations during colorectal-tumor development. *New Engl J Med* 1988;319(9):525–32.
  - [61] Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, et al. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science (New York, NY)* 2008;321(5897):1801–6.
  - [62] Hruban RH, van Mansfeld AD, Offerhaus GJ, van Weering DH, Allison DC, Goodman SN, et al. K-ras oncogene activation in adenocarcinoma of the human pancreas. A study of 82 carcinomas using a combination of mutant-enriched polymerase chain reaction analysis and allele-specific oligonucleotide hybridization. *Am J Pathol* 1993;143(2):545–54.
  - [63] Hezel AF, Deshpande V, Zhu AX. Genetics of biliary tract cancers and emerging targeted therapies. *J Clin Oncol: Off J Am Soc Clin Oncol* 2010;28(21):3531–40.
  - [64] Walker BA, Boyle EM, Wardell CP, Murison A, Begum DB, Dahir NM, et al. Mutational spectrum, copy number changes, and outcome: results of a sequencing study of patients with newly diagnosed myeloma. *J Clin Oncol: Off J Am Soc Clin Oncol* 2015;33(33):3911–20.
  - [65] Bezieau S, Devilder MC, Avet-Loiseau H, Mellerin MP, Puthier D, Pennarun E, et al. High incidence of N and K-Ras activating mutations in multiple myeloma and primary plasma cell leukemia at diagnosis. *Hum Mutat* 2001;18(3):212–24.
  - [66] Cox AD, Fesik SW, Kimmelman AC, Luo J, Der CJ. Drugging the undruggable RAS: mission possible? *Nat Rev Drug Discov* 2014;13(11):828–51.
  - [67] Peng N, Zhao X. Comparison of K-ras mutations in lung, colorectal and gastric cancer. *Oncol Lett* 2014;8(2):561–5.
  - [68] Ayatollahi H, Tavassoli A, Jafarian AH, Alavi A, Shakeri S, Shams SF, et al. KRAS codon 12 and 13 mutations in gastric cancer in the Northeast Iran. *Iran J Pathol* 2018;13(2):167–72.
  - [69] Arber N, Shapira I, Ratan J, Stern B, Hibshoosh H, Moshkowitz M, et al. Activation of c-K-ras mutations in human gastrointestinal tumors. *Gastroenterology* 2000;118(6):1045–50.
  - [70] Moul JW, Theune SM, Chang EH. Detection of RAS mutations in archival testicular germ cell tumors by polymerase chain reaction and oligonucleotide hybridization. *Genes Chromosom Cancer* 1992;5(2):109–18.
  - [71] Hacioglu BM, Kodaz H, Erdogan B, Cinkaya A, Tastekin E, Hacıbekiroglu I, et al. K-

- RAS and N-RAS mutations in testicular germ cell tumors. *Bosnian J Med Sci* 2017;17(2):159–63.
- [72] Ridanpaa M, Lothe RA, Onfelt A, Fossa S, Borresen AL, Husgafvel-Pursiainen K. K-ras oncogene codon 12 point mutations in testicular cancer. *Environ Health Perspect* 1993;101(Suppl 3):185–7.
- [73] Jiang W, Xiang L, Pei X, He T, Shen X, Wu X, et al. Mutational analysis of KRAS and its clinical implications in cervical cancer patients. *J Gynecol Oncol* 2018;29(1):e4.
- [74] Wright AA, Howitt BE, Myers AP, Dahlberg SE, Palescandolo E, Van Hummelen P, et al. Oncogenic mutations in cervical cancer: genomic differences between adenocarcinomas and squamous cell carcinomas of the cervix. *Cancer* 2013;119(21):3776–83.
- [75] Nagel PD, Feld FM, Weissinger SE, Stenzinger A, Moller P, Lennerz JK. Absence of BRAF and KRAS hotspot mutations in primary mediastinal B-cell lymphoma. *Leukemia Lymphoma* 2014;55(10):2389–90.
- [76] Liu QW, Fu JH, Luo KJ, Yang HX, Wang JY, Hu Y, et al. Identification of EGFR and KRAS mutations in Chinese patients with esophageal squamous cell carcinoma. *Dis Esophagus: Off J Int Soc Dis Esophagus* 2011;24(5):374–80.
- [77] Lorenzen S, Schuster T, Porschen R, Al-Batran SE, Hofheinz R, Thuss-Patience P, et al. Cetuximab plus cisplatin-5-fluorouracil versus cisplatin-5-fluorouracil alone in first-line metastatic squamous cell carcinoma of the esophagus: a randomized phase II study of the Arbeitsgemeinschaft Internistische Onkologie. *Ann Oncol: Off J Eur Soc Med Oncol* 2009;20(10):1667–73.
- [78] Essakly A, Loeser H, Kraemer M, Alakus H, Chon SH, Zander T, et al. PIK3CA and KRAS amplification in esophageal adenocarcinoma and their impact on the inflammatory tumor microenvironment and prognosis. *Transl Oncol* 2020;13(2):157–64.
- [79] Hollestelle A, Elstrodt F, Nagel JH, Kallemeijn WW, Schutte M. Phosphatidylinositol-3-OH kinase or RAS pathway mutations in human breast cancer cell lines. *Mol Cancer Res: MCR* 2007;5(2):195–201.
- [80] Pereira CB, Leal MF, de Souza CR, Montenegro RC, Rey JA, Carvalho AA, et al. Prognostic and predictive significance of MYC and KRAS alterations in breast cancer from women treated with neoadjuvant chemotherapy. *PLoS ONE* 2013;8(3):e60576.
- [81] Ahmad EI, Gawish HH, Al Azizi NM, Elhefni AM. The prognostic impact of K-RAS mutations in adult acute myeloid leukemia patients treated with high-dose cytarabine. *OncoTargets Therapy* 2011;4:115–21.
- [82] Stirewalt DL, Kopecsky KJ, Meshinchi S, Appelbaum FR, Slovak ML, Willman CL, et al. FLT3, RAS, and TP53 mutations in elderly patients with acute myeloid leukemia. *Blood* 2001;97(11):3589–95.
- [83] Neubauer A, Dodge RK, George SL, Davey FR, Silver RT, Schiffer CA, et al. Prognostic importance of mutations in the ras proto-oncogenes in de novo acute myeloid leukemia. *Blood* 1994;83(6):1603–11.
- [84] Vendramini E, Bomben R, Pozzo F, Benedetti D, Bittolo T, Rossi FM, et al. KRAS, NRAS, and BRAF mutations are highly enriched in trisomy 12 chronic lymphocytic leukemia and are associated with shorter treatment-free survival. *Leukemia* 2019;33(8):2111–5.
- [85] Gimenez N, Martinez-Trillos A, Montraveta A, Lopez-Guerra M, Rosich L, Nadeu F, et al. Mutations in the RAS-BRAF-MAPK-ERK pathway define a specific subgroup of patients with adverse clinical features and provide new therapeutic options in chronic lymphocytic leukemia. *Haematologica* 2019;104(3):576–86.
- [86] Ouerhani S, Bougateg K, Soltani I, Elgaied AB, Abbas S, Menif S. The prevalence and prognostic significance of KRAS mutation in bladder cancer, chronic myeloid leukemia and colorectal cancer. *Mol Biol Rep* 2013;40(6):4109–14.
- [87] Kodaz HKO, Hacıoglu MB, et al. Frequency of RAS mutations (KRAS, NRAS, HRAS) in human solid cancer. *EJMO* 2017;1(1):1–7.
- [88] Lax SF, Kendall B, Tashiro H, Slebos RJ, Hedrick L. The frequency of p53, K-ras mutations, and microsatellite instability differs in uterine endometrioid and serous carcinoma: evidence of distinct molecular genetic pathways. *Cancer* 2000;88(4):814–24.
- [89] Slebos RJ, Kibbelaar RE, Dalesio O, Kooistra A, Stam J, Meijer CJ, et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *New Engl J Med* 1990;323(9):561–5.
- [90] Shepherd FA, Domerg C, Hainaut P, Janne PA, Pignon JP, Graziano S, et al. Pooled analysis of the prognostic and predictive effects of KRAS mutation status and KRAS mutation subtype in early-stage resected non-small-cell lung cancer in four trials of adjuvant chemotherapy. *J Clin Oncol: Off J Am Soc Clin Oncol* 2013;31(17):2173–81.
- [91] Zer A, Ding K, Lee SM, Goss GD, Seymour L, Ellis PM, et al. Pooled analysis of the prognostic and predictive value of KRAS mutation status and mutation subtype in patients with non-small cell lung cancer treated with epidermal growth factor receptor tyrosine kinase inhibitors. *J Thoracic Oncol: Off Publ Int Assoc Study Lung Cancer* 2016;11(3):312–23.
- [92] Pan W, Yang Y, Zhu H, Zhang Y, Zhou R, Sun X. KRAS mutation is a weak, but valid predictor for poor prognosis and treatment outcomes in NSCLC: a meta-analysis of 41 studies. *Oncotarget* 2016;7(7):8373–88.
- [93] Cerottini JP, Caplin S, Saraga E, Givel JC, Benhattar J. The type of K-ras mutation determines prognosis in colorectal cancer. *Am J Surg* 1998;175(3):198–202.
- [94] Samowitz WS, Curtin K, Schaffer D, Robertson M, Leppert M, Slattery ML. Relationship of Ki-ras mutations in colon cancers to tumor location, stage, and survival: a population-based study. *Cancer Epidemiol Biomarkers Prevent: Publ Am Assoc Cancer Res Cosponsored Am Soc Prevent Oncol* 2000;9(11):1193–7.
- [95] Yoon HH, Tougeron D, Shi Q, Alberts SR, Mahoney MR, Nelson GD, et al. KRAS codon 12 and 13 mutations in relation to disease-free survival in BRAF-wild-type stage III colon cancers from an adjuvant chemotherapy trial (N0147 alliance). *Clin Cancer Res: Off J Am Assoc Cancer Res* 2014;20(11):3033–43.
- [96] Modest DP, Ricard I, Heinemann V, Hegewisch-Becker S, Schmieg W, Porschen R, et al. Outcome according to KRAS-, NRAS- and BRAF-mutation as well as KRAS mutation variants: pooled analysis of five randomized trials in metastatic colorectal cancer by the AIO colorectal cancer study group. *Ann Oncol: Off J Eur Soc Med Oncol* 2016;27(9):1746–53.
- [97] Taieb J, Zaanen A, Le Malicot K, Julie C, Blons H, Mineur L, et al. Prognostic effect of BRAF and KRAS mutations in patients with stage III colon cancer treated with leucovorin, fluorouracil, and oxaliplatin with or without cetuximab: a post hoc analysis of the PETACC-8 trial. *JAMA Oncol* 2016;2(5):643–53.
- [98] Roth AD, Tejpar S, Delorenzi M, Yan P, Fiocca R, Klingbiel D, et al. Prognostic role of KRAS and BRAF in stage II and III resected colon cancer: results of the translational study on the PETACC-3, EORTC 40993, SAKK 60–00 trial. *J Clin Oncol: Off J Am Soc Clin Oncol* 2010;28(3):466–74.
- [99] Andreyev HJ, Norman AR, Cunningham D, Oates J, Dix BR, Iacopetta BJ, et al. Kirsten ras mutations in patients with colorectal cancer: the 'RASCAL II' study. *Br J Cancer* 2001;85(5):692–6.
- [100] Haas M, Ormanns S, Baechmann S, Remold A, Kruger S, Westphalen CB, et al. Extended RAS analysis and correlation with overall survival in advanced pancreatic cancer. *Br J Cancer* 2017;116(11):1462–9.
- [101] Shin SH, Kim SC, Hong SM, Kim YH, Song KB, Park KM, et al. Genetic alterations of K-ras, p53, c-erbB-2, and DPC4 in pancreatic ductal adenocarcinoma and their correlation with patient survival. *Pancreas* 2013;42(2):216–22.
- [102] Windon AL, Loaiza-Bonilla A, Jensen CE, Randall M, Morrisette JJD, Shroff SG. A KRAS wild type mutational status confers a survival advantage in pancreatic ductal adenocarcinoma. *J Gastrointestinal Oncol* 2018;9(1):1–10.
- [103] Bournet B, Muscare F, Buscail C, Assenat E, Barthet M, Hammel P, et al. KRAS G12D mutation subtype is a prognostic factor for advanced pancreatic adenocarcinoma. *Clin Translat Gastroenterol* 2016;7:e157.
- [104] Rachakonda PS, Bauer AS, Xie H, Campa D, Rizzato C, Canzian F, et al. Somatic mutations in exocrine pancreatic tumors: association with patient survival. *PLoS ONE* 2013;8(4):e60870.
- [105] Rodenhuis S, Boerrigter L, Top B, Slebos RJ, Mooi WJ, van't Veer L, et al. Mutational activation of the K-ras oncogene and the effect of chemotherapy in advanced adenocarcinoma of the lung: a prospective study. *J Clin Oncol: Off J Am Soc Clin Oncol* 1997;15(1):285–91.
- [106] Herbst RS, Prager D, Hermann R, Fehrenbacher L, Johnson BE, Sandler A, et al. TRIBUTE: a phase III trial of erlotinib hydrochloride (OSI-774) combined with carboplatin and paclitaxel chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol: Off J Am Soc Clin Oncol* 2005;23(25):5892–9.
- [107] Zhu CQ, da Cunha Santos G, Ding K, Sakurada A, Cutz JC, Liu N, et al. Role of KRAS and EGFR as biomarkers of response to erlotinib in National Cancer Institute of Canada Clinical Trials Group Study BR.21. *J Clin Oncol: Off J Am Soc Clin Oncol* 2008;26(26):4268–75.
- [108] Douillard JY, Shepherd FA, Hirsh V, Mok T, Socinski MA, Gervais R, et al. Molecular predictors of outcome with gefitinib and docetaxel in previously treated non-small-cell lung cancer: data from the randomized phase III INTEREST trial. *J Clin Oncol: Off J Am Soc Clin Oncol* 2010;28(5):744–52.
- [109] Schneider CP, Heigener D, Schott-von-Romer K, Gutz S, Laack E, Digel W, et al. Epidermal growth factor receptor-related tumor markers and clinical outcomes with erlotinib in non-small cell lung cancer: an analysis of patients from german centers in the TRUST study. *J Thoracic Oncol: Off Publ Int Assoc Study Lung Cancer* 2008;3(12):1446–53.
- [110] Kim JH, Kim HS, Kim BJ. Prognostic value of KRAS mutation in advanced non-small-cell lung cancer treated with immune checkpoint inhibitors: a meta-analysis and review. *Oncotarget* 2017;8(29):48248–52.
- [111] Skoulidis F, Goldberg ME, Greenawald DM, Hellmann MD, Awad MM, Gainor JF, et al. STK11/LKB1 mutations and PD-1 inhibitor resistance in KRAS-mutant lung adenocarcinoma. *Cancer Discov* 2018;8(7):822–35.
- [112] Amado RG, Wolf M, Peeters M, Van Cutsem E, Siena S, Freeman DJ, et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol: Off J Am Soc Clin Oncol* 2008;26(10):1626–34.
- [113] De Roock W, Piessevaux H, De Schutter J, Janssens M, De Hertogh G, Personeni N, et al. KRAS wild-type state predicts survival and is associated to early radiological response in metastatic colorectal cancer treated with cetuximab. *Ann Oncol: Off J Eur Soc Med Oncol* 2008;19(3):508–15.
- [114] Karapetis CS, Khambata-Ford S, Jonker DJ, O'Callaghan CJ, Tu D, Tebbutt NC, et al. K-ras mutations and benefit from cetuximab in advanced colorectal cancer. *New Engl J Med* 2008;359(17):1757–65.
- [115] Douillard JY, Oliner KS, Siena S, Tabernero J, Burkes R, Barugel M, et al. Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer. *New Engl J Med* 2013;369(11):1023–34.
- [116] Tejpar S, Celik I, Schlichting M, Sartorius U, Bokemeyer C, Van Cutsem E. Association of KRAS G13D tumor mutations with outcome in patients with metastatic colorectal cancer treated with first-line chemotherapy with or without cetuximab. *J Clin Oncol: Off J Am Soc Clin Oncol* 2012;30(29):3570–7.
- [117] De Roock W, Jonker DJ, Di Nicolantonio F, Sartore-Bianchi A, Tu D, Siena S, et al. Association of KRAS p.G13D mutation with outcome in patients with chemotherapy-refractory metastatic colorectal cancer treated with cetuximab. *Jama* 2010;304(16):1812–20.
- [118] Janssen KP, el-Marjou F, Pinto D, Sastre X, Rouillard D, Fouquet C, et al. Targeted expression of oncogenic K-ras in intestinal epithelium causes spontaneous tumorigenesis in mice. *Gastroenterology* 2002;123(2):492–504.
- [119] Caulin C, Nguyen T, Longley MA, Zhou Z, Wang XJ, Roop DR. Inducible activation of oncogenic K-ras results in tumor formation in the oral cavity. *Cancer Res* 2004;64(15):5054–8.
- [120] Vitale-Cross L, Amornphimoltham P, Fisher G, Molinolo AA, Gutkind JS. Conditional expression of K-ras in an epithelial compartment that includes the



- stem cells is sufficient to promote squamous cell carcinogenesis. *Cancer Res* 2004;64(24):8804–7.
- [121] Collins MA, Brisset JC, Zhang Y, Bednar F, Pierre J, Heist KA, et al. Metastatic pancreatic cancer is dependent on oncogenic Kras in mice. *PLoS ONE* 2012;7(12):e49707.
- [122] Fisher GH, Wellen SL, Klimstra D, Lenczowski JM, Tichelaar JW, Lizak MJ, et al. Induction and apoptotic regression of lung adenocarcinomas by regulation of a K-Ras transgene in the presence and absence of tumor suppressor genes. *Genes Dev* 2001;15(24):3249–62.
- [123] Kwong LN, Costello JC, Liu H, Jiang S, Helms TL, Langsdorf AE, et al. Oncogenic NRAS signaling differentially regulates survival and proliferation in melanoma. *Nat Med* 2012;18(10):1503–10.
- [124] Taveras AG, Remiszewski SW, Doll RJ, Cesarz D, Huang EC, Kirschmeier P, et al. Ras oncoprotein inhibitors: the discovery of potent, ras nucleotide exchange inhibitors and the structural determination of a drug-protein complex. *Bioorg Med Chem* 1997;5(1):125–33.
- [125] Ganguly AK, Wang YS, Pramanik BN, Doll RJ, Snow ME, Taveras AG, et al. Interaction of a novel GDP exchange inhibitor with the Ras protein. *Biochemistry* 1998;37(45):15631–7.
- [126] Jansen B, Zangemeister-Wittke U. Antisense therapy for cancer—the time of truth. *Lancet Oncol* 2002;3(11):672–83.
- [127] Gray GD, Hernandez OM, Hebel D, Root M, Pow-Sang JM, Wickstrom E. Antisense DNA inhibition of tumor growth induced by c-Ha-ras oncogene in nude mice. *Cancer Res* 1993;53(3):577–80.
- [128] Adjei AA, Dy GK, Erlichman C, Reid JM, Sloan JA, Pitot HC, et al. A phase I trial of ISIS 2503, an antisense inhibitor of H-ras, in combination with gemcitabine in patients with advanced cancer. *Clin Cancer Res: Off J Am Assoc Cancer Res* 2003;9(1):115–23.
- [129] Cunningham CC, Holmlund JT, Geary RS, Kwok TJ, Dorr A, Johnston JF, et al. A Phase I trial of H-ras antisense oligonucleotide ISIS 2503 administered as a continuous intravenous infusion in patients with advanced carcinoma. *Cancer* 2001;92(5):1265–71.
- [130] Marshall JL, Eisenberg SG, Johnson MD, Hanfelt J, Dorr FA, El-Ashry D, et al. A phase II trial of ISIS 3521 in patients with metastatic colorectal cancer. *Clin Colorectal Cancer* 2004;4(4):268–74.
- [131] Alberts SR, Schroeder M, Erlichman C, Steen PD, Foster NR, Moore Jr DF, et al. Gemcitabine and ISIS-2503 for patients with locally advanced or metastatic pancreatic adenocarcinoma: a North Central Cancer Treatment Group phase II trial. *J Clin Oncol: Off J Am Soc Clin Oncol* 2004;22(24):4944–50.
- [132] Haluska P, Dy GK, Adjei AA. Farnesyl transferase inhibitors as anticancer agents. *Eur J Cancer (Oxford, England: 1990)* 2002;38(13):1685–700.
- [133] Adjei AA, Mauer A, Bruzek L, Marks RS, Hillman S, Geyer S, et al. Phase II study of the farnesyl transferase inhibitor R115777 in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2003;21(9):1760–6.
- [134] Sharma S, Kemeny N, Kelsen DP, Ilson D, O'Reilly E, Zaknoon S, et al. A phase II trial of farnesyl protein transferase inhibitor SCH 66336, given by twice-daily oral administration, in patients with metastatic colorectal cancer refractory to 5-fluorouracil and irinotecan. *Ann Oncol: Off J Eur Soc Med Oncol* 2002;13(7):1067–71.
- [135] Winquist E, Moore MJ, Chi KN, Ernst DS, Hirte H, North S, et al. A multinomial Phase II study of lonafarnib (SCH 66336) in patients with refractory urothelial cancer. *Urol Oncol* 2005;23(3):143–9.
- [136] Gajewski TF, Salama AKS, Niedzwiecki D, Johnson J, Linette G, Bucher C, et al. Phase II study of the farnesyltransferase inhibitor R115777 in advanced melanoma (CALGB 500104). *J Trans Med* 2012;10(1):246.
- [137] Cohen SJ, Ho L, Ranganathan S, Abbruzzese JL, Alpaugh RK, Beard M, et al. Phase II and pharmacodynamic study of the farnesyltransferase inhibitor R115777 as initial therapy in patients with metastatic pancreatic adenocarcinoma. *J Clin Oncol* 2003;21(7):1301–6.
- [138] Rao S, Cunningham D, de Gramont A, Scheithauer W, Smakal M, Humblet Y, et al. Phase III double-blind placebo-controlled study of farnesyl transferase inhibitor R115777 in patients with refractory advanced colorectal cancer. *J Clin Oncol* 2004;22(19):3950–7.
- [139] Van Cutsem E, van de Velde H, Karasek P, Oettle H, Vervenne WL, Szawlowski A, et al. Phase III trial of gemcitabine plus tipifarnib compared with gemcitabine plus placebo in advanced pancreatic cancer. *J Clin Oncol* 2004;22(8):1430–8.
- [140] Luger SXV, Paietta E, et al. Tipifarnib as maintenance therapy in acute myeloid leukemia (AML) improves survival in a subgroup of patients with high risk disease. Results of the phase III intergroup trial E2902. *Blood* 2015;126(23):1308.
- [141] Kim ES, Kies MS, Fossella FV, Glisson BS, Zaknoon S, Statkevich P, et al. Phase II study of the farnesyltransferase inhibitor lonafarnib with paclitaxel in patients with taxane-refractory/resistant non-small cell lung carcinoma. *Cancer* 2005;104(3):561–9.
- [142] Whyte DB, Kirschmeier P, Hockenberry TN, Nunez-Oliva I, James L, Catino JJ, et al. K- and N-Ras are geranylgeranylated in cells treated with farnesyl protein transferase inhibitors. *J Biol Chem* 1997;272(22):14459–64.
- [143] Ho A BI, Haddad R, et al. Preliminary results from a phase 2 trial of tipifarnib in Head and Neck Squamous Cell Carcinomas (HSCCs) with HRAS mutations. In: Presented at AACR-NCI-EORTC international conference on molecular targets and cancer; October 26–30, 2019; Boston, MA. Abstract 384.
- [144] Karasic TB, Chiorean EG, Sefti SM, O'Dwyer PJ. A phase I study of GGTI-2418 (geranylgeranyl transferase I inhibitor) in patients with advanced solid tumors. *Target Oncol* 2019;14(5):613–8.
- [145] Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, et al. Sorafenib in advanced hepatocellular carcinoma. *New Engl J Med* 2008;359(4):378–90.
- [146] Escudier B, Eisen T, Stadler WM, Szczylak C, Oudard S, Siebels M, et al. Sorafenib in advanced clear-cell renal-cell carcinoma. *New Engl J Med* 2007;356(2):125–34.
- [147] Brose MS, Nutting CM, Jarzab B, Elisei R, Siena S, Bastholt L, et al. Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial. *Lancet (London, England)* 2014;384(9940):319–28.
- [148] Wilhelm SM, Carter C, Tang L, Wilkie D, McNabola A, Rong H, et al. BAY 43–9006 exhibits broad spectrum oral antitumor activity and targets the RAF/MEK/ERK pathway and receptor tyrosine kinases involved in tumor progression and angiogenesis. *Cancer Res* 2004;64(19):7099–109.
- [149] Cascinu S, Berardi R, Sobrero A, Bidoli P, Labianca R, Siena S, et al. Sorafenib does not improve efficacy of chemotherapy in advanced pancreatic cancer: a GISCAD randomized phase II study. *Dig Liver Dis* 2014;46(2):182–6.
- [150] Kindler HL, Wroblewski K, Wallace JA, Hall MJ, Locker G, Nattam S, et al. Gemcitabine plus sorafenib in patients with advanced pancreatic cancer: a phase II trial of the University of Chicago Phase II Consortium. *Invest New Drugs* 2012;30(1):382–6.
- [151] Hauschild A, Grob JJ, Demidov LV, Jouary T, Gutzmer R, Millward M, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet (London, England)* 2012;380(9839):358–65.
- [152] Subbiah V, Kreitman RJ, Wainberg ZA, Cho JY, Schellens JHM, Soria JC, et al. Dabrafenib and trametinib treatment in patients with locally advanced or metastatic BRAF V600-mutant anaplastic thyroid cancer. *J Clin Oncol: Off J Am Soc Clin Oncol* 2018;36(1):7–13.
- [153] Hyman DM, Puzanov I, Subbiah V, Faris JE, Chau I, Blay JY, et al. Vemurafenib in multiple nonmelanoma cancers with BRAF V600 mutations. *New Engl J Med* 2015;373(8):726–36.
- [154] Poulikakos PI, Zhang C, Bollag G, Shokat KM, Rosen N. RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF. *Nature* 2010;464(7287):427–30.
- [155] Heidorn SJ, Milagre C, Whittaker S, Noury A, Niculescu-Duvaz I, Dhomen N, et al. Kinase-dead BRAF and oncogenic RAS cooperate to drive tumor progression through CRAF. *Cell* 2010;140(2):209–21.
- [156] Hatzivassiliou G, Song K, Yen I, Brandhuber BJ, Anderson DJ, Alvarado R, et al. RAF inhibitors prime wild-type RAF to activate the MAPK pathway and enhance growth. *Nature* 2010;464(7287):431–5.
- [157] Dummer R, Ascierto PA, Gogas HJ, Arance A, Mandalà M, Liszkay G, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2018;19(5):603–15.
- [158] Van Cutsem E, Huijberts S, Grothey A, Yaeger R, Cuyle PJ, Elez E, et al. Binimetinib, encorafenib, and cetuximab triplet therapy for patients with BRAF V600E-mutant metastatic colorectal cancer: safety lead-in results from the phase III BEACON colorectal cancer study. *J Clin Oncol: Off J Am Soc Clin Oncol* 2019;37(17):1460–9.
- [159] Planchard D, Besse B, Groen HJM, Souquet PJ, Quoix E, Baik CS, et al. Dabrafenib plus trametinib in patients with previously treated BRAF(V600E)-mutant metastatic non-small cell lung cancer: an open-label, multicentre phase 2 trial. *Lancet Oncol* 2016;17(7):984–93.
- [160] Rinehart J, Adjei AA, Lorusso PM, Waterhouse D, Hecht JR, Natale RB, et al. Multicenter phase II study of the oral MEK inhibitor, CI-1040, in patients with advanced non-small-cell lung, breast, colon, and pancreatic cancer. *J Clin Oncol* 2004;22(22):4456–62.
- [161] Hainsworth JD, Cebotaru CL, Kanarev V, Ciuleanu TE, Danyanov D, Stella P, et al. A phase II, open-label, randomized study to assess the efficacy and safety of AZD6244 (ARRY-142886) versus pemetrexed in patients with non-small cell lung cancer who have failed one or two prior chemotherapeutic regimens. *J Thoracic Oncol: Off Publ Int Assoc Study Lung Cancer* 2010;5(10):1630–6.
- [162] Mirzoeva OK, Collisson EA, Schaefer PM, Hann B, Hom YK, Ko AH, et al. Subtype-specific MEK-P13 kinase feedback as a therapeutic target in pancreatic adenocarcinoma. *Mol Cancer Ther* 2013;12(10):2213–25.
- [163] Carter CA, Mol CA, Keen C, Szabo E, Khozin S, Thomas A, et al. Selumetinib with and without erlotinib in KRAS mutant and KRAS wild-type advanced non-small-cell lung cancer. *Ann Oncol* 2016;27(4):693–9.
- [164] Ko AH, Bekaii-Saab T, Van Ziffle J, Mirzoeva OM, Joseph NM, Talasz A, et al. A Multicenter, open-label phase II clinical trial of combined MEK plus EGFR inhibition for chemotherapy-refractory advanced pancreatic adenocarcinoma. *Clin Cancer Res: Off J Am Assoc Cancer Res* 2016;22(1):61–8.
- [165] Gandara DR, Leighl N, Delord JP, Barlesi F, Bannouna J, Zalcman G, et al. A phase 1/1b study evaluating trametinib plus docetaxel or pemetrexed in patients with advanced non-small cell lung cancer. *J Thoracic Oncol: Off Publ Int Assoc Study Lung Cancer* 2017;12(3):556–66.
- [166] Kawaguchi K, Igarashi K, Miyake K, Lwin TM, Miyake M, Kiyuna T, et al. MEK inhibitor trametinib in combination with gemcitabine regresses a patient-derived orthotopic xenograft (PDOX) pancreatic cancer nude mouse model. *Tissue Cell* 2018;52:124–8.
- [167] Kawaguchi K, Igarashi K, Murakami T, Kiyuna T, Lwin TM, Hwang HK, et al. MEK inhibitors cobimetinib and trametinib, regressed a gemcitabine-resistant pancreatic-cancer patient-derived orthotopic xenograft (PDOX). *Oncotarget* 2017;8(29):47490–6.
- [168] Infante JR, Somer BG, Park JO, Li CP, Scheulen ME, Kasubhai SM, et al. A randomised, double-blind, placebo-controlled trial of trametinib, an oral MEK inhibitor, in combination with gemcitabine for patients with untreated metastatic adenocarcinoma of the pancreas. *Eur J Cancer (Oxford, England: 1990)* 2014;50(12):2072–81.
- [169] Janne PA, van den Heuvel MM, Barlesi F, Cobo M, Mazieres J, Crino L, et al.

- Selumetinib plus docetaxel compared with docetaxel alone and progression-free survival in patients with KRAS-mutant advanced non-small cell lung cancer: the SELECT-1 randomized clinical trial. *JAMA* 2017;317(18):1844–53.
- [170] Bhagwat SV, McMillen WT, Cai S, Zhao B, Whitesell M, Shen W, et al. ERK inhibitor LY3214996 targets ERK pathway-driven cancers: a therapeutic approach toward precision medicine. *Mol Cancer Ther* 2020;19(2):325–36.
- [171] Pant S, Bendell JC, Sullivan RJ, Shapiro G, Millward M, Mi G, et al. A phase I dose escalation (DE) study of ERK inhibitor, LY3214996, in advanced (adv) cancer (CA) patients (pts). 2019;37(15\_suppl):3001.
- [172] Puyol M, Martin A, Dubus P, Mulero F, Pizcueta P, Khan G, et al. A synthetic lethal interaction between K-Ras oncogenes and Cdk4 unveils a therapeutic strategy for non-small cell lung carcinoma. *Cancer Cell* 2010;18(1):63–73.
- [173] Goldman JW, Mazieres J, Barlesi F, Koczywas M, Dragnev KH, Göksel T, et al. A randomized phase 3 study of abemaciclib versus erlotinib in previously treated patients with stage IV NSCLC with KRAS mutation. *JUNIPER* 2018;36(15\_suppl):9025.
- [174] Blumenschein Jr. GR, Smit EF, Planchard D, Kim DW, Cadranet J, De Pas T, et al. A randomized phase II study of the MEK1/MEK2 inhibitor trametinib (GSK1120212) compared with docetaxel in KRAS-mutant advanced non-small-cell lung cancer (NSCLC). *Ann Oncol* 2015;26(5):894–901.
- [175] Ostrem JM, Peters U, Sos ML, Wells JA, Shokat KM. K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions. *Nature* 2013;503(7477):548–51.
- [176] Patricelli MP, Janes MR, Li LS, Hansen R, Peters U, Kessler LV, et al. Selective inhibition of oncogenic KRAS output with small molecules targeting the inactive state. *Cancer Discov* 2016;6(3):316–29.
- [177] Lito P, Solomon M, Li LS, Hansen R, Rosen N. Allele-specific inhibitors inactivate mutant KRAS G12C by a trapping mechanism. *Science* 2016;351(6273):604–8.
- [178] Janes MR, Zhang J, Li LS, Hansen R, Peters U, Guo X, et al. Targeting KRAS mutant cancers with a covalent G12C-specific inhibitor. *Cell* 2018;172(3): 578–89.e17.
- [179] Canon J, Rex K, Saiki AY, Mohr C, Cooke K, Bagal D, et al. The clinical KRAS (G12C) inhibitor AMG 510 drives anti-tumour immunity. *Nature* 2019;575(7781):217–23.
- [180] Hallin J, Engstrom LD, Hargis L, Calinisan A, Aranda R, Briere DM, et al. The KRAS (G12C) inhibitor MRTX849 provides insight toward therapeutic susceptibility of KRAS-mutant cancers in mouse models and patients. *Cancer Discov* 2020;10(1):54–71.
- [181] Bar-Sagi D, Knelson EH, Sequist LV. A bright future for KRAS inhibitors. *Nature Cancer* 2020;1(1):25–7.
- [182] Neumann J, Zeindl-Eberhart E, Kirchner T, Jung A. Frequency and type of KRAS mutations in routine diagnostic analysis of metastatic colorectal cancer. *Pathol Res Pract* 2009;205(12):858–62.
- [183] Miller MS, Miller LD. RAS mutations and oncogenesis: not all RAS mutations are created equally. *Front Genet* 2011;2:100.
- [184] Govindan RFM, Price T, et al. Phase 1 study of safety, tolerability, PK, and efficacy of AMG 510, a novel KRAS G12C inhibitor, evaluated in NSCLC. In: Presented at the IASLC 2019 world lung cancer conference; September 7–10, 2019; Barcelona, Spain.
- [185] Amodio V, Yaeger R, Arcella P, Cancelliere C, Lamba S, Lorenzato A, et al. EGFR blockade reverts resistance to KRAS G12C inhibition in colorectal cancer. *Cancer Discov* 2020.
- [186] Hofmann MHGM, Ramharter J, et al. Discovery of BI-3406: a potent and selective SOS1:: KRAS inhibitor opens a new approach for treating KRAS-driven tumors. In: Presented at AACR-NCI-EORTC international conference on molecular targets and cancer therapeutics; October 26–30, 2019; Boston, MA. Abstract PL06-01.
- [187] Dance M, Montagner A, Salles JP, Yart A, Raynal P. The molecular functions of Shp2 in the Ras/Mitogen-activated protein kinase (ERK1/2) pathway. *Cell Signal* 2008;20(3):453–9.
- [188] Chen YN, LaMarche MJ, Chan HM, Fekkes P, Garcia-Fortanet J, Acker MG, et al. Allosteric inhibition of SHP2 phosphatase inhibits cancers driven by receptor tyrosine kinases. *Nature* 2016;535(7610):148–52.
- [189] Mainardi S, Mulero-Sanchez A, Prahallad A, Germano G, Bosma A, Krimpenfort P, et al. SHP2 is required for growth of KRAS-mutant non-small-cell lung cancer in vivo. *Nat Med* 2018;24(7):961–7.
- [190] Hernandez G, Vazquez-Pianzola P. Functional diversity of the eukaryotic translation initiation factors belonging to eIF4 families. *Mech Dev* 2005;122(7–8):865–76.
- [191] Thompson PAEB, Young NP, et al. eFT226, a potent and selective inhibitor of eIF4A, is efficacious in preclinical models of lymphoma. In: Presented at AACR Annual Meeting 2019; March 29–April 3, 2019; Atlant, GA. Abstract 2698.
- [192] Thompson PAYN, Stumpf CR, et al. eFT226, a first in class inhibitor of eIF4A1, targets FEGFR1/2 and HER2 driven cancers. In: Presented at AACR-NCI-EORTC International Conference on Molecular Targets and Cancer Therapeutics; October 26–30, 2019; Boston, MA. Abstract B133.
- [193] Sahin U, Derhovanessian E, Miller M, Kloeke BP, Simon P, Löwer M, et al. Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature* 2017;547(7662):222–6.
- [194] Sakamoto K, Kamada Y, Sameshima T, Yaguchi M, Niida A, Sasaki S, et al. K-Ras (G12D)-selective inhibitory peptides generated by random peptide T7 phage display technology. *Biochem Biophys Res Commun* 2017;484(3):605–11.
- [195] Niida A, Sasaki S, Yonemori K, Sameshima T, Yaguchi M, Asami T, et al. Investigation of the structural requirements of K-Ras(G12D) selective inhibitory peptide KRpep-2d using alanine scans and cysteine bridging. *Bioorg Med Chem Lett* 2017;27(12):2757–61.
- [196] Sogabe S, Kamada Y, Miwa M, Niida A, Sameshima T, Kamaura M, et al. Crystal structure of a human K-ras G12D mutant in complex with GDP and the cyclic inhibitory peptide KRpep-2d. *ACS Med Chem Lett* 2017;8(7):732–6.