

Math Refresher
Winter Institute in Data Science

Ryan T. Moore

2021-01-03

Functions

Vectors

Matrix Algebra

Systems of Linear Equations

Warming Up

1. Let “ $A \circledast B$ ” be defined as $A^B + A \cdot B$. Calculate $4 \circledast 3$.
2. Solve this system of two linear equations:

$$2x - y = 4$$

$$x + y = 5$$

Functions

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.
- ▶ Range (“Image’ ”): set of all possible (obtained) output values. Set of elements of Y assigned to elements of X by $f(x)$. $f(X) = \{y : y = f(x), x \in X\}$

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.
- ▶ Range (“Image’ ”): set of all possible (obtained) output values. Set of elements of Y assigned to elements of X by $f(x)$. $f(X) = \{y : y = f(x), x \in X\}$
- ▶ Mapping notation examples

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.
- ▶ Range (“Image’ ”): set of all possible (obtained) output values. Set of elements of Y assigned to elements of X by $f(x)$. $f(X) = \{y : y = f(x), x \in X\}$
- ▶ Mapping notation examples
 - ▶ $f : X \rightarrow Y$

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.
- ▶ Range (“Image’ ”): set of all possible (obtained) output values. Set of elements of Y assigned to elements of X by $f(x)$. $f(X) = \{y : y = f(x), x \in X\}$
- ▶ Mapping notation examples
 - ▶ $f : X \rightarrow Y$
 - ▶ Function of one variable: $f : \mathbb{R}^1 \rightarrow \mathbb{R}^1$

Introduction to Functions

- ▶ A **function** is a relation (or mapping or transformation) that assigns every member of the domain to exactly one member of the range.
- ▶ Domain: set of all possible inputs. Set X over which $f(x)$ defined.
- ▶ Range (“Image’ ”): set of all possible (obtained) output values. Set of elements of Y assigned to elements of X by $f(x)$. $f(X) = \{y : y = f(x), x \in X\}$
- ▶ Mapping notation examples
 - ▶ $f : X \rightarrow Y$
 - ▶ Function of one variable: $f : \mathbb{R}^1 \rightarrow \mathbb{R}^1$
 - ▶ Function of two variables: $f : \mathbb{R}^2 \rightarrow \mathbb{R}^1, f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

Functions

Examples:

► $f(x) = x + 1$

For each x in \mathbb{R}^1 , $f(x)$ assigns the number $x + 1$.

Functions

Examples:

► $f(x) = x + 1$

For each x in \mathbb{R}^1 , $f(x)$ assigns the number $x + 1$.

► $f(x, y) = x^2 + y^2$

For each ordered pair (x, y) in \mathbb{R}^2 , $f(x, y)$ assigns the number $x^2 + y^2$

Functions

Examples:

► $f(x) = x + 1$

For each x in \mathbb{R}^1 , $f(x)$ assigns the number $x + 1$.

► $f(x, y) = x^2 + y^2$

For each ordered pair (x, y) in \mathbb{R}^2 , $f(x, y)$ assigns the number $x^2 + y^2$

► Often use one variable x as input and another y as output.

Example: $y = x + 1$

Functions

Examples:

- ▶ $f(x) = x + 1$

For each x in \mathbb{R}^1 , $f(x)$ assigns the number $x + 1$.

- ▶ $f(x, y) = x^2 + y^2$

For each ordered pair (x, y) in \mathbb{R}^2 , $f(x, y)$ assigns the number $x^2 + y^2$

- ▶ Often use one variable x as input and another y as output.

Example: $y = x + 1$

- ▶ Input variable: *predictor, covariate, indep var*
- ▶ Output variable: *outcome, response, dep var*

Example 1

$$f(x) = \frac{3}{1+x^2}$$

Domain $X =$

Example 1

$$f(x) = \frac{3}{1+x^2}$$

Domain $X = \mathbb{R}^1$

Example 1

$$f(x) = \frac{3}{1+x^2}$$

Domain $X = \mathbb{R}^1$


Range $f(X) =$

Example 1

$$f(x) = \frac{3}{1+x^2}$$

Domain $X = \mathbb{R}^1$

Range $f(X) = (0, 3]$



Example 2

$$f(x) = \begin{cases} x + 1, & 1 \leq x \leq 2 \\ 0, & x = 0 \\ 1 - x & -2 \leq x \leq -1 \end{cases}$$

Domain $X =$

Example 2

$$f(x) = \begin{cases} x + 1, & 1 \leq x \leq 2 \\ 0, & x = 0 \\ 1 - x & -2 \leq x \leq -1 \end{cases}$$

Domain $X = [-2, -1] \cup \{0\} \cup [1, 2]$

Example 2

$$f(x) = \begin{cases} x + 1, & 1 \leq x \leq 2 \\ 0, & x = 0 \\ 1 - x & -2 \leq x \leq -1 \end{cases}$$

Domain $X = [-2, -1] \cup \{0\} \cup [1, 2]$


Range $f(X) =$

Example 2

$$f(x) = \begin{cases} x + 1, & 1 \leq x \leq 2 \\ 0, & x = 0 \\ 1 - x & -2 \leq x \leq -1 \end{cases}$$

Domain $X = [-2, -1] \cup \{0\} \cup [1, 2]$

Range $f(X) = [2, 3] \cup \{0\}$



Example 3

$$f(x) = \frac{1}{x}$$

Domain $X =$

Example 3

$$f(x) = \frac{1}{x}$$

Domain $X = \mathbb{R}^1 \setminus \{0\}$

Example 3

$$f(x) = \frac{1}{x}$$

Domain $X = \mathbb{R}^1 \setminus \{0\}$


Range $f(X) =$

Example 3

$$f(x) = \frac{1}{x}$$

Domain $X = \mathbb{R}^1 \setminus \{0\}$

Range $f(X) = \mathbb{R}^1 \setminus \{0\}$



Example 4

$$f(x, y) = x^2 + y^2$$

Domain $X = \mathbb{R}^2$

Example 4

$$f(x, y) = x^2 + y^2$$

Domain $X = \mathbb{R}^2$


Image $f(X, Y) =$

Example 4

$$f(x, y) = x^2 + y^2$$

Domain $X = \mathbb{R}^2$

Image $f(X, Y) = \mathbb{R}_+^1 \cup \{0\}$



General Types of Functions

► **Monomials:** $f(x) = ax^k$

a is the coefficient. k is the degree.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

General Types of Functions

► **Monomials:** $f(x) = ax^k$

a is the coefficient. k is the degree.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

► **Polynomials:** sum of monomials.

Examples: $y = -\frac{1}{2}x^3 + x^2$, $y = 3x + 5$

The degree of a polynomial is the highest degree of its monomial terms. Write polynomials with terms in decreasing degree.

General Types of Functions

► **Monomials:** $f(x) = ax^k$

a is the coefficient. k is the degree.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

► **Polynomials:** sum of monomials.

Examples: $y = -\frac{1}{2}x^3 + x^2$, $y = 3x + 5$

The degree of a polynomial is the highest degree of its monomial terms. Write polynomials with terms in decreasing degree.

► **Rational Functions:** ratio of two polynomials.

Examples: $y = \frac{x}{2}$, $y = \frac{x^2+1}{x^2-2x+1}$

► **Exponential Functions:** Example: $y = 2^x$

General Types of Functions

► **Monomials:** $f(x) = ax^k$

a is the coefficient. k is the degree.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

► **Polynomials:** sum of monomials.

Examples: $y = -\frac{1}{2}x^3 + x^2$, $y = 3x + 5$

The degree of a polynomial is the highest degree of its monomial terms. Write polynomials with terms in decreasing degree.

► **Rational Functions:** ratio of two polynomials.

Examples: $y = \frac{x}{2}$, $y = \frac{x^2+1}{x^2-2x+1}$

► **Exponential Functions:** Example: $y = 2^x$

► **Trigonometric Functions:** Examples: $y = \cos(x)$,
 $y = 3 \sin(4x)$

Trigonometric Functions: Gill & Casella (2004)





Fig. 1 A highly multimodal surface.

$$f(x, y) = |(x \sin(20y - 90) - y \cos(20x + 45))^3 a \cos(\sin(90y + 42)x) + (x \cos(10y + 10) - y \sin(10x + 15))^2 a \cos(\cos(10x + 24)y)|$$

Linear and Nonlinear Functions

- **Linear:** polynomial of degree 1.


Example: $y = mx + b$, where m = slope and b = y -intercept.



Linear and Nonlinear Functions

- **Linear:** polynomial of degree 1.

Example: $y = mx + b$, where m = slope and b = y -intercept.




- **Nonlinear:** anything that isn't constant or polynomial of degree 1.

Examples: $y = x^2 + 2x + 1$, $y = \sin(x)$, $y = \ln(x)$, $y = e^x$

Linear and Nonlinear Functions


- **Linear:** polynomial of degree 1.

Example: $y = mx + b$, where m = slope and b = y -intercept.



- **Nonlinear:** anything that isn't constant or polynomial of degree 1.

Examples: $y = x^2 + 2x + 1$, $y = \sin(x)$, $y = \ln(x)$, $y = e^x$



Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Important properties of logarithms:

1. $b^{\log_b x} = x$
2. $\log_b(xy) = \log_b(x) + \log_b(y)$
3. $\log_b(\frac{x}{y}) = \log_b(x) - \log_b(y)$
4. $\log_b(y^x) = x \log_b y$
5. $\log_b(\frac{1}{x}) = -\log_b(x)$

Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Important properties of logarithms:

1. $b^{\log_b x} = x$
2. $\log_b(xy) = \log_b(x) + \log_b(y)$
3. $\log_b(\frac{x}{y}) = \log_b(x) - \log_b(y)$
4. $\log_b(y^x) = x \log_b y$
5. $\log_b(\frac{1}{x}) = -\log_b(x) \quad b^{-x} = \frac{1}{b^x}$

Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Important properties of logarithms:

1. $b^{\log_b x} = x$
2. $\log_b(xy) = \log_b(x) + \log_b(y)$
3. $\log_b(\frac{x}{y}) = \log_b(x) - \log_b(y)$
4. $\log_b(y^x) = x \log_b y$
5. $\log_b(\frac{1}{x}) = -\log_b(x) \quad b^{-x} = \frac{1}{b^x}$
6. $\log_b(1) = 0$

Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Important properties of logarithms:

1. $b^{\log_b x} = x$
2. $\log_b(xy) = \log_b(x) + \log_b(y)$
3. $\log_b(\frac{x}{y}) = \log_b(x) - \log_b(y)$
4. $\log_b(y^x) = x \log_b y$
5. $\log_b(\frac{1}{x}) = -\log_b(x) \quad b^{-x} = \frac{1}{b^x}$
6. $\log_b(1) = 0 \quad b^0 = 1$

Exponents and Logarithms

Definition:

$$(\log_b P = Q) \iff (b^Q = P)$$

Important properties of logarithms:

1. $b^{\log_b x} = x$
2. $\log_b(xy) = \log_b(x) + \log_b(y)$
3. $\log_b\left(\frac{x}{y}\right) = \log_b(x) - \log_b(y)$
4. $\log_b(y^x) = x \log_b y$
5. $\log_b\left(\frac{1}{x}\right) = -\log_b(x) \quad b^{-x} = \frac{1}{b^x}$
6. $\log_b(1) = 0 \quad b^0 = 1$
7. $\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$

Examples

► $\log(\sqrt{10}) =$

Examples

► $\log(\sqrt{10}) = \frac{1}{2}$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) =$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) =$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) =$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$
- ▶ $\ln(1) =$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$
- ▶ $\ln(1) = 0$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$
- ▶ $\ln(1) = 0$
- ▶ $\ln(e) =$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$
- ▶ $\ln(1) = 0$
- ▶ $\ln(e) = 1$

Examples

- ▶ $\log(\sqrt{10}) = \frac{1}{2}$
- ▶ $\log(1) = 0$
- ▶ $\log(10) = 1$
- ▶ $\log(100) = 2$
- ▶ $\ln(1) = 0$
- ▶ $\ln(e) = 1$

Summation Notation

$$\sum_{i=1}^n x_i = x_1 + x_2 + x_3 + \cdots + x_n$$

Summation Notation

$$\sum_{i=1}^n x_i = x_1 + x_2 + x_3 + \cdots + x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\sum_{i=1}^2 x_i$

► $\sum_{i=1}^3 x_i^{y_i}$

► $\sum_{i=2}^3 (x_i + y_{i-1})$

Summation Notation

$$\sum_{i=1}^n x_i = x_1 + x_2 + x_3 + \cdots + x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\sum_{i=1}^2 x_i$

► $\sum_{i=1}^3 x_i^{y_i}$

► $\sum_{i=2}^3 (x_i + y_{i-1})$

► $\sum_{i=1}^2 x_i = 4 + (-2) = 2$

Summation Notation

$$\sum_{i=1}^n x_i = x_1 + x_2 + x_3 + \cdots + x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\sum_{i=1}^2 x_i$

► $\sum_{i=1}^3 x_i^{y_i}$

► $\sum_{i=2}^3 (x_i + y_{i-1})$

► $\sum_{i=1}^2 x_i = 4 + (-2) = 2$

► $\sum_{i=1}^3 x_i^{y_i} = 4^{-1} + (-2)^0 + 3^1 = \frac{1}{4} + 1 + 3 = 4\frac{1}{4}$

Summation Notation

$$\sum_{i=1}^n x_i = x_1 + x_2 + x_3 + \cdots + x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\sum_{i=1}^2 x_i$

► $\sum_{i=1}^3 x_i^{y_i}$

► $\sum_{i=2}^3 (x_i + y_{i-1})$

► $\sum_{i=1}^2 x_i = 4 + (-2) = 2$

► $\sum_{i=1}^3 x_i^{y_i} = 4^{-1} + (-2)^0 + 3^1 = \frac{1}{4} + 1 + 3 = 4\frac{1}{4}$

► $\sum_{i=2}^3 (x_i + y_{i-1}) = (-2 + -1) + (3 + 0) = 0$

Summation Notation

Properties:

$$\blacktriangleright \sum_{i=1}^n cx_i = c \sum_{i=1}^n x_i$$

$$\blacktriangleright \sum_{i=1}^n (x_i + y_i) = \sum_{i=1}^n x_i + \sum_{i=1}^n y_i$$

$$\blacktriangleright \sum_{i=1}^n c = nc$$

Product Notation

$$\prod_{i=1}^n x_i = x_1 x_2 x_3 \cdots x_n$$

Product Notation

$$\prod_{i=1}^n x_i = x_1 x_2 x_3 \cdots x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\prod_{i=1}^2 x_i$

► $\prod_{i=1}^3 x_i^{y_i}$

► $\prod_{i=2}^3 (x_i + y_{i-1})$

Product Notation

$$\prod_{i=1}^n x_i = x_1 x_2 x_3 \cdots x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\prod_{i=1}^2 x_i$

► $\prod_{i=1}^3 x_i^{y_i}$

► $\prod_{i=2}^3 (x_i + y_{i-1})$

► $\prod_{i=1}^2 x_i = 4 \cdot -2 = -8$

Product Notation

$$\prod_{i=1}^n x_i = x_1 x_2 x_3 \cdots x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

► $\prod_{i=1}^2 x_i$

► $\prod_{i=1}^3 x_i^{y_i}$

► $\prod_{i=2}^3 (x_i + y_{i-1})$

► $\prod_{i=1}^2 x_i = 4 \cdot -2 = -8$

► $\prod_{i=1}^3 x_i^{y_i} = 4^{-1} \cdot (-2)^0 \cdot 3^1 = \frac{1}{4} \cdot 1 \cdot 3 = \frac{3}{4}$

Product Notation

$$\prod_{i=1}^n x_i = x_1 x_2 x_3 \cdots x_n$$

Let $x = (4, -2, 3)$, $y = (-1, 0, 1)$. Calculate

▶ $\prod_{i=1}^2 x_i$

▶ $\prod_{i=1}^3 x_i^{y_i}$

▶ $\prod_{i=2}^3 (x_i + y_{i-1})$

▶ $\prod_{i=1}^2 x_i = 4 \cdot -2 = -8$

▶ $\prod_{i=1}^3 x_i^{y_i} = 4^{-1} \cdot (-2)^0 \cdot 3^1 = \frac{1}{4} \cdot 1 \cdot 3 = \frac{3}{4}$

▶ $\prod_{i=2}^3 (x_i + y_{i-1}) = (-2 + -1) \cdot (3 + 0) = -9$

Product Notation

Properties:

$$\blacktriangleright \prod_{i=1}^n cx_i = c^n \prod_{i=1}^n x_i$$

$$\blacktriangleright \prod_{i=1}^n (x_i + y_i) = (x_1 + y_1)(x_2 + y_2) \dots$$

$$\blacktriangleright \prod_{i=1}^n c = c^n$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\log \left(\prod_{i=1}^n cx_i \right)$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\log\left(\prod_{i=1}^n cx_i\right) = \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n)$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\begin{aligned}\log\left(\prod_{i=1}^n cx_i\right) &= \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n) \\ &= \log(cx_1) + \log(cx_2) + \dots + \log(cx_n)\end{aligned}$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\begin{aligned}\log\left(\prod_{i=1}^n cx_i\right) &= \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n) \\ &= \log(cx_1) + \log(cx_2) + \dots + \log(cx_n) \\ &= \sum_{i=1}^n \log(cx_i)\end{aligned}$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\begin{aligned}\log\left(\prod_{i=1}^n cx_i\right) &= \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n) \\ &= \log(cx_1) + \log(cx_2) + \dots + \log(cx_n) \\ &= \sum_{i=1}^n \log(cx_i) \\ &= [(\log c + \log x_1) + (\log c + \log x_2) + \\ &\quad \dots + (\log c + \log x_n)]\end{aligned}$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\begin{aligned}\log\left(\prod_{i=1}^n cx_i\right) &= \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n) \\&= \log(cx_1) + \log(cx_2) + \dots + \log(cx_n) \\&= \sum_{i=1}^n \log(cx_i) \\&= [(\log c + \log x_1) + (\log c + \log x_2) + \\&\quad \dots + (\log c + \log x_n)] \\&= \sum_{i=1}^n \log c + \sum_{i=1}^n \log(x_i)\end{aligned}$$

Sums, Products, and Logs

Logarithms can help us change products into sums.

$$\begin{aligned}\log\left(\prod_{i=1}^n cx_i\right) &= \log(cx_1 \cdot cx_2 \cdot \dots \cdot cx_n) \\&= \log(cx_1) + \log(cx_2) + \dots + \log(cx_n) \\&= \sum_{i=1}^n \log(cx_i) \\&= [(\log c + \log x_1) + (\log c + \log x_2) + \\&\quad \dots + (\log c + \log x_n)] \\&= \sum_{i=1}^n \log c + \sum_{i=1}^n \log(x_i) \\&= n \log(c) + \sum_{i=1}^n \log(x_i)\end{aligned}$$

Vectors

Vectors

- ▶ **Vector:** A vector in n -space is an ordered list of n numbers. These numbers can be represented as either a row vector or a column vector:

$$\mathbf{v} = \begin{pmatrix} v_1 & v_2 & \dots & v_n \end{pmatrix}, \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix}$$

Vectors

- ▶ **Vector:** A vector in n -space is an ordered list of n numbers. These numbers can be represented as either a row vector or a column vector:

$$\mathbf{v} = \begin{pmatrix} v_1 & v_2 & \dots & v_n \end{pmatrix}, \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix}$$

- ▶ We can also think of a vector as defining a point in n -dimensional space, usually \mathbf{R}^n ; each element of the vector defines the coordinate of the point in a particular direction.

Vector Arithmetic

- ▶ **Vector Addition:** Vector addition is defined for two vectors \mathbf{u} and \mathbf{v} iff they have the same number of elements:

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} u_1 + v_1 & u_2 + v_2 & \cdots & u_n + v_n \end{pmatrix}$$

Vector Arithmetic

- **Vector Addition:** Vector addition is defined for two vectors \mathbf{u} and \mathbf{v} iff they have the same number of elements:

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} u_1 + v_1 & u_2 + v_2 & \cdots & u_n + v_n \end{pmatrix}$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

Vector Arithmetic

- **Vector Addition:** Vector addition is defined for two vectors \mathbf{u} and \mathbf{v} iff they have the same number of elements:

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} u_1 + v_1 & u_2 + v_2 & \cdots & u_n + v_n \end{pmatrix}$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$\mathbf{u} + \mathbf{v} =$

Vector Arithmetic

- **Vector Addition:** Vector addition is defined for two vectors \mathbf{u} and \mathbf{v} iff they have the same number of elements:

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} u_1 + v_1 & u_2 + v_2 & \cdots & u_n + v_n \end{pmatrix}$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} 3 + 2 & -2 + 0 & 1 + 1 \end{pmatrix}$$

Vector Arithmetic

- **Vector Addition:** Vector addition is defined for two vectors \mathbf{u} and \mathbf{v} iff they have the same number of elements:

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} u_1 + v_1 & u_2 + v_2 & \cdots & u_n + v_n \end{pmatrix}$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$$\mathbf{u} + \mathbf{v} = \begin{pmatrix} 3 + 2 & -2 + 0 & 1 + 1 \end{pmatrix} = \begin{pmatrix} 5 & -2 & 2 \end{pmatrix}$$

```
u <- c(3, -2, 1)
```

```
v <- c(2, 0, 1)
```

```
u + v
```

```
## [1] 5 -2 2
```

Vector Arithmetic

- **Scalar Multiplication:** The product of a scalar c and vector \mathbf{v} is:

$$c\mathbf{v} = \begin{pmatrix} cv_1 & cv_2 & \dots & cv_n \end{pmatrix}$$

Vector Arithmetic

- **Scalar Multiplication:** The product of a scalar c and vector \mathbf{v} is:

$$c\mathbf{v} = \begin{pmatrix} cv_1 & cv_2 & \dots & cv_n \end{pmatrix}$$

Let $\mathbf{v} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $c = 6$.

Vector Arithmetic

- **Scalar Multiplication:** The product of a scalar c and vector \mathbf{v} is:

$$c\mathbf{v} = \begin{pmatrix} cv_1 & cv_2 & \dots & cv_n \end{pmatrix}$$

Let $\mathbf{v} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $c = 6$.

$$c\mathbf{v} =$$

Vector Arithmetic

- **Scalar Multiplication:** The product of a scalar c and vector \mathbf{v} is:

$$c\mathbf{v} = (cv_1 \quad cv_2 \quad \dots \quad cv_n)$$

Let $\mathbf{v} = (3 \quad -2 \quad 1)$, $c = 6$.

$$c\mathbf{v} = (6 \cdot 3 \quad 6 \cdot -2 \quad 6 \cdot 1)$$

Vector Arithmetic

- **Scalar Multiplication:** The product of a scalar c and vector \mathbf{v} is:

$$c\mathbf{v} = (cv_1 \quad cv_2 \quad \dots \quad cv_n)$$

Let $\mathbf{v} = (3 \quad -2 \quad 1)$, $c = 6$.

$$c\mathbf{v} = (6 \cdot 3 \quad 6 \cdot -2 \quad 6 \cdot 1) = (18 \quad -12 \quad 6)$$

```
c <- 6
```

```
v <- c(3, -2, 1)
```

```
c * v
```

```
## [1] 18 -12 6
```

Vector Properties

Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors, c, d be constants, $\mathbf{0}$ be vector \mathbf{z} s.t.
 $z_i = 0$.

Vector Properties

Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors, c, d be constants, $\mathbf{0}$ be vector \mathbf{z} s.t.
 $z_i = 0$.

Then, assuming $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{0}$ *conformable*,

Vector Properties

Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors, c, d be constants, $\mathbf{0}$ be vector \mathbf{z} s.t. $z_i = 0$.

Then, assuming $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{0}$ *conformable*,

- ▶ Commutativity: $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- ▶ Associativity: $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
- ▶ Distributivity: $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
- ▶ Scalar Distributivity: $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$

Vector Properties

Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors, c, d be constants, $\mathbf{0}$ be vector \mathbf{z} s.t. $z_i = 0$.

Then, assuming $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{0}$ conformable,

- ▶ Commutativity: $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- ▶ Associativity: $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
- ▶ Distributivity: $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
- ▶ Scalar Distributivity: $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
- ▶ Additive Identity: $\mathbf{u} + \mathbf{0} = \mathbf{u}$

Vector Properties

Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors, c, d be constants, $\mathbf{0}$ be vector \mathbf{z} s.t. $z_i = 0$.

Then, assuming $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{0}$ conformable,

- ▶ Commutativity: $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- ▶ Associativity: $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
- ▶ Distributivity: $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
- ▶ Scalar Distributivity: $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
- ▶ Additive Identity: $\mathbf{u} + \mathbf{0} = \mathbf{u}$
- ▶ Scalar Multiplicative Identity: $1\mathbf{u} = \mathbf{u}$

Transpose

- ▶ **Transpose:** The transpose of a vector \mathbf{u} changes \mathbf{u} 's row/column status.

Transpose

- ▶ **Transpose:** The transpose of a vector \mathbf{u} changes \mathbf{u} 's row/column status.

$$\mathbf{u}^T = \mathbf{u}'$$

Transpose

- ▶ **Transpose:** The transpose of a vector \mathbf{u} changes \mathbf{u} 's row/column status.

$$\mathbf{u}^T = \mathbf{u}'$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$ be 1×3 .

Transpose

- **Transpose:** The transpose of a vector \mathbf{u} changes \mathbf{u} 's row/column status.

$$\mathbf{u}^T = \mathbf{u}'$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$ be 1×3 . Then, $\mathbf{u}' = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix}$ is 3×1 .

Inner Product

- **Inner Product:** The Euclidean inner product (also, the “dot product”) of two vectors \mathbf{u} and \mathbf{v} is defined iff they have the same number of elements

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n = \sum_{i=1}^n u_iv_i$$

Inner Product

- **Inner Product:** The Euclidean inner product (also, the “dot product”) of two vectors \mathbf{u} and \mathbf{v} is defined iff they have the same number of elements

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n = \sum_{i=1}^n u_iv_i$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

Inner Product

- **Inner Product:** The Euclidean inner product (also, the “dot product”) of two vectors \mathbf{u} and \mathbf{v} is defined iff they have the same number of elements

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n = \sum_{i=1}^n u_iv_i$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$\mathbf{u} \cdot \mathbf{v} =$

Inner Product

- **Inner Product:** The Euclidean inner product (also, the “dot product”) of two vectors \mathbf{u} and \mathbf{v} is defined iff they have the same number of elements

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n = \sum_{i=1}^n u_iv_i$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$$\mathbf{u} \cdot \mathbf{v} = 3 \cdot 2 + -2 \cdot 0 + 1 \cdot 1$$

Inner Product

- **Inner Product:** The Euclidean inner product (also, the “dot product”) of two vectors \mathbf{u} and \mathbf{v} is defined iff they have the same number of elements

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n = \sum_{i=1}^n u_iv_i$$

Let $\mathbf{u} = \begin{pmatrix} 3 & -2 & 1 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix}$.

$$\mathbf{u} \cdot \mathbf{v} = 3 \cdot 2 + (-2) \cdot 0 + 1 \cdot 1 = 6 + 1 = 7.$$

```
u %*% v %>% drop() # Simplify array dims
```

```
## [1] 7
```


Inner Product and Orthogonality

If $\mathbf{u} \cdot \mathbf{v} = 0$, the vectors are *orthogonal* (or perpendicular).

Let $\mathbf{u} = \begin{pmatrix} 5 & 0 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 0 & -2 \end{pmatrix}$.

Inner Product and Orthogonality

If $\mathbf{u} \cdot \mathbf{v} = 0$, the vectors are *orthogonal* (or perpendicular).

Let $\mathbf{u} = \begin{pmatrix} 5 & 0 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 0 & -2 \end{pmatrix}$.

$$\mathbf{u} \cdot \mathbf{v} = 5 \cdot 0 + 0 \cdot -2 = 0 + 0 = 0.$$

Inner Product

- Think about the $\mathbf{u} \cdot \mathbf{v}$ inner product as

$$\underbrace{\mathbf{u}}_{1 \times k} \cdot \underbrace{\mathbf{v}}_{k \times 1} = \underbrace{w}_{1 \times 1}$$

Inner Product

- Think about the $\mathbf{u} \cdot \mathbf{v}$ inner product as

$$\underbrace{\mathbf{u}}_{1 \times k} \cdot \underbrace{\mathbf{v}}_{k \times 1} = \underbrace{w}_{1 \times 1}$$

- Or, assume \mathbf{u}, \mathbf{v} both $k \times 1$ columns, then $\mathbf{u} \cdot \mathbf{v} \equiv \mathbf{v}^T \mathbf{u}$

Inner Product

- Think about the $\mathbf{u} \cdot \mathbf{v}$ inner product as

$$\underbrace{\mathbf{u}}_{1 \times k} \cdot \underbrace{\mathbf{v}}_{k \times 1} = \underbrace{w}_{1 \times 1}$$

- Or, assume \mathbf{u}, \mathbf{v} both $k \times 1$ columns, then $\mathbf{u} \cdot \mathbf{v} \equiv \mathbf{v}^T \mathbf{u}$

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{v}' \mathbf{u} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix}$$

Inner Product

- Think about the $\mathbf{u} \cdot \mathbf{v}$ inner product as

$$\underbrace{\mathbf{u}}_{1 \times k} \cdot \underbrace{\mathbf{v}}_{k \times 1} = \underbrace{w}_{1 \times 1}$$

- Or, assume \mathbf{u}, \mathbf{v} both $k \times 1$ columns, then $\mathbf{u} \cdot \mathbf{v} \equiv \mathbf{v}^T \mathbf{u}$

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{v}' \mathbf{u} = \begin{pmatrix} 2 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} = 7$$

Inner Product Properties

- ▶ Commutativity: $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
- ▶ Associativity: $c(\mathbf{u} \cdot \mathbf{v}) = (c\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (c\mathbf{v})$
- ▶ Distributivity: $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$
- ▶ Zero Product: $\mathbf{u} \cdot \mathbf{0} = 0$

Vector Norm

- ▶ **Vector Norm:** The *norm* of a vector measures its length. There are many norms; most common: Euclidean norm (corresponding to usual conception of distance in 3D space):

Vector Norm

- **Vector Norm:** The *norm* of a vector measures its length. There are many norms; most common: Euclidean norm (corresponding to usual conception of distance in 3D space):

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1v_1 + v_2v_2 + \cdots + v_nv_n}$$

Vector Norm

- **Vector Norm:** The *norm* of a vector measures its length. There are many norms; most common: Euclidean norm (corresponding to usual conception of distance in 3D space):

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1v_1 + v_2v_2 + \cdots + v_nv_n}$$

$$\|\mathbf{v}\| = \left\| \begin{pmatrix} 2 & 0 & 1 \end{pmatrix} \right\|$$

Vector Norm

- **Vector Norm:** The *norm* of a vector measures its length. There are many norms; most common: Euclidean norm (corresponding to usual conception of distance in 3D space):

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1v_1 + v_2v_2 + \cdots + v_nv_n}$$

$$\begin{aligned}\|\mathbf{v}\| &= \left\| \begin{pmatrix} 2 & 0 & 1 \end{pmatrix} \right\| \\ &= \sqrt{2^2 + 0^2 + 1^2}\end{aligned}$$

Vector Norm

- **Vector Norm:** The *norm* of a vector measures its length. There are many norms; most common: Euclidean norm (corresponding to usual conception of distance in 3D space):

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1v_1 + v_2v_2 + \cdots + v_nv_n}$$

$$\begin{aligned}\|\mathbf{v}\| &= \left\| \begin{pmatrix} 2 & 0 & 1 \end{pmatrix} \right\| \\ &= \sqrt{2^2 + 0^2 + 1^2} \\ &= \sqrt{5}\end{aligned}$$

Vector Norm Properties

- ▶ Scalar Multiplication: $\|c\mathbf{u}\| = |c| \cdot \|\mathbf{u}\|$
- ▶ Vector Distance: $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|$
- ▶ Norm Squared: $\|\mathbf{u}\|^2 = \mathbf{u} \cdot \mathbf{u}$
- ▶ Cosine Rule: $\cos(\theta) = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \cdot \|\mathbf{v}\|}$
- ▶ Difference Norm: $\begin{aligned} \|\mathbf{u} - \mathbf{v}\|^2 &= \|\mathbf{u}\|^2 - 2(\mathbf{u} \cdot \mathbf{v}) + \|\mathbf{v}\|^2 \\ &= \|\mathbf{u}\|^2 - 2\|\mathbf{u}\| \cdot \|\mathbf{v}\|(\cos \theta) + \|\mathbf{v}\|^2 \end{aligned}$
- ▶ Triangle Inequality: $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$
- ▶ Cauchy-Schwartz Inequality: $\|\mathbf{u} \cdot \mathbf{v}\| \leq \|\mathbf{u}\| \cdot \|\mathbf{v}\|$

Mahalanobis Distance

$$MD_{ij} = \sqrt{(\mathbf{x}_i - \mathbf{x}_j)' \Sigma^{-1} (\mathbf{x}_i - \mathbf{x}_j)}$$

Mahalanobis Distance

$$MD_{ij} = \sqrt{(\mathbf{x}_i - \mathbf{x}_j)' \Sigma^{-1} (\mathbf{x}_i - \mathbf{x}_j)}$$

Like Euclidean distance, but scaled by inverse covariances

$$\hat{\Sigma} = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})'$$

where

$$\bar{\mathbf{x}} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i$$

is the vector of sample means.

A Bit of R

```
a <- c(3, 0, 0)
b <- c(0, 2, 0)
a %*% b ## inner prod
```

```
##      [,1]
## [1,]    0
```


Dependence and Independence

- ▶ **Linear combinations:** The vector \mathbf{u} is a linear combination of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ if

$$\mathbf{u} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k$$

Dependence and Independence

- ▶ **Linear combinations:** The vector \mathbf{u} is a linear combination of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ if

$$\mathbf{u} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k$$

- ▶ **Linear independence:** A set of vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ is linearly independent if the only solution to the equation

$$c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k = \mathbf{0}$$

is $c_1 = c_2 = \dots = c_k = 0$. If another solution exists, the set of vectors is linearly dependent.

Linear Dependence

- ▶ A set S of vectors is linearly dependent iff at least one of the vectors in S can be written as a linear combination of the other vectors in S .
- ▶ Linear independence is only defined for sets of vectors with the same number of elements
- ▶ Any linearly independent set of vectors in n -space contains at most n vectors.

Example 1

Are the following sets of vectors linearly independent?

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

Example 1

Are the following sets of vectors linearly independent?

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

Yes. $(c_3 = 0) \Rightarrow (c_2 = 0) \Rightarrow (c_1 = 0)$

Example 2

Are the following sets of vectors linearly independent?

$$\mathbf{v}_1 = \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} -2 \\ 2 \\ 4 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix}$$

Example 2

Are the following sets of vectors linearly independent?

$$\mathbf{v}_1 = \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} -2 \\ 2 \\ 4 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix}$$

No. $\mathbf{c} = \begin{pmatrix} 2 & 1 & -2 \end{pmatrix}$ (e.g.)

Matrix Algebra

Matrices

- **Matrix:** A matrix is an array of mn real numbers arranged in m rows by n columns.

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

Matrices

- ▶ Vectors are special cases of matrices
 - ▶ Col vector of length k is a $k \times 1$ matrix
 - ▶ Row vector of length k is a $1 \times k$ matrix

Matrices

- ▶ Vectors are special cases of matrices
 - ▶ Col vector of length k is a $k \times 1$ matrix
 - ▶ Row vector of length k is a $1 \times k$ matrix
- ▶ Think of larger matrices as made up of row or column vectors. E.g.,

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{pmatrix}$$

Matrices

- ▶ Vectors are special cases of matrices
 - ▶ Col vector of length k is a $k \times 1$ matrix
 - ▶ Row vector of length k is a $1 \times k$ matrix
- ▶ Think of larger matrices as made up of row or column vectors. E.g.,

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \\ \cdots \\ \mathbf{b}_n \end{pmatrix}$$

}

Special Matrices

► Identity: $\mathbf{I}_n = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$

► $\mathbf{J}_n = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & \ddots & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix}$

► Zero: $\mathbf{0} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}$

Special Matrices

► Diagonal:
$$\begin{pmatrix} a_{11} & 0 & 0 & 0 \\ 0 & a_{22} & 0 & 0 \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{pmatrix}$$

► Lower Triangular:
$$\begin{pmatrix} a_{11} & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 \\ a_{31} & a_{32} & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}$$

► Upper Triangular:
$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{pmatrix}$$

► Triangular: Either upper triangular or lower triangular

Matrix Equality

- Let \mathbf{A} and \mathbf{B} be two $m \times n$ matrices. Then

$$\mathbf{A} = \mathbf{B}$$

iff

$$a_{ij} = b_{ij}$$

$$\forall i \in \{1, \dots, m\}, \quad j \in \{1, \dots, n\}$$

Matrix Addition

- ▶ Let \mathbf{A} and \mathbf{B} be two $m \times n$ matrices. Then

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{pmatrix}$$

- ▶ \mathbf{A} and \mathbf{B} must be same size – *conformable* for addition

Matrix Addition Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \end{pmatrix}$$

$$\mathbf{A} + \mathbf{B} =$$

Matrix Addition Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 1 & 2 & 1 \\ 2 & 1 & 2 \end{pmatrix}$$

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} 2 & 4 & 4 \\ 6 & 6 & 8 \end{pmatrix}$$

A + B

```
##          [,1] [,2] [,3]
## [1,]      2    4    4
## [2,]      6    6    8
```

Scalar Multiplication

Scalar Multiplication: Given scalar c , the scalar multiplication $c\mathbf{A}$ is

$$c\mathbf{A} = c \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} = \begin{pmatrix} ca_{11} & ca_{12} & \cdots & ca_{1n} \\ ca_{21} & ca_{22} & \cdots & ca_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ ca_{m1} & ca_{m2} & \cdots & ca_{mn} \end{pmatrix}$$

Scalar Multiplication Example

$$c = 2 \quad \mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$$

$$c\mathbf{A} =$$

Scalar Multiplication Example

$$c = 2 \quad \mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$$

$$c\mathbf{A} = \begin{pmatrix} 2 & 4 & 6 \\ 8 & 10 & 12 \end{pmatrix}$$

```
c <- 2
```

```
c * A
```

```
##      [,1] [,2] [,3]  
## [1,]    2    4    6  
## [2,]    8   10   12
```

Matrix Multiplication

- ▶ **Matrix Multiplication:** If \mathbf{A} is $m \times k$ and \mathbf{B} is $k \times n$, then their product $\mathbf{C} = \mathbf{AB}$ is $m \times n$ matrix where

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ik}b_{kj}$$

Matrix Multiplication

- ▶ **Matrix Multiplication:** If \mathbf{A} is $m \times k$ and \mathbf{B} is $k \times n$, then their product $\mathbf{C} = \mathbf{AB}$ is $m \times n$ matrix where

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ik}b_{kj}$$

- ▶ Consider \mathbf{A} to be composed of stacked rows

$$\mathbf{a}_i = \begin{pmatrix} a_{i1} & a_{i2} & \cdots & a_{in} \end{pmatrix},$$

$$\mathbf{B} \text{ to be composed of stacked columns } \mathbf{b}_j = \begin{pmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{mj} \end{pmatrix}.$$

Then, $\mathbf{AB} = \mathbf{C}$, where

$$c_{ij} = \mathbf{a}_i \cdot \mathbf{b}_j$$

Notes on Matrix Multiplication

- ▶ To be *conformable* for multiplication, number of cols of first matrix must equal number of rows of second matrix

Notes on Matrix Multiplication

- ▶ To be *conformable* for multiplication, number of cols of first matrix must equal number of rows of second matrix
- ▶ The sizes of the matrices and product must be

$$(m \times k)(k \times n) = (m \times n)$$

Notes on Matrix Multiplication

- ▶ To be *conformable* for multiplication, number of cols of first matrix must equal number of rows of second matrix
- ▶ The sizes of the matrices and product must be

$$(m \times k)(k \times n) = (m \times n)$$

- ▶ Given \mathbf{AB} , say \mathbf{B} is pre-multiplied by \mathbf{A} or \mathbf{B} is left-multiplied by \mathbf{A} or \mathbf{A} is post-multiplied by \mathbf{B} or \mathbf{A} is right-multiplied by \mathbf{B}

Warning!

- ▶ Commutative law for multiplication does **not** hold – order of multiplication matters:

$$\mathbf{AB} \neq \mathbf{BA}$$

- ▶ \mathbf{AB} may exist, while \mathbf{BA} does not.

Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{AB} =$$

Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{AB} = \begin{pmatrix} 2 & 3 \\ -2 & 2 \end{pmatrix}$$

Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{AB} = \begin{pmatrix} 2 & 3 \\ -2 & 2 \end{pmatrix}$$

$$\mathbf{BA} =$$

Example

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{AB} = \begin{pmatrix} 2 & 3 \\ -2 & 2 \end{pmatrix}$$

$$\mathbf{BA} = \begin{pmatrix} 1 & 7 \\ -1 & 3 \end{pmatrix}$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T =$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .
- ▶ Examples:

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T =$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .
- ▶ Examples:

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T =$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .
- ▶ Examples:

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T = \begin{pmatrix} 4 & 0 \\ -2 & 5 \\ 3 & -1 \end{pmatrix}$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .
- ▶ Examples:

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T = \begin{pmatrix} 4 & 0 \\ -2 & 5 \\ 3 & -1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \quad \mathbf{B}^T =$$

Transpose

- ▶ **Transpose:** The transpose of the $m \times n$ matrix \mathbf{A} is $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}') obtained by interchanging rows and columns of \mathbf{A} .

- ▶ Examples:

$$\mathbf{A} = \begin{pmatrix} 4 & -2 & 3 \\ 0 & 5 & -1 \end{pmatrix} \quad \mathbf{A}^T = \begin{pmatrix} 4 & 0 \\ -2 & 5 \\ 3 & -1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \quad \mathbf{B}^T = (2 \quad -1 \quad 3)$$

Example Property

Example of $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$:

Example Property

Example of $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$:

$$\mathbf{A} = \begin{pmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{pmatrix}$$

$$(\mathbf{AB})^T =$$

Example Property

Example of $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$:

$$\mathbf{A} = \begin{pmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{pmatrix}$$

$$(\mathbf{AB})^T =$$

$$\left[\begin{pmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{pmatrix} \right]^T = \begin{pmatrix} 12 & 7 \\ 5 & -3 \end{pmatrix}$$

Example Continued

$$\mathbf{B}^T \mathbf{A}^T =$$

Example Continued

$$\mathbf{B}^T \mathbf{A}^T =$$

$$\begin{pmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{pmatrix} =$$

Example Continued

$$\mathbf{B}^T \mathbf{A}^T =$$

$$\begin{pmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{pmatrix} =$$
$$\begin{pmatrix} 12 & 7 \\ 5 & -3 \end{pmatrix}$$

Identity Matrix

The $n \times n$ identity matrix \mathbf{I}_n has diagonal elements = 1 and off-diagonal elements = 0.

Examples:

$$\mathbf{I}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \mathbf{I}_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Systems of Linear Equations

Linear Equations

- ▶ **Linear Equation:** $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$
 a_i are parameters or coefficients. x_i are variables or unknowns.

Linear Equations

- ▶ **Linear Equation:** $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$
 a_i are parameters or coefficients. x_i are variables or unknowns.
- ▶ “Linear”: only one variable per term and degree at most 1.

Linear Equations

- ▶ **Linear Equation:** $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$
 a_i are parameters or coefficients. x_i are variables or unknowns.
- ▶ “Linear”: only one variable per term and degree at most 1.
 1. \mathbb{R}^2 : line
$$x_2 = \frac{b}{a_2} - \frac{a_1}{a_2}x_1$$

Linear Equations

- ▶ **Linear Equation:** $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$
 a_i are parameters or coefficients. x_i are variables or unknowns.
- ▶ “Linear”: only one variable per term and degree at most 1.
 1. \mathbb{R}^2 : line
$$x_2 = \frac{b}{a_2} - \frac{a_1}{a_2}x_1$$
 2. \mathbb{R}^3 : plane
$$x_3 = \frac{b}{a_3} - \frac{a_1}{a_3}x_1 - \frac{a_2}{a_3}x_2$$

Linear Equations

- ▶ **Linear Equation:** $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$
 a_i are parameters or coefficients. x_i are variables or unknowns.
- ▶ “Linear”: only one variable per term and degree at most 1.
 1. \mathbb{R}^2 : line
$$x_2 = \frac{b}{a_2} - \frac{a_1}{a_2}x_1$$
 2. \mathbb{R}^3 : plane
$$x_3 = \frac{b}{a_3} - \frac{a_1}{a_3}x_1 - \frac{a_2}{a_3}x_2$$
 3. \mathbb{R}^n : hyperplane

Systems of Linear Equations

Often interested in solving linear systems like

More Systems of Linear Equations

- More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

More Systems of Linear Equations

- More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

- This scalar system is equivalent to the matrix equation

$$\mathbf{Ax} = \mathbf{b}$$

More Systems of Linear Equations

- More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

- This scalar system is equivalent to the matrix equation

$$\mathbf{Ax} = \mathbf{b}$$

- A **solution** to a linear system of m equations in n unknowns is a set of n numbers x_1, x_2, \dots, x_n that satisfy each of the m equations.

More Systems of Linear Equations

- ▶ More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

- ▶ This scalar system is equivalent to the matrix equation

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ A **solution** to a linear system of m equations in n unknowns is a set of n numbers x_1, x_2, \dots, x_n that satisfy each of the m equations.
 1. \mathbb{R}^2 : intersection of the lines.

More Systems of Linear Equations

- More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

- This scalar system is equivalent to the matrix equation

$$\mathbf{Ax} = \mathbf{b}$$

- A **solution** to a linear system of m equations in n unknowns is a set of n numbers x_1, x_2, \dots, x_n that satisfy each of the m equations.
 1. \mathbb{R}^2 : intersection of the lines.
 2. \mathbb{R}^3 : intersection of the planes.

More Systems of Linear Equations

- More generally, we might have a system of m equations in n unknowns

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

- This scalar system is equivalent to the matrix equation

$$\mathbf{Ax} = \mathbf{b}$$

- A **solution** to a linear system of m equations in n unknowns is a set of n numbers x_1, x_2, \dots, x_n that satisfy each of the m equations.
 1. \mathbb{R}^2 : intersection of the lines.
 2. \mathbb{R}^3 : intersection of the planes.
 3. \mathbb{R}^n : intersection of the hyperplanes.

How many Solutions?

- ▶ Does a linear system have one, no, or multiple solutions?
For a system of 2 equations in 2 unknowns (i.e., two lines):

How many Solutions?

- ▶ Does a linear system have one, no, or multiple solutions?
For a system of 2 equations in 2 unknowns (i.e., two lines):
 1. One solution: lines intersect at exactly one point.

How many Solutions?

- ▶ Does a linear system have one, no, or multiple solutions?
For a system of 2 equations in 2 unknowns (i.e., two lines):
 1. One solution: lines intersect at exactly one point.
 2. No solution: lines are parallel.

How many Solutions?

- ▶ Does a linear system have one, no, or multiple solutions?
For a system of 2 equations in 2 unknowns (i.e., two lines):
 1. One solution: lines intersect at exactly one point.
 2. No solution: lines are parallel.
 3. Infinite solutions: lines coincide.

Matrices to Represent Linear Systems

Matrices are an efficient way to represent linear systems such as

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

Matrices to Represent Linear Systems

Matrices are an efficient way to represent linear systems such as

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & & & \vdots & & & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

as $\mathbf{Ax} = \mathbf{b}$

Coefficient Matrix

The $m \times n$ **coefficient matrix** \mathbf{A} is an array of mn real numbers arranged in m rows by n columns:

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

Variable & Output Vectors

- ▶ The unknown quantities are represented by the vector

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Variable & Output Vectors

- ▶ The unknown quantities are represented by the vector

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

- ▶ The RHS of the linear system is represented by the vector

$$\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}.$$

Inverse of a Matrix

- ▶ An $n \times n$ matrix \mathbf{A} is *nonsingular* or *invertible* if there exists an $n \times n$ matrix \mathbf{A}^{-1} such that

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n$$

Inverse of a Matrix

- ▶ An $n \times n$ matrix \mathbf{A} is *nonsingular* or *invertible* if there exists an $n \times n$ matrix \mathbf{A}^{-1} such that

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n$$

- ▶ Then, \mathbf{A}^{-1} is the inverse of \mathbf{A} .

Inverse of a Matrix

- ▶ An $n \times n$ matrix \mathbf{A} is *nonsingular* or *invertible* if there exists an $n \times n$ matrix \mathbf{A}^{-1} such that

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n$$

- ▶ Then, \mathbf{A}^{-1} is the inverse of \mathbf{A} .
- ▶ If there is no such \mathbf{A}^{-1} , then \mathbf{A} is *singular* or *noninvertible*.

Example of Inverses

► Example: Let

$$\mathbf{A} = \begin{pmatrix} 2 & 3 \\ 2 & 2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} -1 & \frac{3}{2} \\ 1 & -1 \end{pmatrix}$$

Example of Inverses

► Example: Let

$$\mathbf{A} = \begin{pmatrix} 2 & 3 \\ 2 & 2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} -1 & \frac{3}{2} \\ 1 & -1 \end{pmatrix}$$

Since

$$\mathbf{AB} = \mathbf{BA} = \mathbf{I}_n$$

we conclude that \mathbf{B} is the inverse of \mathbf{A} , \mathbf{A}^{-1} , and that \mathbf{A} is nonsingular.

Calculating the Inverse in R

```
A <- matrix(c(1, 4, 5, 6), 2, 2, byrow = TRUE)
```

```
A
```

```
##      [,1] [,2]  
## [1,]    1    4  
## [2,]    5    6
```


Calculating the Inverse in R

```
A <- matrix(c(1, 4, 5, 6), 2, 2, byrow = TRUE)
```

```
A
```

```
##      [,1] [,2]
```

```
## [1,]    1    4
```

```
## [2,]    5    6
```

```
solve(A)
```

```
##      [,1]      [,2]
```

```
## [1,] -0.4285714  0.28571429
```

```
## [2,]  0.3571429 -0.07142857
```

Calculating the Inverse in R

```
solve(A) %*% A
```

```
##                [,1] [,2]  
## [1,]  1.000000e+00    0  
## [2,] -5.551115e-17    1
```

Calculating the Inverse in R

```
solve(A) %*% A
```

```
##           [,1] [,2]  
## [1,] 1.000000e+00 0  
## [2,] -5.551115e-17 1
```

```
A %*% solve(A)
```

```
##           [,1] [,2]  
## [1,] 1 -5.551115e-17  
## [2,] 0 1.000000e+00
```

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.
- ▶ Suppose \mathbf{A} is nonsingular $\implies \mathbf{A}^{-1}$ exists.

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.
- ▶ Suppose \mathbf{A} is nonsingular $\implies \mathbf{A}^{-1}$ exists.
- ▶ To solve this system, we can premultiply each side by \mathbf{A}^{-1} and simplify:

$$\mathbf{A}^{-1}(\mathbf{Ax}) = \mathbf{A}^{-1}\mathbf{b}$$

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.
- ▶ Suppose \mathbf{A} is nonsingular $\implies \mathbf{A}^{-1}$ exists.
- ▶ To solve this system, we can premultiply each side by \mathbf{A}^{-1} and simplify:

$$\begin{aligned}\mathbf{A}^{-1}(\mathbf{Ax}) &= \mathbf{A}^{-1}\mathbf{b} \\ (\mathbf{A}^{-1}\mathbf{A})\mathbf{x} &= \mathbf{A}^{-1}\mathbf{b}\end{aligned}$$

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.
- ▶ Suppose \mathbf{A} is nonsingular $\implies \mathbf{A}^{-1}$ exists.
- ▶ To solve this system, we can premultiply each side by \mathbf{A}^{-1} and simplify:

$$\begin{aligned}\mathbf{A}^{-1}(\mathbf{Ax}) &= \mathbf{A}^{-1}\mathbf{b} \\ (\mathbf{A}^{-1}\mathbf{A})\mathbf{x} &= \mathbf{A}^{-1}\mathbf{b} \\ \mathbf{I}_n\mathbf{x} &= \mathbf{A}^{-1}\mathbf{b}\end{aligned}$$

Linear Systems and Inverses

- ▶ Matrix representation of a linear system

$$\mathbf{Ax} = \mathbf{b}$$

- ▶ If \mathbf{A} is an $n \times n$ matrix, then $\mathbf{Ax} = \mathbf{b}$ is a system of n equations in n unknowns.
- ▶ Suppose \mathbf{A} is nonsingular $\implies \mathbf{A}^{-1}$ exists.
- ▶ To solve this system, we can premultiply each side by \mathbf{A}^{-1} and simplify:

$$\mathbf{A}^{-1}(\mathbf{Ax}) = \mathbf{A}^{-1}\mathbf{b}$$

$$(\mathbf{A}^{-1}\mathbf{A})\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$

$$\mathbf{I}_n\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$

$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)
- ▶ Then

$$\mathbf{Xb} = \mathbf{y}$$

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)
- ▶ Then

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

- ▶ Since usually $n \gg k$, \mathbf{X} not square

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)

- ▶ Then

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

- ▶ Since usually $n \gg k$, \mathbf{X} not square
- ▶ To isolate \mathbf{b} , how to make premultiplying matrix square?

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)

- ▶ Then

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

- ▶ Since usually $n \gg k$, \mathbf{X} not square
- ▶ To isolate \mathbf{b} , how to make premultiplying matrix square?
- ▶ Observe:

$$\underset{(n \times k)}{\mathbf{X}} \underset{(k \times n)}{\mathbf{X}'} = \underset{(n \times n)}{\quad}$$

can't premultiply \mathbf{b} .

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)

- ▶ Then

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

- ▶ Since usually $n \gg k$, \mathbf{X} not square
- ▶ To isolate \mathbf{b} , how to make premultiplying matrix square?
- ▶ Observe:

$$\underset{(n \times k)}{\mathbf{X}} \underset{(k \times n)}{\mathbf{X}'} = \underset{(n \times n)}{\quad}$$

can't premultiply \mathbf{b} .

The Linear Regression Parameters

- ▶ We have linear systems with n obs and k predictors
 - ▶ \mathbf{X} is $n \times k$ matrix of predictor variable data
 - ▶ \mathbf{y} is $n \times 1$ vector of outcome variable values
 - ▶ \mathbf{b} is $k \times 1$ vector of linear parameters (β 's)

- ▶ Then

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

- ▶ Since usually $n \gg k$, \mathbf{X} not square
- ▶ To isolate \mathbf{b} , how to make premultiplying matrix square?
- ▶ Observe:

$$\underset{(n \times k)}{\mathbf{X}} \underset{(k \times n)}{\mathbf{X}'} = \underset{(n \times n)}{\quad}$$

can't premultiply \mathbf{b} . But this can:

$$\underset{(k \times n)}{\mathbf{X}'} \underset{(n \times k)}{\mathbf{X}} = \underset{(k \times k)}{\quad}$$

The Linear Regression Parameters

$$\mathbf{X}\mathbf{b} = \mathbf{y}$$

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ?

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

$$(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\mathbf{b} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$$

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

$$\begin{aligned}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{I}_k\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}\end{aligned}$$

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

$$\begin{aligned}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{I}_k\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}\end{aligned}$$

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

$$\begin{aligned}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{I}_k\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}\end{aligned}$$

Befriend $(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$.

The Linear Regression Parameters

$$\begin{aligned}\mathbf{X}\mathbf{b} &= \mathbf{y} \\ \mathbf{X}'\mathbf{X}\mathbf{b} &= \mathbf{X}'\mathbf{y}\end{aligned}$$

How to isolate \mathbf{b} ? Multiply by an inverse.

$$\begin{aligned}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{I}_k\mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} \\ \mathbf{b} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}\end{aligned}$$

Befriend $(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$. If you understand it, its cousins, and their properties (both strengths and weaknesses), your data-analytic future will be bright.

Solving a System in R

$$\mathbf{y} = \mathbf{X}\mathbf{b}$$

```
x1 <- c(1, 3, 5)
x2 <- c(3, 1, 2)
x3 <- c(1, 1, 1)
y <- 4 * x1 + 3 * x2 + x3
(X <- cbind(x1, x2, x3))
```

```
##      x1 x2 x3
## [1,]  1  3  1
## [2,]  3  1  1
## [3,]  5  2  1
```

Solving a System in R

```
Xinv <- solve(X)
b <- Xinv %*% y
b
```

```
##      [,1]
## x1      4
## x2      3
## x3      1
```