

Anatomically-Constrained LDMs for Rare Case Augmentation



SAPIENZA
UNIVERSITÀ DI ROMA

Context and objective:


Periodontitis is a severe gum infection that destroys the soft tissue and bone supporting the teeth, representing one of the leading causes of tooth loss worldwide

Goal:

LDM



Anatomically-
Constrained
LDMs




Downstream

Ablation study

Data organisation

Requirement : Keypoints in the image to allow us to constrain the LMD anatomically

One major issue → only 193 samples



Dataset already used in Periodontal Bone Loss Analysis via Keypoint Detection With Heuristic Post-Processing, Ryan Banks


Our models:

LDM

- Standard LDM without spatial guidance
- Baseline for comparing the impact of anatomical constraints
- Generated using only random noise


LDM with BBOX

- LDM guided by a simple binary mask
- Low spatial constraint (approximate region of the tooth)



LDM with gaussian heatmap

- Guided by a continuous heatmap (0 to 1)
- Precise anatomical guidance (CEJ points, bone level, apex)



Evaluation:

- Three models and 193 samples

Downstream


- Compare performance with and without synthetic data
- Metrics: mAP, AP (Severe)

Ablation Study

- Analyze the influence of the anatomical loss weight
- Identify the most effective design choices

Downstream:


impact of the increase in data on the mAP



rate of increase for severe patches


Ablation Study:

Impact of anatomic weight on FID



α anatomic weight

Best FID for each model



Conclusion

Conclusion

Terminons avec optimisme – la biotech façonne un monde meilleur, guidée par l'éthique et la responsabilité.

Rejoignez la révolution

+123-456-7890
@reallygreatsite
www.reallygreatsite.com
hello@reallygreatsite.com

