Geometric approaches to solving Diophantine equations

Alex J. Best

Tomorrows Mathematicians Today 2013

16-2-2013

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

Kational points on surfaces



Introduction

Named for Diophantus of Alexandria ($\approx 250 AD$) Some Diophantine equations: Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry o Numbers

ational points on urfaces



Introduction

Named for Diophantus of Alexandria ($\approx 250 AD$) Some Diophantine equations:

$$x^2 + y^2 = z^2$$

(Pythagorean triples)

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

cational points on surfaces



$$x^2 + y^2 = z^2$$

(Pythagorean triples)

$$x^2 - ny^2 = 1$$

(Pell's equation)

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

surfaces

$$x^2 + y^2 = z^2$$

(Pythagorean triples)

$$x^2 - ny^2 = 1$$

(Pell's equation)

$$x^3 + 48 = y^4$$

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

surfaces



Named for Diophantus of Alexandria ($\approx 250 AD$) Some Diophantine equations:

$$x^2 + y^2 = z^2$$

(Pythagorean triples)

$$x^2 - ny^2 = 1$$

(Pell's equation)

$$x^3 + 48 = y^4$$

$$9^x - 8^y = 1$$

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Kational points on surfaces

► Are there any solutions?

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry o Numbers

> Kational points on surfaces



Geometric
approaches to
solving
Diophantine
equations

Alex J. Best

Introduction

Numbers

surfaces

- ► Are there any solutions?
- ► How many are there?

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

Kational points on surfaces

- Are there any solutions?
- How many are there?
- ► Can we classify them?

Geometric
approaches to
solving
Diophantine
equations

Alex J. Best

Introduction

Numbers

Kational points on surfaces

- Are there any solutions?
- How many are there?
- ► Can we classify them?
- ► How can we compute them?

In this talk

Two general ideas:

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

tational points on urfaces

In this talk

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry o Numbers

> Rational points on surfaces

Two general ideas:

► Geometry of numbers

In this talk

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry o Numbers

> Rational points on surfaces

Two general ideas:

- ► Geometry of numbers
- Rational points on surfaces

Minkowski and the Geometry of Numbers

1910 - Hermann Minkowski publishes his paper "Geometrie der Zahlen" and sparks a new field called the Geometry of Numbers.





Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Kational points on surfaces



A motivating example

We want to find when integers x, y such that $x^2 + y^2 = p$ where p is prime.

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on aurfaces

surfaces

We want to find when integers x, y such that $x^2 + y^2 = p$ where p is prime.

We can try a few primes, and notice that $2^2 + 1^2 = 5$, $3^2 + 2^2 = 13$, $4^2 + 1^2 = 17$, ... all work.

We want to find when integers x, y such that $x^2 + y^2 = p$ where p is prime.

We can try a few primes, and notice that $2^2 + 1^2 = 5$, $3^2 + 2^2 = 13$, $4^2 + 1^2 = 17$, ... all work. But 7, 11, 19, ... do not.

We want to find when integers x, y such that $x^2 + y^2 = p$ where p is prime.

We can try a few primes, and notice that $2^2 + 1^2 = 5$, $3^2 + 2^2 = 13$, $4^2 + 1^2 = 17$, ... all work.


But 7, 11, 19, ... do not.

So maybe $x^2 + y^2 = p$ when $p \equiv 1 \pmod{4}$.

Theorem (Fermat's Christmas Theorem)

An odd prime p can be written as $p = x^2 + y^2$ if and only if $p \equiv 1 \pmod{4}$.

When p = 5 we can look at points satisfying our criteria:



Geometric approaches to solving Diophantine equations

Alex J. Best


Introduction

Geometry of Numbers

surfaces



When p = 5 we can look at points satisfying our criteria:




Geometric approaches to solving Diophantine equations

Alex J. Best

Geometry of Numbers



When p = 5 we can look at points satisfying our criteria:




Geometric approaches to solving Diophantine equations

Alex J. Best

Geometry of Numbers



When p = 5 we can look at points satisfying our criteria:



Geometric approaches to solving Diophantine equations

Alex J. Best

Geometry of Numbers



Minkowski's first theorem

We say a set B in \mathbb{R}^n is symmetric if $x \in B \implies -x \in B$. We say a set B is \mathbb{R}^n is convex if $x, y \in B \implies x + \lambda(y - x) \in B$ for $0 \le \lambda \le 1$. Geometric approaches to solving Diophantine equations

Alex J. Best

Introduct

Geometry of Numbers

Rational points on surfaces

We say a set B in \mathbb{R}^n is symmetric if $x \in B \implies -x \in B$. We say a set B is \mathbb{R}^n is convex if $x, y \in B \implies x + \lambda(y - x) \in B$ for $0 \le \lambda \le 1$.

Theorem (Minkowski)

If B is a convex symmetric body and Λ a lattice in \mathbb{R}^n then B contains a non-zero point of the lattice if:

$$Vol(B) > 2^d i$$

Where i is the area of a single cell of the lattice Λ . We can find it using determinants, or the order of our lattice as a subgroup of \mathbb{Z}^n for the more algebraically inclined.

Let's apply it

A circle in \mathbb{R}^2 is symmetric and convex, great!

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

surfaces

Let's apply it

A circle in \mathbb{R}^2 is symmetric and convex, great! The area of our lattice cell is p.

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on surfaces

Let's apply it

A circle in \mathbb{R}^2 is symmetric and convex, great! The area of our lattice cell is p. The area of our circle is $\pi 2p$

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on surfaces

A circle in \mathbb{R}^2 is symmetric and convex, great! The area of our lattice cell is p. The area of our circle is $\pi 2p$ So Minkowski tells us that as:

$$\pi 2p = \text{Vol}(B) > 2^d i = 2^2 p = 4p$$

We have a point which satisfies $x^2 + y^2 = kp$ for some $k \ge 1$, and also $x^2 + y^2 < 2p$. So $x^2 + y^2 = p$ and we are done.

Euler looked at

$$A^4 + B^4 = C^4 + D^4$$

with $A, B, C, D \in \mathbb{Q}$ This defines a surface in \mathbb{R}^4 . Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

Rational points on surfaces

Euler looked at

$$A^4 + B^4 = C^4 + D^4$$

with $A, B, C, D \in \mathbb{Q}$

This defines a surface in \mathbb{R}^4 .

One parametrisation of solutions[1]:

$$a(s,t) = s^7 + s^5t^2 - 2s^3t^4 + 3s^2t^5 + st^6$$

$$b(s,t) = s^6t - 3s^5t^2 - 2s^4t^3 + s^2t^5 + t^7$$

$$c(s,t) = s^7 + s^5t^2 - 2s^3t^4 - 3s^2t^5 + st^6$$

$$d(s,t) = s^6t + 3s^5t^2 - 2s^4t^3 + s^2t^5 + t^7$$

Euler looked at

$$A^4 + B^4 = C^4 + D^4$$

with $A, B, C, D \in \mathbb{Q}$

This defines a surface in \mathbb{R}^4 .

One parametrisation of solutions[1]:

$$a(s,t) = s^7 + s^5t^2 - 2s^3t^4 + 3s^2t^5 + st^6$$

$$b(s,t) = s^6t - 3s^5t^2 - 2s^4t^3 + s^2t^5 + t^7$$

$$c(s,t) = s^7 + s^5t^2 - 2s^3t^4 - 3s^2t^5 + st^6$$

$$d(s,t) = s^6t + 3s^5t^2 - 2s^4t^3 + s^2t^5 + t^7$$

Unfortunately this does not give us all solutions.

Counting solutions

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on surfaces

Q: How many solutions are there?

Counting solutions

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on surfaces

Q: How many solutions are there?

A: ∞

Can we find a better way of counting them?

We want to estimate the density of our solutions.

We want a different way of describing the size of a rational number.

Take

$$x = \frac{a}{b} \in \mathbb{Q}, \ a, b \in \mathbb{Z}$$
 $\gcd(a, b) = 1$

We want a different way of describing the size of a rational number.

Take

$$x = \frac{a}{b} \in \mathbb{Q}, \ a, b \in \mathbb{Z}$$

$$\gcd(a, b) = 1$$

Then we say height of x is:

$$H(x) := \max\{\log(|a|), \log(|b|)\}$$

We want a different way of describing the size of a rational number.

Take

$$x = \frac{a}{b} \in \mathbb{Q}, \ a, b \in \mathbb{Z}$$

$$\gcd(a, b) = 1$$

Then we say height of x is:

$$H(x) := \max\{\log(|a|), \log(|b|)\}$$

Hurrah! There are only finitely many points with height less than a given value.

Counting points

We now want to count points in a set X with bounded height. We do this in a simple way and define:

$$N(X, B) := \#\{x \in X | H(x) \le B\}$$

We can analyse the growth of this function as B grows.

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry o Numbers

Rational points on surfaces



Numbers

Rational points on surfaces

We now want to count points in a set X with bounded height. We do this in a simple way and define:

$$N(X, B) := \#\{x \in X | H(x) \le B\}$$

We can analyse the growth of this function as B grows. Manin and others have conjectured that the growth of N(X,B) is asymptotically governed by geometric properties of the surface for many problems[2].

Morals

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Geometry of Numbers

Rational points on surfaces

Geometry can help us show solutions exist to Diophantine equations.

Geometric properties govern the density of the solutions to some problems.

Alex J. Best

Introduction

Numbers

Rational points on surfaces

Geometry can help us show solutions exist to Diophantine equations.

Geometric properties govern the density of the solutions to some problems.

And there is a lot more interplay between these two areas.

References

Geometric approaches to solving Diophantine equations

Alex J. Best

Introduction

Numbers

Rational points on surfaces

Hardy, G.H. and Wright, E.M. and Heath-Brown, D.R. and Silverman, J.H., An Introduction to the Theory of Numbers, Oxford University Press, 2008.

Heath-Brown, D.R., Diophantine equations, Algebra, Geometry, Analysis & Logic, Talk at Warwick Mathematics Institute.