

# Part III Algebraic Topology 2014

Jacob Rasmussen  
DPMMS  
University of Cambridge

Alex J. Best  
DPMMS  
University of Cambridge

November 14, 2014

## Abstract

These are lecture notes for the 2014 Part III Algebraic Topology course taught by Dr. Jacob Rasmussen.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Homotopy</b>	<b>2</b>
<b>3</b>	<b>Homology</b>	<b>3</b>
	<b>Notation</b>	<b>4</b>

# 1 Introduction

The recommended books are:

- [Algebraic Topology](#) - Allen Hatcher,
- Homology Theory - James W. Vick,
- Differential Forms in Algebraic Topology - Raoul Bott and Loring W. Tu.

Generated: November 14, 2014, 01:38:39 (Z)

## 2 Homotopy

### 2.1 Homotopies

**Definition 2.1** (Homotopic maps). Maps  $f_0, f_1: X \rightarrow Y$  are said to be **homotopic** if there is a continuous map  $F: X \times I \rightarrow Y$  such that

$$F(x, 0) = f_0(x) \text{ and } F(x, 1) = f_1(x) \quad \forall x \in X.$$

We let  $\text{Map}(X, Y) = \{f: X \rightarrow Y \text{ continuous}\}$ . Then letting  $f_t(x) = F(x, t)$  in the above definition we see that  $f_t$  is a path from  $f_0$  to  $f_1$  in  $\text{Map}(X, Y)$ .

**Example 2.2.** 1.  $X = Y = \mathbf{R}^n$ ,  $f_0(\bar{x}) = \bar{0}$  and  $f_1(\bar{x}) = \bar{x}$  are homotopic via  $f_t(\bar{x}) = t\bar{x}$ .

2.  $S^1 = \{z \in \mathbf{C} : |z| = 1\}$  then

3.  $S^n = \{\bar{x} \in \mathbf{R}^n : |\bar{x}| = 1\}$

**Lemma 2.3.** Homotopy is an equivalence relation on  $\text{Map}(X, Y)$ .

**Lemma 2.4.** If  $f_0 \sim f_1: X \rightarrow Y$  and  $g_0 \sim g_1: Y \rightarrow Z$  then  $g_0 \circ f_0 \sim g_1 \circ f_1$ .

**Corollary 2.5.** For any space  $X$  the set  $[X, \mathbf{R}^n]$  has one element.

*Proof.* Define  $0_X: X \rightarrow \mathbf{R}^n$  by  $0_X(x) = 0 \in \mathbf{R}^n$  for any  $x \in X$ . □

**Definition 2.6** (Contractible space).  $X$  is **contractible** if  $1_X$  is homotopic to a constant map.

**Proposition 2.7.**  $Y$  is contractible  $\iff [X, Y]$  has one element for any space  $X$ .

*Proof.*  $(\Rightarrow)$  as in corollary.  $(\Leftarrow)$   $[X, Y]$  has one element so  $1_Y \sim$  a constant map. □

Given a space  $X$  how can we tell if  $X$  is contractible? If  $X$  is contractible then it must be path connected for one.

*Proof.* Contractible implies that  $[S^0, X]$  has one element and so  $f: S^0 \rightarrow X$  extends to  $D^1$ , and therefore  $X$  is path connected. □

Similarly if  $[S^1, X]$  has more than one element then  $X$  is not contractible.

**Definition 2.8** (Simply connected). We say  $X$  is **simply connected** if  $[S^1, X]$  has only one element.

We say two space  $X$  and  $Y$  are *homotopy equivalent* if there exists  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  such that  $g \circ f \sim 1_X$  and  $f \circ g \sim 1_Y$ .

**Example 2.9.**  $X$  is contractible if and only if  $X \sim \{p\}$ .

*Proof.*  $X$  contractible  $\implies 1_X \sim c$ , a constant map. Choose  $f: X \rightarrow \{p\}$ ,  $f(x) = p$  and  $g: \{p\} \rightarrow X$ ,  $g(p) = c$ . Then  $g \circ f = c \sim 1$  and  $f \circ g = 1_{\{p\}}$ . Converse: exercise.  $\square$

**Exercise 2.10.**

Given  $X$  and  $Y$  how can we determine if  $X \sim Y$ ? How do we determine  $[X, Y]$ ? For example is  $S^n \sim S^m$ .

## 2.2 Homotopy groups

**Definition 2.11** (Map of pairs). A **map of pairs**  $f: (X, A) \rightarrow (Y, B)$  is a map  $f: X \rightarrow Y$  with sets  $A \subset X$  and  $B \subset Y$  such that  $f(A) \subset B$ .

If we have maps of pairs  $f_0, f_1: (X, A) \rightarrow (Y, B)$  then we write  $f_0 \sim f_1$  if there exists  $F: (X \times I, A \times I) \rightarrow (Y, B)$  such that  $F(x, 0) = f_0(x)$  and  $F(x, 1) = f_1(x)$ .

**Definition 2.12** (Homotopy groups). If  $*$   $\in X$  then the  $n$ th homotopy group is

$$\pi_n(X, *) = [(D^n, S^{n-1}) \rightarrow (X, \{*\})].$$

We now note several properties of this definition:

1.  $\pi_0(X, *)$  = set of path components of  $X$ .
2.  $\pi_1(X, *)$  is a group.  $\pi_n(X, *)$  is an abelian group.
3.  $\pi_n$  is a functor

$$\left\{ \begin{array}{c} \text{pointed spaces} \\ \text{pointed maps} \end{array} \right\} \rightarrow \left\{ \begin{array}{c} \text{groups} \\ \text{group homomorphisms} \end{array} \right\}.$$

So given

$$f: (X, p) \rightarrow (Y, q)$$

we get

$$f_*: \pi_n(X, p) \rightarrow \pi_n(Y, q)$$

defined by

$$f_*(\gamma) = f \circ \gamma.$$

$n$	1	2	3	4	5	6	7
$\pi_n(S^2)$	0	$\mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z}/2$	$\mathbf{Z}/2$	$\mathbf{Z}/12$	$\mathbf{Z}/15$

**Example 2.13** (Homotopy groups of  $S^2$ ).

## 3 Homology

Our goal is to construct a functor "from the category of topological spaces and continuous maps to the category of  $\mathbf{Z}$ -modules and  $\mathbf{Z}$ -linear maps. This means to each space  $X$  we associate an abelian group  $H(X) = \bigoplus_{n \geq 0} H_n(X)$ , and to each map  $f: X \rightarrow Y$  a function  $f_*: H_n(X) \rightarrow H_n(Y)$  satisfying  $(1_X)_* = 1_{H_n(X)}$  and  $(f \circ g)_* = f_* \circ g_*$ .

Some properties we would like to have for our construction are:

1. Homotopy invariance, if  $f \sim g: X \rightarrow Y$  then  $f_* = g_*$ .
2. The dimension axiom,  $H_n(X) = 0$  for any  $n > \dim X$ .

### 3.1 Chain complexes

**Definition 3.1** (Chain complex). If  $R$  is a commutative ring then a **chain complex** over  $R$  is a pair  $(C, d)$  satisfying:

1.  $C = \bigoplus_{n \in \mathbf{Z}} C_n$  for  $R$ -modules  $C_n$ .
2.  $d: C \rightarrow C$  where  $d = \bigoplus d_n$  for  $R$ -linear maps  $d_n$ .
3.  $d \circ d = 0$ .

The indexing by  $n$  is called a **grading**. Usually we take  $C_n = 0$  for  $n < 0$ . An element of  $\ker d_n$  is called **closed** or a **cycle**. An element of  $\operatorname{im} d_n$  is called a **boundary**.  $d$  is the **boundary map** or **differential**.

**Definition 3.2** (Homology groups). If  $(C, d)$  is a chain complex, its  $n$ **th homology group** is

$$H_n(C, d) = \ker d_n / \operatorname{im} d_{n+1}.$$

If  $x \in \ker d_n$  we write  $[x]$  for its image in  $H_n(C)$ .

**Example 3.3.** 1.  $C_0 = C_1 = \mathbf{Z}$ ,  $C_i = 0$  otherwise,

$$0 \rightarrow \mathbf{Z} \xrightarrow{\cdot 3} \mathbf{Z} \rightarrow 0.$$

Then  $H_1 = 0$ ,  $H_0 = \mathbf{Z}/3$ .

2.

$\mathbf{Z}$

## Notation

Symbol	Description	Page
--------	-------------	------

2