

MA3H6 Algebraic Topology - Lecture Notes

Based on lectures by Dr. Saul Schleimer

Typeset by Alex J. Best

May 6, 2014

Contents

1	Introduction	1
2	Basics	2
2.1	Topological review	2
3	Simplicial homology	2
3.1	Simplices	2
3.2	Δ -complexes	3
3.3	Abelian groups	4
3.4	Chains	4
3.5	Boundary operators	5
3.6	Chain complexes	6
3.7	Computations	7
4	Singular homology	7
4.1	Reduced Homology	9
4.2	Functoriality	10
4.3	Chain maps	10
4.4	Homotopic spaces	11
4.5	Prisms	12

1 Introduction

These are lecture notes that I typeset for MA3H6 Algebraic Topology in 2014, they are currently full of gaps, mistakes, wrong statements, notation abuse and lots of other badness. However they might be useful to someone, despite the fact they lack very many pictures at present. If you find anything else that can be improved send me an email at a.j.best@warwick.ac.uk, thanks.

2 Basics

2.1 Topological review

Notation.

$\mathbb{R}^n = \{(x_1, \dots, x_n) \mid x_j \in \mathbb{R}\}$ with the product topology of open intervals.

$$\|x\| = \sqrt{\sum x_i^2}.$$

$B^n = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$ the $n - 1$ sphere.

$S^{n-1} = \{x \in \mathbb{R}^n \mid \|x\| = 1\}.$

$\mathbb{R}^0 = \{()\}.$

Exercise.

$$B^n \times B^m \cong B^{n+m}.$$

Exercise.

$$S^n \times S^m \not\cong S^{n+m}.$$

Hint. Find an invariant of topological spaces that distinguishes them.

Invariants Connectedness, Hausdorffness, π_1 , compactness, Euler characteristic. But none of these work.

Quotients We recall that the quotient topology is defined by $a \subseteq X/\sim$ is open iff its preimage under the map $f: X \rightarrow X/\sim$ is open. This topology makes as many of the sets of the quotient as possible open while keeping the quotient map continuous.

There are more ways to produce S^1 , for example

$$S^1 \cong [0, 1]/0 \sim 1$$

when equipped with the quotient topology.

Another way is to consider $\mathbb{R}/\mathbb{Z} = \mathbb{R}/\{x \sim y \iff x - y \in \mathbb{Z}\}$. So there is a map $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ which is the covering map of \mathbb{R}/\mathbb{Z} by its universal cover.

3 Simplicial homology

3.1 Simplices

Definition. We define the n -simplex to be

$$\Delta^n = \left\{ x \in \mathbb{R}^{n+1} \mid x_i \geq 0 \ \forall i, \sum x_i = 1 \right\}.$$

In general if $v_i \in \mathbb{R}^m$ are a collection of $n + 1$ affinely independent points (do not lie in an $n - 1$ dimensional subspace) then we define

$$[v] = [v_0, v_1, \dots, v_n] = \left\{ \sum x_i v_i \mid x_i \in \Delta^n \right\}.$$

If we omit some of the v_i we obtain a facet of $[v]$. If we only omit one of them we get a face. This is denoted by

$$[v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$$

where the v_i is read to be omitted.

The vertices are ordered and if $[v], [w]$ are simplices of the same dimension then there exists a unique affine map extending the ordering of the vertices. The standard map $f: [v] \rightarrow [w]$ sends v_i to w_i and respects barycentric coordinates.

Definition. A facet of Δ is a subsimplex (i.e. pick some x_i and set them to zero).

Definition. A face is a codimension one facet.

Definition. The boundary of Δ^n is denoted by $\partial\Delta^n$ and consists of the union of its faces.

We have that $\mathring{\Delta} = \Delta - \partial\Delta$.

Example.

Exercise. Count the k -dimensional faces of Δ^n .

3.2 Δ -complexes

Definition. Fix X a topological space and a collection of maps

$$\{\sigma_\alpha: \Delta_\alpha \rightarrow X \mid \alpha \in A\}.$$

This is known as a Δ -complex structure on X if:

- (i) (Partition) for all α $\sigma_\alpha|_{\mathring{\Delta}_\alpha}$ is injective and for $x \in X$ there is a *unique* $\alpha \in A$ s.t. $x \in \sigma_\alpha(\mathring{\Delta}_\alpha)$.
- (ii) (Tiling) If $\Delta \subset \Delta_\alpha$ is a face then there is a unique $\beta \in A$ s.t. $\sigma_\alpha|_\Delta = \sigma_\beta \circ f$ where $f: \Delta \rightarrow \Delta_\beta$ is the canonical map.
- (iii) (Topology) $U \subset X$ is open iff $\forall \alpha \sigma_\alpha^{-1}(U) \subset \Delta_\alpha$ is open.

We can state this equivalently as: X must be homeomorphic to the quotient space

$$\bigsqcup_{\alpha \in A} \Delta_\alpha / \text{face gluings}.$$

Example.

Example. $\partial\Delta^n$ gives a Δ -complex structure on S^{n-1} .

Example. If we double Δ^n across $\partial\Delta$ we get a Δ -complex structure on S^n .

Example. Check these are homeomorphic to S^n .

Non Example. Violates tiling on the edge marked $[0, 2]$ and so is not a Δ -complex structure.

Exercise. 1. Find a Δ -complex structure on the space in the non-example above.

2. Show that every graph admits a Δ -complex structure.

Example. Here the indexing set $A = \mathbb{R}$ (very big!).

Definition. A Δ -complex is *finite dimensional* if there exists n s.t. for all α $\dim(\Delta_\alpha) \leq n$.

Definition. A Δ -complex structure is *finite* if $|A| < \infty$ (where as above A is the index set).

Exercise. Show that if X admits a Δ -complex structure then X is Hausdorff.

Exercise. Show that if $\{\sigma_\alpha\}$ is a Δ -complex structure on X and $K \subset X$ is compact then K meets the interiors of only finitely many of the σ_α 's.

Exercise. If X, Y admit Δ -complex structures then so does $X \times Y$.

3.3 Abelian groups

Fix A a set. Define $\mathbb{Z}[A]$ to be the *free abelian group* on A given by

$$\mathbb{Z}[A] = \left\{ \sum_{\alpha \in A} n_\alpha \cdot \alpha \mid n_\alpha \in \mathbb{Z} \text{ and all but finitely many are non-zero} \right\}$$

= all finite \mathbb{Z} -linear sums of elements of A .

Example.

$$\mathbb{Z}[\{\alpha, \beta\}] \cong \mathbb{Z}^2 = \{n\alpha + m\beta \mid m, n \in \mathbb{Z}\}.$$

If A is finite then $\mathbb{Z}[A] \cong \mathbb{Z}^A$. But if $|A| = \infty$ then this is false.

Exercise. \mathbb{Q} is *not* a free abelian group.

3.4 Chains

Suppose $(X, \{\sigma\})$ is a space equipped with a Δ -complex structure.

Definition. We define the set of n -chains to be

$$C_n^\Delta = \mathbb{Z}[\{\sigma_\alpha \mid \dim(\Delta_\alpha) = n\}].$$

Example.

3.5 Boundary operators

Recall $\Delta_v = [v_0, v_1, \dots, v_n]$ is an n -simplex.

The i th face of Δ is $[v_0, v_1, \dots, \hat{v}_i, \dots, v_n]$.

Definition. We define the *boundary operator* as follows. First suppose $\sigma: \Delta \rightarrow X$ is a map.

We then define

$$\partial\sigma = \sum_{i=0}^n (-1)^i \sigma \mid [v_0, \dots, \hat{v}_i, \dots, v_n].$$

Which is an $(n-1)$ -chain.

So we extend linearly to define

$$\partial: C_n^\Delta(X) \rightarrow C_{n-1}^\Delta(X)$$

given by

$$\sum n_\alpha \sigma_\alpha \mapsto \sum n_\alpha \partial\sigma_\alpha.$$

Example.

Lemma.

$$\partial_{n-1} \circ \partial_n = 0.$$

“The extremes of the extremes are empty”.

Proof. It suffices to check this on a basis element

$$\sigma: \Delta^n \rightarrow X$$

so

$$\partial_n \sigma = \sum_{i=0}^n (-1)^i \sigma \mid [v_0, \dots, \hat{v}_i, \dots, v_n]$$

now we apply ∂_{n-1} :

$$\begin{aligned}
\partial_{n-1}\partial_n\sigma &= \partial_{n-1} \left(\sum_{i=0}^n (-1)^i \sigma \mid [v_0, \dots, \hat{v}_i, \dots, v_n] \right) \\
&= \sum_{i=0}^n (-1)^i \partial_{n-1} (\sigma \mid [v_0, \dots, \hat{v}_i, \dots, v_n]) \\
&= \sum_{i=0}^n (-1)^i \sum_{j=0}^{n-1} (-1)^j (\sigma \mid [v_0, \dots, \hat{v}_i, \dots, v_n]) \mid [w_0, \dots, \hat{w}_j, \dots, w_{n-1}] \\
&= \sum_{i=0}^n (-1)^i \left(\sum_{j<i} (-1)^j \sigma \mid [v_0, \dots, \hat{v}_j, \dots, \hat{v}_i, \dots, v_n] \right. \\
&\quad \left. + \sum_{j>i} (-1)^{j+1} \sigma \mid [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_n] \right) \\
&= \sum_{j<i} (-1)^{j+i} \sigma \mid [v_0, \dots, \hat{v}_j, \dots, \hat{v}_i, \dots, v_n] \\
&\quad - \sum_{j>i} (-1)^{j+i} \sigma \mid [v_0, \dots, \hat{v}_i, \dots, \hat{v}_j, \dots, v_n] \\
&= 0
\end{aligned}$$

□

3.6 Chain complexes

Definition. A sequence $\{C_n\}_{n=0}^\infty$ of abelian groups with homomorphisms

$$\partial_n: C_n \rightarrow C_{n-1}$$

such that $\partial^2 = 0$ is called a *chain complex*.

By convention we take C_{-1} to be 0.

Example.

$$0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow 0.$$

Given two chain complexes we can form the direct sum by taking the direct sum of each of the groups and letting the operators act elementwise.

Terminology If $c \in C_n$ we call c an *n-chain*.

If $z \in Z_n = \ker(\partial_n)$ we call z an *n-cycle*.

If $b \in B_n = \text{im}(\partial_{n-1})$ we call b an *n-boundary*.

If $h \in Z_n/B_n = H_n$ we call h a *homology class*.

Since $\partial^2 = 0$ we deduce that $B_n \leq Z_n$ and $H_n = Z_n/B_n$ makes sense.

Example. For

$$0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow 0$$

we have $H_1 = 0$, $H_0 = \mathbb{Z}/2\mathbb{Z}$ and $H_k = 0$ for all $k \geq 1$.

Definition. If (X, σ) is a Δ -complex then set $C_n^\Delta(X) = \mathbb{Z}[\{\sigma_\alpha \mid \dim(\Delta_\alpha) = n\}]$ and $\partial_n: C_n^\Delta(X) \rightarrow C_{n-1}^\Delta(X)$ is the boundary operator.

Then $H_n^\Delta(X)$ are called the *simplicial homology groups* of X .

Theorem. This is independent of the choice of Δ -complex structure on X .

3.7 Computations

1. $X = \{\text{pt}\}$. $C_0^\Delta(X) \cong \mathbb{Z}$ and all others are 0, so we have the chain complex:

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \rightarrow 0.$$

So $H_0^\Delta(\text{pt}) \cong \mathbb{Z}$ and $H_k^\Delta(\text{pt}) \cong 0$ if $k \geq 1$.

2. $X = S^1$. $C_0^\Delta(X) \cong \mathbb{Z}$, $C_1^\Delta(X) \cong \mathbb{Z}$ and all others are 0, so we have the chain complex:

$$\cdots \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{\partial} \mathbb{Z} \rightarrow 0.$$

We see that $\partial e = \sum_{i=0}^1 (-1)^i e|[v_0, \dots, \hat{v}_i, \dots, v_1] = e|[v_1] - e|[v_0] = v - v = 0$. So

$$H_k^\Delta(S^1) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0 \text{ or } 1, \\ 0 & \text{otherwise.} \end{cases}$$

Exercise. Compute $H_\bullet^\Delta(S^1)$ for the Δ -complex structure on S^1 with k vertices and k edges.

Exercise. Compute $H_\bullet^\Delta(X)$ for the $X = B^2$, S^1 and K^2 (the Klein bottle).

Exercise. Using the fact that Δ^n is a Δ -complex structure on B^n compute $H_\bullet^\Delta(B^n)$.

In general you'll want to make use of the Smith normal form.

4 Singular homology

Definition. A *singular n -simplex* in X is a map $\sigma: \Delta^n \rightarrow X$.

Definition.

$$C_n^{\text{sing}}(X) = \mathbb{Z}[\{\sigma: \Delta^n \rightarrow X\}].$$

We call $c \in C_n^{\text{sing}}(X)$ a singular n -chain.

Definition. We define $\partial: C_n^{\text{sing}}(X) \rightarrow C_{n-1}^{\text{sing}}(X)$ exactly as before by

$$\partial\sigma = \sum_{i=0}^n (-1)^i \sigma|[v_0, \dots, \hat{v}_i, \dots, v_n].$$

And again we define $Z_n^{\text{sing}}(X)$ (resp. $B_n^{\text{sing}}(X)$) exactly as above and call it the group of singular n -cycles (resp. n -boundaries).

Definition. Now $H_n^{\text{sing}}(X) = Z_n^{\text{sing}}(X)/B_n^{\text{sing}}(X)$ is the n -th *singular homology group*.

Remark. We have that $\partial_{n-1} \circ \partial_n = 0$ exactly as before.

Example. Suppose X is a single point, then there is a unique Δ -complex structure on X . We say $\sigma^0: \mathbb{Z} \rightarrow X$ is the “constant map”. So $0 \rightarrow \mathbb{Z} \rightarrow 0$ is the chain complex $C_{\bullet}^{\Delta}(X)$. So

$$H_n^{\Delta} = \begin{cases} \mathbb{Z} & n = 0, \\ 0 & n \geq 1. \end{cases}$$

Suppose X is as above again, then we can compute

$$H_n^{\text{sing}} = \begin{cases} \mathbb{Z} & n = 0, \\ 0 & n \geq 1. \end{cases}$$

This is as in dimension n there is only the constant map

$$\sigma^n: \Delta^n \rightarrow X$$

so $C_n^{\text{sing}}(X) \cong \mathbb{Z}$ and we also have that

$$\begin{aligned} \partial\sigma^n &= \sum_{i=0}^n (-1)^i \sigma|[v_0, \dots, \hat{v}_i, \dots, v_n] = \sum_{i=0}^n (-1)^i \sigma^{n-1} \\ &= \left(\sum_{i=0}^n (-1)^i \right) \sigma^{n-1} = \begin{cases} 0 & n \text{ odd}, \\ \sigma^{n-1} & n \text{ even}, \end{cases} \end{aligned}$$

except if $n = 0$. So C_n is

$$\dots \rightarrow \mathbb{Z} \xrightarrow{\times 0} \mathbb{Z} \xrightarrow{\times 1} \mathbb{Z} \xrightarrow{\times 0} \mathbb{Z} \xrightarrow{\times 1} \mathbb{Z} \xrightarrow{\times 0} \mathbb{Z} \rightarrow 0$$

and so the singular homology groups are as claimed.

Challenge Compute $H_n^{\text{sing}}(S^1)$ from the definitions.

Theorem. If X admits a Δ -complex structure then

$$H_{\bullet}^{\Delta}(X) \cong H_{\bullet}^{\text{sing}}(X).$$

The left hand side is generally easier to compute, but the right can be easier to prove theorems with.

Proposition. If $X = \sqcup X_{\alpha}$ where all X_{α} are path connected spaces then

$$H_n^{\text{sing}}(X) = \bigoplus_{\alpha} H_n^{\text{sing}}(X_{\alpha}).$$

Proof.

$$C_n^{\text{sing}}(X) = \bigoplus_{\alpha} C_n^{\text{sing}}(X_{\alpha}).$$

and ∂ respects this “splitting”. □

Proposition. If $X \neq \emptyset$ and X is path connected then $H_0^{\text{sing}}(X) \cong \mathbb{Z}$.

Proof. Define $\epsilon: C_0(X) \rightarrow \mathbb{Z}$ by $\sum n_{\alpha}v_{\alpha} \mapsto \sum n_{\alpha}$, the augmentation map, then ϵ is surjective. We claim that $\ker(\epsilon) = \text{im}(\partial_1)$. Given any $\tau: \Delta^1 \rightarrow X$ that goes from v to w we have that $\partial\tau = w - v$ so $\epsilon(\partial\tau) = 1 - 1 = 0$ and the image is contained in the kernel. Now fix $\sum n_{\alpha}v_{\alpha}$ s.t. $\epsilon(\sum n_{\alpha}v_{\alpha}) = 0$. Also fix some $u \in X$ and for all α pick $\tau_{\alpha}: \Delta^1 \rightarrow X$ a path from u to v_{α} . Consider $\sum n_{\alpha}\tau_{\alpha} \in C_1(X)$

$$\begin{aligned} \partial(\sum n_{\alpha}\tau_{\alpha}) &= \sum \partial(n_{\alpha}\tau_{\alpha}) \\ &= \sum n_{\alpha}\partial\tau_{\alpha} \\ &= \sum n_{\alpha}(v_{\alpha} - u) \\ &= \sum n_{\alpha}v_{\alpha} - \sum n_{\alpha}u \\ &= \sum n_{\alpha}v_{\alpha} - u \sum n_{\alpha} \\ &= \sum n_{\alpha}v_{\alpha} - u \cdot 0 \in \text{im} \end{aligned}$$

Hence $\text{im} = \ker$ as claimed.

And so $H_0 = \ker(\partial_0)/\text{im}(\partial_1) = \ker(\partial_0)/\ker(\epsilon) = C_0(X)/\ker(\epsilon) \cong \mathbb{Z}$. □

4.1 Reduced Homology

Definition. If X has k path components, then $H_0(X) \cong \mathbb{Z}^k$ so we define the *augmented chain complex*

$$\cdots \rightarrow C_2(X) \xrightarrow{\partial_2} C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0,$$

where ϵ is the augmentation map from above. Define the *reduced homology groups* $\tilde{H}_\bullet(X)$ to be the homology groups of this chain complex. So $\tilde{H}_n(X) = H_n(X)$ is $n > 0$ and $\tilde{H}_0(X) = \ker(\epsilon)/\text{im}(\partial_1)$. Hence if X has k path components

$$\tilde{H}_0(X) \cong \mathbb{Z}^{k-1}.$$

Recall that $H_\bullet(X \sqcup Y) = H_\bullet(X) \oplus H_\bullet(Y)$, the reduced homology groups behave nicely with respect to many operations such as 1-point unions. In a 1-point union $X \vee Y = X \sqcup Y / x \sim y$ for some designated point $x \in X$ and $y \in Y$. So $\tilde{H}_\bullet(X \vee Y) = \tilde{H}_\bullet(X) \oplus \tilde{H}_\bullet(Y)$.

4.2 Functoriality

Definition. Suppose $f: X \rightarrow Y$ is a (continuous) map. Let $f_n: C_n(X) \rightarrow C_n(Y)$ by $\sigma \mapsto f \circ \sigma$. The function $f \circ \sigma$ is again a map from Δ^n to Y and so still lies in $C_n(Y)$.

The key property of this definition is that $\partial_n \circ f_n = f_{n-1} \circ \partial_n$. This is saying that the square

$$\begin{array}{ccc} C_n(X) & \xrightarrow{\partial_n} & C_{n-1}(X) \\ \downarrow f_n & & \downarrow f_{n-1} \\ C_n(Y) & \xrightarrow{\partial_n} & C_{n-1}(Y) \end{array}$$

commutes. We denote the family of these maps f_n as $f_\#$,

Exercise. If $X \xrightarrow{f} Y \xrightarrow{g} Z$ then $(g \circ f)_\# = g_\# \circ f_\#$.

4.3 Chain maps

Definition. If C_\bullet, D_\bullet are chain complexes we say that a family of homomorphisms $f_\#: C_\bullet \rightarrow D_\bullet$ is a *chain map* if

$$f_\# \circ \partial = \partial \circ f_\#$$

Example. If $f: X \rightarrow Y$ is continuous then $f_\#$ is a chain map from $C_\bullet(X) \rightarrow C_\bullet(Y)$.

Example. Suppose (X, σ) is a Δ -complex then

$$i: C_n^\Delta(X) \rightarrow C_n^{\text{sing}}(X)$$

is also a chain map.

Proposition. If $f_\#: C_\bullet \rightarrow D_\bullet$ is a chain map then $f_\#$ induces a homomorphism

$$f_*: H_\bullet(C) \rightarrow H_\bullet(D)$$

given by

$$f_*([z]) = [f_\#(z)].$$

Proof. Check that $f_{\#}(Z_n^C) \leq Z_n^D$ (exercise) and that $f_{\#}(B_n^C) \leq B_n^D$. So $f_{\#}(b) = f_{\#}(\partial c) = \partial f_{\#}(c)$. \square

Remark. If $f: X \rightarrow Y$ is a homomorphism then there exists a continuous inverse $g: Y \rightarrow X$ such that

$$f_*: H_{\bullet}(X) \rightarrow H_{\bullet}(Y)$$

is inverse to

$$g_*: H_{\bullet}(Y) \rightarrow H_{\bullet}(X).$$

4.4 Homotopic spaces

Definition. We say two maps f and g from $X \rightarrow Y$ are *homotopic* if there is a map $F: X \times [0, 1] \rightarrow Y$ such that $f(x) = F(x, 0)$ and $g(x) = F(x, 1)$. We write $f \sim g$.

We then say two spaces X and Y are *homotopy equivalent* if there exists maps $f: X \rightarrow Y$ and $g: Y \rightarrow X$ such that

$$(g \circ f) \sim \text{Id}_X \text{ and } (f \circ g) \sim \text{Id}_Y.$$

Example.

$$S^n \sim \mathbb{R}^{n+1} \setminus \{0\}$$

via (for $n = 1$)

$$\begin{aligned} i: S^1 &\rightarrow \mathbb{R}^2 \setminus \{0\} \\ x &\mapsto x \end{aligned}$$

and

$$\begin{aligned} r: \mathbb{R}^2 \setminus \{0\} &\rightarrow S^1 \\ x &\mapsto \frac{x}{\|x\|}. \end{aligned}$$

We also have

$$S^n \sim B^{n+1} \setminus \{0\}$$

Theorem. If X is homotopy equivalent to Y via f and g then

$$f_* = g_*^{-1}: H_{\bullet}(X) \rightarrow H_{\bullet}(Y).$$

Corollary. If X is homotopy equivalent to Y via f then

$$f_*: H_{\bullet}(X) \rightarrow H_{\bullet}(Y)$$

is an isomorphism.

Proof.

$$(\text{Id}_X)_* = \text{Id}_{H_{\bullet}}.$$

\square

Definition. Suppose $f_{\#}, g_{\#}: C_{\bullet} \rightarrow D_{\bullet}$ are chain maps. A sequence of homomorphisms $P_n: C_n \rightarrow D_{n+1}$ is called a *chain homotopy* if

$$\partial_{n-1}P_n + P_{n-1}\partial_n = g_{\#} - f_{\#}$$

if there is a chain homotopy between two chain maps $f_{\#}, g_{\#}$ we write $f_{\#} \sim g_{\#}$.

$$\begin{array}{ccccc} C_{n+1} & \xrightarrow{\partial} & C_n & \xrightarrow{\partial} & C_{n-1} \\ \left(\begin{smallmatrix} f \\ \downarrow \end{smallmatrix}\right)_g & & \left(\begin{smallmatrix} f \\ \downarrow \end{smallmatrix}\right)_g & & \left(\begin{smallmatrix} f \\ \downarrow \end{smallmatrix}\right)_g \\ D_{n+1} & \xrightarrow{\partial} & D_n & \xrightarrow{\partial} & D_{n-1} \end{array}$$

Proposition. If $f_{\#} \sim g_{\#}: C_{\bullet} \rightarrow D_{\bullet}$ then

$$f_* = g_*: H_{\bullet}(C) \rightarrow H_{\bullet}(D).$$

Proof. Pick any $h \in H_{\bullet}(C)$, we want to show $(g_* - f_*)(h) = 0$. Choose some $x \in Z_n(C)$ such that $h = [z]$ and compute

$$\begin{aligned} (g_* - f_*)(h) &= (g_* - f_*)([z]) \\ &= [(g_{\#} - f_{\#})(z)] \\ &= [(P\partial + \partial P)(z)] \\ &= [P\partial z + \partial Pz] \\ &= [P0 + \partial Pz] \\ &= [\partial(Pz)] \\ &= 0 \text{ (as } B_n = 0 \text{ in homology)}. \end{aligned}$$

□

4.5 Prisms

Definition. A *prism* is a copy of $\Delta^n \times I$.

We can subdivide $\Delta \times I$ into $n + 1$ dimensional simplices of the form

$$[v_0, v_1, \dots, v_i, w_{i+1}, \dots, w_n],$$

where v are the vertices of the simplex at one end of the interval and w are the vertices at the other.

If we have $F: X \times I \rightarrow Y$ and $\sigma: \Delta^n \rightarrow X$ we let

$$F\sigma = F \circ (\sigma \times \text{Id}_I): \Delta^n \times I \rightarrow Y.$$