

Part III Algebraic Topology 2014

Part III Algebraic Topology 2014

Contents

1	Homology	1
1.1	Introduction	1
1.2	Homotopy	1
1.3	Homology	3
1.4	Homology of a pair	5
1.5	Subdivision and Excision	9
1.6	Degree and Orientations	10
1.7	Cell Complexes	10
2	Cohomology and Products	11
2.1	Homology with Coefficients and Cohomology	11
2.2	Notation	13

Chapter 1

Homology

1.1 Introduction

These are lecture notes for the 2014 Part III Algebraic Topology course taught by Dr. Jacob Rasmussen.

The recommended books are:

- [Algebraic Topology](#) - Allen Hatcher,
- Homology Theory - James W. Vick,
- Differential Forms in Algebraic Topology - Raoul Bott and Loring W. Tu.

Generated: November 17, 2014, 21:56:42 (Z)

1.2 Homotopy

1.2.1 Homotopies

Definition 1.2.1 (Homotopic maps). Maps $f_0, f_1: X \rightarrow Y$ are said to be **homotopic** if there is a continuous map $F: X \times I \rightarrow Y$ such that

$$F(x, 0) = f_0(x) \text{ and } F(x, 1) = f_1(x) \quad \forall x \in X.$$

We let $\text{Map}(X, Y) = \{f: X \rightarrow Y \text{ continuous}\}$. Then letting $f_t(x) = F(x, t)$ in the above definition we see that f_t is a path from f_0 to f_1 in $\text{Map}(X, Y)$.

Example 1.2.2. 1. $X = Y = \mathbf{R}^n$, $f_0(\bar{x}) = \bar{0}$ and $f_1(\bar{x}) = \bar{x}$ are homotopic via $f_t(\bar{x}) = t\bar{x}$.

2. $S^1 = \{z \in \mathbf{C} : |z| = 1\}$ then

3. $S^n = \{\bar{x} \in \mathbf{R}^n : |\bar{x}| = 1\}$

Lemma 1.2.3. *Homotopy is an equivalence relation on $\text{Map}(X, Y)$.*

Lemma 1.2.4. *If $f_0 \sim f_1: X \rightarrow Y$ and $g_0 \sim g_1: Y \rightarrow Z$ then $g_0 \circ f_0 \sim g_1 \circ f_1$.*

Corollary 1.2.5. *For any space X the set $[X, \mathbf{R}^n]$ has one element.*

Proof. Define $0_X: X \rightarrow \mathbf{R}^n$ by $0_X(x) = 0 \in \mathbf{R}^n$ for any $x \in X$. □

Definition 1.2.6 (Contractible space). X is **contractible** if 1_X is homotopic to a constant map.

Proposition 1.2.7. Y is contractible $\iff [X, Y]$ has one element for any space X .

Proof. (\implies) as in corollary. (\impliedby) $[X, Y]$ has one element so $1_Y \sim$ a constant map. \square

Given a space X how can we tell if X is contractible? If X is contractible then it must be path connected for one.

Proof. Contractible implies that $[S^0, X]$ has one element and so $f: S^0 \rightarrow X$ extends to D^1 , and therefore X is path connected. \square

Similarly if $[S^1, X]$ has more than one element then X is not contractible.

Definition 1.2.8 (Simply connected). We say X is **simply connected** if $[S^1, X]$ has only one element.

We say two space X and Y are *homotopy equivalent* if there exists $f: X \rightarrow Y$ and $g: Y \rightarrow X$ such that $g \circ f \sim 1_X$ and $f \circ g \sim 1_Y$.

Example 1.2.9. X is contractible if and only if $X \sim \{p\}$.

Proof. X contractible $\implies 1_X \sim c$, a constant map. Choose $f: X \rightarrow \{p\}$, $f(x) = p$ and $g: \{p\} \rightarrow X$, $g(p) = c$. Then $g \circ f = c \sim 1$ and $f \circ g = 1_{\{p\}}$. Converse: exercise. \square

Exercise 1.2.10.

Given X and Y how can we determine if $X \sim Y$? How do we determine $[X, Y]$? For example is $S^n \sim S^m$.

1.2.2 Homotopy groups

Definition 1.2.11 (Map of pairs). A **map of pairs** $f: (X, A) \rightarrow (Y, B)$ is a map $f: X \rightarrow Y$ with sets $A \subset X$ and $B \subset Y$ such that $f(A) \subset B$.

If we have maps of pairs $f_0, f_1: (X, A) \rightarrow (Y, B)$ then we write $f_0 \sim f_1$ if there exists $F: (X \times I, A \times I) \rightarrow (Y, B)$ such that $F(x, 0) = f_0(x)$ and $F(x, 1) = f_1(x)$.

Definition 1.2.12 (Homotopy groups). If $*$ $\in X$ then the n th **homotopy group** is

$$\pi_n(X, *) = [(D^n, S^{n-1}) \rightarrow (X, \{*\})].$$

We now note several properties of this definition:

1. $\pi_0(X, *)$ = set of path components of X .
2. $\pi_1(X, *)$ is a group. $\pi_n(X, *)$ is an abelian group.
3. π_n is a functor

$$\left\{ \begin{array}{c} \text{pointed spaces} \\ \text{pointed maps} \end{array} \right\} \rightarrow \left\{ \begin{array}{c} \text{groups} \\ \text{group homomorphisms} \end{array} \right\}.$$

So given

$$f: (X, p) \rightarrow (Y, q)$$

we get

$$f_*: \pi_n(X, p) \rightarrow \pi_n(Y, q)$$

defined by

$$f_*(\gamma) = f \circ \gamma.$$

n	1	2	3	4	5	6	7
$\pi_n(S^2)$	0	\mathbf{Z}	\mathbf{Z}	$\mathbf{Z}/2$	$\mathbf{Z}/2$	$\mathbf{Z}/12$	$\mathbf{Z}/15$

Example 1.2.13 (Homotopy groups of S^2).

1.3 Homology

Our goal is to construct a functor H_* from the category of topological spaces and continuous maps to the category of \mathbf{Z} -modules and \mathbf{Z} -linear maps. This means to each space X we associate an abelian group $H_*(X) = \bigoplus_{n \geq 0} H_n(X)$, and to each map $f: X \rightarrow Y$ a function $f_*: H_n(X) \rightarrow H_n(Y)$ satisfying $(1_X)_* = 1_{H_n(X)}$ and $(f \circ g)_* = f_* \circ g_*$.

Some properties we would like to have for our construction are:

1. Homotopy invariance, if $f \sim g: X \rightarrow Y$ then $f_* = g_*$.
2. The dimension axiom, $H_n(X) = 0$ for any $n > \dim X$.

1.3.1 Chain complexes

Definition 1.3.1 (Chain complex). If R is a commutative ring then a **chain complex** over R is a pair (C, d) satisfying:

1. $C = \bigoplus_{n \in \mathbf{Z}} C_n$ for R -modules C_n .
2. $d: C \rightarrow C$ where $d = \bigoplus d_n$ for R -linear maps d_n .
3. $d \circ d = 0$.

The indexing by n is called a **grading**. Usually we take $C_n = 0$ for $n < 0$. An element of $\ker d_n$ is called **closed** or a **cycle**. An element of $\operatorname{im} d_n$ is called a **boundary**. d is the **boundary map** or **differential**.

Definition 1.3.2 (Homology groups). If (C, d) is a chain complex, its n **th homology group** is

$$H_n(C, d) = \ker d_n / \operatorname{im} d_{n+1}.$$

If $x \in \ker d_n$ we write $[x]$ for its image in $H_n(C)$.

Example 1.3.3. 1. $C_0 = C_1 = \mathbf{Z}$, $C_i = 0$ otherwise,

$$0 \rightarrow \mathbf{Z} \xrightarrow{-3} \mathbf{Z} \rightarrow 0.$$

Then $H_1 = 0$, $H_0 = \mathbf{Z}/3$.

2.

$$\mathbf{Z} = \langle e \rangle \rightarrow \mathbf{Z}^2 = \langle f_1, f_2 \rangle \rightarrow \mathbf{Z} = \langle g \rangle \rightarrow 0$$

with $d(e) = f_1 - f_2$, $d(f_1) = d(f_2) = g$, then $H_*(C) = 0$ (exercise).

1.3.2 The chain complex of a simplex

Definition 1.3.4 (n -simplex). The n -dimensional simplex Δ^n is

$$\Delta^n = \left\{ (x_0, \dots, x_n) \in \mathbf{R}^n : \sum_i x_i = 1, x_i \geq 0 \forall i \right\}.$$

Δ^n has **vertices** v_0, \dots, v_n which are the intersections with the coordinate axes. The k -dimensional **faces** are in bijection with the $k+1$ element subsets of $\{0, \dots, n\}$.

Definition 1.3.5 (Simplicial chain complex). $S_*(\Delta^n)$ is the chain complex with $S_k(\Delta^n)$ the free \mathbf{Z} -module generated by the k -dimensional faces of Δ^n . So

$$S_k(\Delta^n) = \langle e_I : I = \{i_0, \dots, i_k : 0 \leq i_0 \leq \dots \leq i_k \leq n\} \rangle.$$

To define d it suffices to define $d(e_I)$, we let

$$d(e_I) = \sum_{j=0}^k (-1)^j e_{i_0, \dots, i_{j-1}, i_{j+1}, \dots, i_k} \in S_{k-1}(\Delta^n).$$

Example 1.3.6. 1. For Δ^1 we have $d(e_{0,1}) = e_1 - e_0$.

2. For Δ^2 we have $d(e_{0,1,2}) = e_{12} - e_{02} + e_{01}$ and so $d^2(e_I) = (e_2 - e_1) - (e_2 - e_0) + (e_1 - e_0) = 0$.

Lemma 1.3.7. $d^2 = 0$

Proof. It suffices to show $d^2(e_I) = 0$ for all I .

$$\begin{aligned} d^2(e_I) &= d \left(\sum_{j=0}^k (-1)^j e_{i_0, \dots, \hat{i}_j, \dots, i_k} \right) \\ &= \sum_{j=0}^k (-1)^j d(e_{i_0, \dots, \hat{i}_j, \dots, i_k}) \\ &= \sum_{j=0}^k (-1)^j \left(\sum_{l < j} (-1)^l e_{i_0, \dots, \hat{i}_l, \dots, \hat{i}_j, \dots, i_k} + \sum_{l > j} (-1)^{l-1} e_{i_0, \dots, \hat{i}_j, \dots, \hat{i}_l, \dots, i_k} \right) \\ &= \sum_{j=0}^k \left(\sum_{l < j} (-1)^{l+j} e_{I - i_l - i_j} - \sum_{l > j} (-1)^{l+j} e_{I - i_j - i_l} \right) = 0. \end{aligned}$$

□

Example 1.3.8 (Computing $H_*(S_*(\Delta^2))$).

$$0 \rightarrow \mathbf{Z} = \langle e_{012} \rangle \rightarrow \mathbf{Z}^3 = \langle e_{01}, e_{02}, e_{12} \rangle \rightarrow \mathbf{Z}^3 = \langle e_0, e_1, e_2 \rangle \rightarrow 0$$

with $d(e_{012}) = e_{12} - e_{02} + e_{01}$. So $\ker d_2 = 0 \implies H_2 = 0$.

$$d(ae_{01} + be_{02} + ce_{12}) = a(e_1 - e_0) + b(e_2 - e_1) + c(e_2 - e_0)$$

so $ae_{01} + be_{02} + ce_{12} \in \ker d_1 \iff -a - b = 0, a - c = 0, c + b = 0 \iff a = -b = c$ hence $\ker d_1 = \langle e_{01} - e_{02} + e_{12} \rangle = \text{im } d_2$ and so $H_1 = 0$. $\ker(d_0) = 0$, $\text{im } d_1 = \langle e_1 - e_0, e_2 - e_1, e_2 - e_0 \rangle$ so $H_0 = \mathbf{Z} = \langle [e_0] \rangle$.

Exercise 1.3.9. Show that $H_*(S_*(\Delta^n)) = 0$ if $k \neq 0$ and \mathbf{Z} if $k = 0$.

1.3.3 The singular chain complex

Definition 1.3.10 (Singular chain complex). If X is a space, the **singular chain complex** of X , $C_*(X)$ is defined by

$$C_n(x) = \langle e_\sigma : \sigma : \Delta^n \rightarrow X \text{ any continuous map} \rangle.$$

Where

$$d(e_\sigma) = \sum_{j=0}^n (-1)^j e_{\sigma \circ F_j} \in C_{n-1}(X)$$

where $F_j : \Delta^{n-1} \rightarrow \Delta^n$ is given by $F_j(x_0, \dots, x_{n-1}) = (x_0, \dots, 0, \dots, x_{n-1})$ with the 0 in the j th place.

1.3.3.1 Homotopy invariance

Definition 1.3.11 (Chain homotopic maps). Suppose $\phi, \psi : C_* \rightarrow C'_*$ are chain maps, we say that ϕ are **chain homotopic** if there exists an R -linear map $h : C_* \rightarrow C'_{*+1}$ such that $d' \circ h + h \circ d = \phi - \psi$. We denote this relation by $\phi \sim \psi$.

Lemma 1.3.12. *If $\phi \sim \psi$ then $\phi_* = \psi_*$.*

Proof.

$$\begin{aligned}\phi_*([x]) - \psi_*([x]) &= [\phi(x) - \psi(x)] \\ &= [d'hx + hdx] = [d'hx] = 0 \in H_*(C').\end{aligned}$$

□

Theorem 1.3.13. *Suppose $f \sim g: X \rightarrow Y$ via H then $f_{\#} \sim g_{\#} \implies f_* = g_*$.*

Proof.

□

Corollary 1.3.14. *If $X \sim Y$ then $H_*(X) \cong H_*(Y)$.*

Corollary 1.3.15. *If X is contractible then $H_*(X) \cong H_*(\{p\}) \cong \mathbf{Z}$ if $*$ = 0, 0 otherwise.*

1.4 Homology of a pair

1.4.1 Exact sequences

Definition 1.4.1 (Exact sequence). A sequence

$$\cdots \rightarrow A_{n+1} \rightarrow A_n \rightarrow A_{n-1} \rightarrow \cdots$$

of R -modules and R -linear maps is **exact at** A_n if $\ker f_n = \operatorname{im} f_{n+1}$.

A sequence is **exact** if it is exact at all A_n , then (A_*, f) is known as a **acyclic** chain complex (the homology is zero).

Example 1.4.2. 1. $0 \rightarrow A \xrightarrow{f} B$ is exact if and only if f is surjective.

2. $B \xrightarrow{g} C \rightarrow 0$ is exact if and only if g is injective.

3. $0 \rightarrow A \xrightarrow{f} A' \rightarrow 0$ is exact if and only if f is an isomorphism.

4. $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is exact if and only if $A \subset B$ and $C \cong B/A$.

Definition 1.4.3 (Short exact sequence). A sequence

$$0 \rightarrow A_* \xrightarrow{i} B_* \xrightarrow{\pi} C_* \rightarrow 0$$

is a **short exact sequence** of chain complexes if

1. A_*, B_*, C_* are chain complexes.

2. i, π are chain maps.

3.

$$0 \rightarrow A_n \xrightarrow{i} B_n \xrightarrow{\pi} C_n \rightarrow 0$$

is exact for all n .

Lemma 1.4.4 (Snake lemma). *If*

$$0 \rightarrow A_* \xrightarrow{i} B_* \xrightarrow{\pi} C_* \rightarrow 0$$

*is a short exact sequence of chain complexes then there is an associated **long exact sequence** of homology groups*

1.4.2 Homology of a pair

Definition 1.4.9 (Homology of a pair). $H_*(X, A) = H_*(C_*(X, A))$ is the **homology of the pair** (X, A) .

From this we obtain:

Definition 1.4.10 (Long exact sequence of a pair). The **long exact sequence of the pair** (X, A) is

$$\cdots \rightarrow H_n(A) \rightarrow H_n(X) \rightarrow H_n(X, A) \xrightarrow{\partial} H_{n-1}(A) \rightarrow \cdots$$

If $f: (X, A) \rightarrow (Y, B)$ is a map of pairs then we get an **induced map** $f_\#: C_*(X) \rightarrow C_*(Y)$ defined by

$$(\sigma: \Delta^n \rightarrow A) \mapsto f \circ \sigma.$$

Observe that $f_\#(C_*(A)) \subset C_*(B)$ and so $f_\#$ descends to a map

$$f_\#: C_*(X)/C_*(A) \rightarrow C_*(Y)/C_*(B)$$

or equivalently $f_\#: C_*(X, A) \rightarrow C_*(Y, B)$ this then induces $f_*: H_*(X, A) \rightarrow H_*(Y, B)$.

Proposition 1.4.11 (Homotopy invariance). *If $f, g: (X, A) \rightarrow (Y, B)$ are homotopic as maps of pairs then $f_* = g_*: H_*(X, A) \rightarrow H_*(Y, B)$.*

Proof. Let $H: (X \times I, A \times I) \rightarrow (Y, B)$ be the homotopy, H induces a chain homotopy $h: C_*(X) \rightarrow C_{*+1}(Y)$ where $dh + hd = f_\# - g_\#$. $H(A \times I) \subset B$ so $h(C_*(A)) \subset C_{*+1}(B)$ this implies that h descends to a map

$$h: C_*(X)/C_*(A) \rightarrow C_{*+1}(Y)/C_{*+1}(B)$$

with $hd + dh = f_\# - g_\#$ as any relation satisfied will remain satisfied in the quotient. So we have $h: C_*(X, A) \rightarrow C_{*+1}(Y, A)$ and hence $f_\#, g_\#: C_*(X, A) \rightarrow C_*(Y, B)$ are chain homotopic. \square

1.4.2.1 Visualising relative homology classes

If W^{n+1} is a connected oriented compact manifold then we'll show that $H_{n+1}(W, \partial W) \cong \mathbf{Z} = \langle [W, \partial] \rangle$ (where the ∂ notation means relative to boundary). So given $f: (W, \partial) \rightarrow (X, A)$ we get $f_*([W, \partial W]) \in H_{n+1}(X, A)$.

Example 1.4.12. Let $X = \mathbf{R}^3$ and $A = S^1$ then $W = D^2$ defines a class in $H_2(\mathbf{R}^3, S^1)$ (the boundary of W lies inside of A).

1.4.3 Good pairs

Definition 1.4.13 (Good pair). (X, A) is a **good pair** if

1. $\exists U \subset X$ open and $A \subset U$
2. $\exists \pi: U \rightarrow A$ with $\pi|_A = \text{id}_A$.
3. $\pi \sim 1_{(U, A)}$ as maps of pairs from (U, A) to itself.
(i.e. A is a deformation retract of U).

Example 1.4.14 (Good pairs). 1. (smooth manifold, closed submanifold).

2. (simplicial complex, subcomplex).

Example 1.4.15 (Not good pairs). 1. (\mathbf{R}, \mathbf{Q}) .

2. Letting $H \subset \mathbf{R}^2$ be the Hawaiian earring then (\mathbf{R}^2, H) is not a good pair.

Theorem 1.4.16. *Take $A \subset X$ and let π be the natural map $\pi: (X, A) \rightarrow (X/A, A/A) \cong (X/A, *)$. Then if (X, A) is a good pair the induced map $\pi_*: H_*(X, A) \rightarrow H_*(X/A, *)$ is an isomorphism.*

Proof. Postponed. □

Exercise 1.4.17. The composite map ϕ in

$$\tilde{H}_*(X) \rightarrow H_*(X) \rightarrow H_*(X, *)$$

is an isomorphism. i.e.

$$H_*(X, *) = \begin{cases} H_*(X), & * > 0, \\ H_0(X)/\mathbf{Z}, & * \leq 0. \end{cases}$$

Proposition 1.4.18.

$$\tilde{H}_*(S^n) = \begin{cases} \mathbf{Z}, & * = n, \\ 0, & * \neq n. \end{cases}$$

Proof. We proceed by induction on n . For $n = 0$ we have $S^0 = \{p_+, p_-\}$ implying

$$H_*(S^0) = H_*(p_+) \oplus H_*(p_-) = \begin{cases} \mathbf{Z}^2, & * = 0, \\ 0, & * \neq 0, \end{cases}$$

giving

$$\tilde{H}_*(S^0) = \begin{cases} \mathbf{Z}, & * = 0, \\ 0, & * \neq 0. \end{cases}$$

Now consider the long exact sequence of the pair (D^n, S^{n-1})

$$H_*(D^n) \rightarrow H_*(D^n, S^{n-1}) \rightarrow H_{*-1}(S^{n-1}) \xrightarrow{\phi} H_{*-1}(D^n),$$

we can break this up using the kernel and cokernel to get

$$0 \rightarrow \text{coker } \phi_m \rightarrow H_m(D^n, S^{n-1}) \rightarrow \ker \phi_{m-1} \rightarrow 0. \quad (1.4.1)$$

D^n is contractible and so we get

$$H_*(D^n) = \begin{cases} \mathbf{Z}, & * = 0, \\ 0, & * \neq 0. \end{cases}$$

We have $\phi: H_0(S^{n-1}) \rightarrow H_0(D^n) \cong \mathbf{Z} = \langle e_p \rangle$ given by $\phi(e_p) = e_p$. We see that $\ker \phi = \tilde{H}_*(S^{n-1})$ and $\text{coker } \phi = 0$. Now by looking at (1.4.1) we see that

$$0 \rightarrow \tilde{H}_*(S^n) = H_*(S^n, *) = H_*(D^n, S^{n-1}) \rightarrow \ker \phi = \tilde{H}_*(S^{n-1}) \rightarrow 0$$

is exact giving $\tilde{H}_*(S^n) \cong \tilde{H}_*(S^{n-1})$. The claim then follows by induction. □

Corollary 1.4.19.

$$S^n \sim S^m \implies m = n.$$

Example 1.4.20 (Chains generating $H_n(S^n, *)$). 1. Choose $f: \Delta^n \rightarrow \Delta^n$ a homeomorphism and let $e: \Delta^n \rightarrow \Delta^n$ be the identity map. Then $a_{n-1} = f_{\#}(de) \in C_*(S^{n-1})$ and we have $da_{n-1} = df_{\#}(de) = f_{\#}(d^2e) = 0$ and so a_{n-1} is closed.

2. Choose $g: (D^n/S^{n-1}, S^{n-1}/S^{n-1}) \rightarrow (S^n, *)$. Now let $b_n = g_\#(f_\#(e)) \in C_n(S^n, *)$. Then $db_n = g_\#(a_{n-1}) \in C_{n-1}(S^n, *)$ implies b_n is closed in $C_*(S^n, *)$. For example if $n = 2$ we are crushing the boundary of the 2-simplex to a point.

Proposition 1.4.21. $[a_n]$ generates $\tilde{H}_n(S^n) = \mathbf{Z}$ (statement (A_n)).
 $[b_n]$ generates $\tilde{H}_n(S^n, *) = \mathbf{Z}$ (statement (B_n)).

Proof. Statement (A_0) : $S^0 = \{p_+, p_-\}$, $de = e_{p_+} - e_{p_-}$ so de generates $\tilde{H}_0(S^0)$. We'll show that $A_{n-1} \implies B_n$. Here $\partial: H_n(S^n, *) = H_n(D^n, S^{n-1}) \xrightarrow{\sim} \tilde{H}_{n-1}(S^{n-1})$ (by 1.4.18). So it suffices to check that $\partial[b_n] = [a_{n-1}]$ \square

Definition 1.4.22 (Wedge product). If (X_i, p_i) are pointed spaces the **wedge product**

$$\bigvee_{i \in I} (X_i, p_i) \text{ is } \coprod_{i \in I} X_i / \{p_i : i \in I\}.$$

If X_i is such that for any $p, q \in X_i$ there exists a homeomorphism $f: X_i \rightarrow X_i$ with $f(p) = q$ we can drop p_i from the notation (for example in the case of X_i a connected manifold we can do this).

Example 1.4.23.

Corollary 1.4.24. If (X_i, p_i) are good pairs then

$$\tilde{H}_* \left(\bigvee_i (X_i, p_i) \right) = \bigoplus_i \tilde{H}_*(X_i).$$

Proof.

$$\begin{aligned} \tilde{H}_* \left(\bigvee_{i \in I} X_i \right) &\cong H_* \left(\bigvee_{i \in I} X_i, p \right) \cong H_* \left(\coprod_{i \in I} X_i, \{p_i : i \in I\} \right) \\ &\cong \bigoplus_{i \in I} H_*(X_i, p_i) \cong \bigoplus_{i \in I} \tilde{H}_*(X_i). \end{aligned}$$

\square

Example 1.4.25.

$$H_*(S^1 \vee S^2) = \begin{cases} \mathbf{Z}, & * = 0, 1, 2 \\ 0, & \text{otherwise.} \end{cases}$$

1.5 Subdivision and Excision

Definition 1.5.1. If $\mathcal{U} = \{U_i\}_{i \in I}$ is an open cover of X , let

$$C_n^{\mathcal{U}}(X) = \langle e_\sigma : \sigma : \Delta^n \rightarrow X, \text{ im } \sigma \subset U_i \text{ for some } i \rangle \subset C_n(X).$$

Observe that $\text{im } \sigma \subset U_i$ implies $\text{im } \sigma \circ F_j \subset U_i$ and so $C_*^{\mathcal{U}}$ is a subcomplex of C_* . Let $H_*^{\mathcal{U}}$ be the homology of this complex, then we have a map

$$i: C_*^{\mathcal{U}}(X) \hookrightarrow C_*(X).$$

Lemma 1.5.2 (Subdivision). $C_*: H_*^{\mathcal{U}}(X) \rightarrow H_*(X)$ is an isomorphism.

1.6 Degree and Orientations

1.7 Cell Complexes

Definition 1.7.1 (Attaching of cells). If $f: S^{n-1} \rightarrow X$ then

$$X \cup_f D^n = X \amalg D^n / \sim$$

is the space obtained by **attaching** an n -dimensional cell to X via the map f .

Example 1.7.2. If $X = \{p\}$ and $f: S^{n-1} \rightarrow X$ then

$$X \cup_f D^n \cong D^n / S^{n-1} \cong S^n.$$

Definition 1.7.3 (Finite cell complex). A 0-dimensional **finite cell complex** is a finite disjoint union of points.

A k -dimensional finite cell complex is a space obtained by attaching finitely many k -cells to a $(k-1)$ -dimensional finite cell complex.

Example 1.7.4.

Example 1.7.5. If a finite cell complex X has one 0-cell and one n -cell then $X \cong S^n$. Similarly if X has one 0-cell and kn -cells then $X \cong \bigvee_{i=1}^k S^n$.

Example 1.7.6. T^2 is a finite cell complex with one 0-cell, two 1-cells and one 2-cell.

Example 1.7.7. Any simplicial complex is a finite cell complex and any closed manifold can be given the structure of a finite cell complex.

Chapter 2

Cohomology and Products

2.1 Homology with Coefficients and Cohomology

2.1.1 Hom and \otimes for modules

Definition 2.1.1 (Tensor product of R -modules). Let M, N be R -modules. Then the **tensor product** $M \otimes N$ is the R -modules generated by all pairs $m \otimes n$ for $m \in M, n \in N$ modulo the relations:

1. $(m_1 + m_2) \otimes (n_1 + n_2) = \sum m_i \otimes n_j$.
2. $r(m \otimes n) = (rm) \otimes n = m \otimes (rn)$.

We have the following properties of this definition:

1. $(M_1 \oplus M_2) \otimes (N_1 \oplus N_2) = \bigoplus M_i \otimes N_j$.
2. $M \otimes N \cong N \otimes M$.
3. $M \otimes R = M$

Example 2.1.2 (Tensor products). 1. $R^n \otimes R^m \cong R^{mn}$.

2. Letting $R = \mathbf{Z}, \mathbf{Q} \otimes \mathbf{Z}/a \cong 0$.
3. $\mathbf{Z}/a \otimes \mathbf{Z}/b \cong \mathbf{Z}/(a, b)$.

If $f: M_1 \rightarrow M_2$ and $g: N_1 \rightarrow N_2$ then there is a map

$$\begin{aligned} f \otimes g: M_1 \otimes N_1 &\rightarrow M_2 \otimes N_2, \\ m \otimes n &\mapsto f(m) \otimes g(n). \end{aligned}$$

Example 2.1.3. If $f: R^n \rightarrow R^m$ is given by multiplication by $A \in \text{Mat}_{n \times m}(R)$ then $f \otimes 1_M: M^n \rightarrow M^m$ is given by multiplication by A .

Definition 2.1.4 (Hom).

$$\text{Hom}(M, N) = \{f: M \rightarrow N : f \text{ is } R\text{-linear}\}$$

is an R -module, via $(f + rg)(m) = f(m) + rg(m)$.

From this definition we see that

1. $\text{Hom}(\bigoplus M_i, \bigoplus N_j) \cong \bigoplus_{i,j} \text{Hom}(M_i, N_j)$.
2. $\text{Hom}(R, M) \cong M$.

Note however that we do not have $\text{Hom}(M, N) = \text{Hom}(N, M)$ as for example $\text{Hom}(\mathbf{Z}/2, \mathbf{Z}) = 0$ but $\text{Hom}(\mathbf{Z}, \mathbf{Z}/2) = \mathbf{Z}/2$.

Definition 2.1.5 (Dual module). Given an R -module M the **dual** of M is $M^* = \text{Hom}(M, R)$.

Now if we have $f: M \rightarrow N$ we get a map

$$f^*: \text{Hom}(N, O) \rightarrow \text{Hom}(M, O)$$

given by $f^*g = g \circ f$.

Example 2.1.6. If $f: R^n \rightarrow R^m$ is multiplication by A then

$$f^*: \text{Hom}(R^m, O) \cong O^m \rightarrow \text{Hom}(R^n, O) \cong O^n$$

is multiplication by A^\top .

2.1.2 Hom and \otimes for chain complexes

If (C, d) is a chain complex defined over R then so are $(C_* \otimes M, d \otimes 1_M) = C_* \otimes M$ and $(\text{Hom}(C_*, M), d^*) = \text{Hom}(C_*, M)$.

2.2 Notation

Symbol	Description	Page
		1