

Parity

MA842 at BU Spring 2019

Céline Maistret

February 7, 2019

1 Parity

These are notes for Céline Maistret's course MA842 at BU Spring 2019.

The course webpage is https://sites.google.com/view/cmaistret/teaching#h.p_BYGoPzU848FJ.

Course overview:

Main references:

1. Vlad
2. Silverman
3. Milne

1.1 Mordell-Weil

Lecture 4 5/2/2018

Remark 1.1 A homomorphism $\phi: \text{Gal}(\bar{K}/K) \rightarrow G$ for a finite group G is continuous if it comes from a finite Galois extension, i.e.

$$\exists F/K \text{ finite Galois, } \tilde{\phi}: \text{Gal}(F/K) \rightarrow G$$

s.t. ϕ is the composition $\text{Gal}(\bar{K}/K) \rightarrow \text{Gal}(F/K) \xrightarrow{\tilde{\phi}} G$. So $\phi(g)$ only cares about what g does to F .

Proposition 1.2 Let E/K be an elliptic curve

$$y^2 = (x - \alpha)(x - \beta)(x - \gamma)$$

for $P \in E(K)$ have $\frac{1}{2}P \in E(\bar{K})$ s.t. $\frac{1}{2}P \oplus \frac{1}{2}P = P$.

1. $K(\frac{1}{2}P)/K$ is a Galois extension and $\text{Gal}(K(\frac{1}{2}P)/K) = C_2 \times C_2$ from Lemma 1.

2.

$$\phi_P: \text{Gal}(\bar{K}/K) \rightarrow E(K)[2]$$

$$g \mapsto Q^g - Q = g\left(\frac{1}{2}P\right) - \frac{1}{2}P$$

is well defined and has kernel $\text{Gal}(K/K(\frac{1}{2}P))$.

3.

$$\phi: E(K)/2E(K) \rightarrow \text{Hom}_{cts}(\text{Gal}(\bar{K}/K), E(K)[2])$$

$$P \mapsto \phi_P$$

is well defined and injective. Now ϕ_P is continuous by 2. and so

$$\begin{aligned}\phi_{P \oplus Q}(g) &= g\left(\frac{1}{2}(P \oplus Q)\right) - \left(\frac{1}{2}P \oplus \frac{1}{2}Q\right) \\ &= g\left(\frac{1}{2}P\right) \oplus g\left(\frac{1}{2}Q\right) - \frac{1}{2}P \oplus \frac{1}{2}Q \\ &= \phi_P(g) \oplus \phi_Q(g)\end{aligned}$$

a homomorphism.

$$\phi_{2Q}(g) = g\left(\frac{1}{2}2Q\right) - \frac{1}{2}2(Q) = g(Q) - Q = 0$$

for all $g \in \text{Gal}(\bar{K}/K)$ if $Q \in E(K)$ so this is well defined. For injectivity:

$$\begin{aligned}\phi_P(g) = 0 &\implies g\left(\frac{1}{2}P\right) = \frac{1}{2}P \forall g \in \text{Gal}(\bar{K}/K) \\ &\implies \frac{1}{2}P \in E(K) \implies P \in 2E(K)\end{aligned}$$

which gives injectivity.

4.

$$\eta: \text{Hom}_{cts}(\text{Gal}(\bar{K}/K), E(K)[2]) \rightarrow K^\times/K^{\times 2} \times K^\times/K^{\times 2} \times K^\times/K^{\times 2}$$

$$\psi \mapsto \psi_\alpha, \psi_\beta, \psi_\gamma$$

$$\psi(g) \in \{0, (\alpha, 0)\} \subseteq E(K) \iff g \in \text{Gal}(\bar{K}/K(\sqrt{\psi_\alpha}))$$

then η is an injective homomorphism. It is an isomorphism to the subgroup of triples a, b, c s.t. $abc \in K^{\times 2}$. Proof:

$$\text{Hom}_{cts}(\text{Gal}(\bar{K}/K), C_2) \simeq K^\times/K^{\times 2}$$

with ψ s.t. $\ker \psi = \text{Gal}(\bar{K}/K\sqrt{d}) \leftrightarrow d$. It is an isomorphism:

$$\ker \psi_i = \text{Gal}(\bar{K}/K(\sqrt{d_i})), i = 1, 2$$

$$\ker \psi_1 \psi_2 = \text{Gal}(\bar{K}/K(\sqrt{d_1 d_2}))$$

Now apply this to $E(K)[2] = C_2 \times C_2$ to get an isomorphism to $K^\times/K^{\times 2} \times K^\times/K^{\times 2}$. Record this third homomorphism to get η .

5. If $P = (x_0, y_0) \in E(K)$ then

$$\eta(\phi_P) = (x_0 - \alpha, x_0 - \beta, x_0 - \gamma).$$

Proof sketch: If

$$E: y^2 = x^3 + Ax^2 + Bx$$

then for $Q = (x_0, y_0) \in E(K)$.

$$2Q = \left(\left(\frac{x_0 - B}{2y_0} \right)^2, \dots \right)$$

Hence if $2Q = P = (x_1, y_1)$ then $\sqrt{x_1} \in K(\frac{1}{2}P)$. So if

$$E: y^2 = (x - \alpha)(x - \beta)(x - \gamma)$$

then

$$P = (x_2, y_2)$$

then

$$\begin{aligned} \sqrt{x_2 - \alpha}, \sqrt{x_2 - \beta}, \sqrt{x_2 - \gamma} &\in K(\frac{1}{2}P) \\ K(\sqrt{x_2 - \alpha}), K(\sqrt{x_2 - \beta}), K(\sqrt{x_2 - \gamma}) &\subseteq K(\frac{1}{2}P) \\ \implies K(\frac{1}{2}P) &= K(\sqrt{x_2 - \alpha}, \sqrt{x_2 - \beta}, \sqrt{x_2 - \gamma}) \end{aligned}$$

Example 1.3 Let

$$E: y^2 = x(x - 1)(x + 1)$$

for $P \in E(\mathbf{Q})$, $\mathbf{Q}(\frac{1}{2}P)/\mathbf{Q}$ can only ramify at 2.

$$\mathbf{Q}(\frac{1}{2}P) \subseteq \mathbf{Q}(i, \sqrt{2})$$

$$P = (x_0, y_0) \mapsto x_0, x_0 - 1, x_0 + 1 \in \mathbf{Q}^\times / \mathbf{Q}^{\times 2}$$

is a homomorphism so $x_0, x_0 - 1, x_0 + 1$ are $\pm 1, \pm 2$ up to square.

x_0	$x_0 - 1$	$x_0 + 1$	rat?
1	1	1	1) rat
1	-1	-1	2) non-rat
1	2	2	1) rat
1	-2	-2	2) non-rat
-1	1	-1	2) non-rat
-1	-1	1	1) rat
-1	2	-1	2) non-rat
-1	-2	2	1) rat
2	1	2	3) non-rat
2	-1	-2	2) non-rat
2	2	1	4) rat
2	-2	-1	2) non-rat
-2	1	-2	rat
-2	-1	2	rat
-2	2	-1	rat
-2	-2	1	rat

Table 1.4: Images

1) The 2-torsion points $P = 0, (0, 0), (1, 0), (-1, 0) \in E(\mathbf{Q})$ give us some rows.
2) As we have $x_0 > -1$ we get $x_0 + 1 > 0$ so $x_0(x_0 - 1) > 0$ for the product to be a square (and hence > 0). 3) $x_0 = 2A^2$, $x_0 - 1 = B^2$, $x_0 + 1 = 2C^2$ with $A, B, C \in \mathbf{Q} \setminus \{0\}$. Let $A = m/n$ so $2m^2/n^2 - 1 = B^2$

$$2m^2 - n^2 = (Bn)^2$$

and

$$2m^2 + n^2 = 2(Cn)^2$$

if $m \equiv 0(2) \implies -1 \equiv \square \pmod{8}$ a contradiction.

$$m \equiv 1 \pmod{2} \implies m^2 \equiv 1 \pmod{8}.$$

So $2 - n^2 \equiv \square \pmod{8} \implies n^2 \equiv 1 \pmod{8}$

$$2 + n^2 \equiv 2 \pmod{8} \implies n^2 \equiv 0 \pmod{8}$$

$$|E(\mathbf{Q})/2E(\mathbf{Q})| = 4$$

$$|E(\mathbf{Q})[2]| = 4 \implies \text{rk} = 0$$

$$E(\mathbf{Q}) \cong E(\mathbf{Q})[2].$$

4) Use the group structure!

□

Theorem 1.5 Complete 2-decent. Let K be a field of characteristic 0 and

$$E: y^2 = (x - \alpha)(x - \beta)(x - \gamma), \alpha, \beta, \gamma \text{ distinct.}$$

The map

$$P \mapsto (x_0 - \alpha, x_0 - \beta, x_0 - \gamma)$$

replacing $x_0 - \alpha$ with $(x_0 - \beta)(x_0 - \gamma)$ if 0.

$$E(K)/2E(K) \rightarrow (K^\times/K^{\times 2})^3$$

Triples (a, b, c) that lie in the image satisfy $abc \in K^{\times 2}$. A triple a, b, c with $abc \in K^{\times 2}$ lies in the image iff it is in the image of $E(K)[2]$ or

$$cz_3^2 - \alpha + \gamma = az_1^2$$

$$cz_3^2 - \beta + \gamma = bz_1^2$$

is soluble with $z_i \in K^\times$. In which case

$$P = (az_1^2 + \alpha, \sqrt{abc}, z_1 z_2 z_3) \mapsto (a, b, c)$$

iii) If K is a number field and (a, b, c) is in the image then

$$K(\sqrt{a}, \sqrt{b}, \sqrt{c})/K$$

only ramifies at primes dividing $2(\alpha - \beta)(\alpha - \gamma)(\beta - \gamma)$.

Exercise 1.6

$$E: y^2 = x(x - 5)(x + 5).$$

Lecture 5 7/2/2018

Recall:

$$\phi: E(K)/2E(K) \rightarrow \text{Hom}_{cts}(G_K, E(K)[2])$$

$$P \mapsto \phi_P$$

where $\phi_P: \sigma \mapsto Q^\sigma - Q$ where $Q = 2P$. Which is well-defined and injective.

Elements of

$$\text{Hom}_{cts}(G_K, E[2]) \leftrightarrow a, b, c \in (K^\times/K^{\times 2}) \text{ s.t. } abc \in K^{\times 2}$$

$$(x_0, y_0) \mapsto (x_0 - \alpha, x_0 - \beta, x_0 - \gamma).$$

Lemma 1.7 Let $n \geq 1$

1.

$$\psi: E(K)/nE(K) \rightarrow \{K \subseteq F \subseteq \overline{K}\}$$

$$P \mapsto K(\frac{1}{n}P, E[n])$$

is well defined.

2. $K(\frac{1}{n}P, E[n])/K$ only ramifies at $\mathfrak{p} | n\Delta_E$.

3.

$$\text{Gal}(K(\frac{1}{n}P, E[n])/K) \leq \mathbf{Z}/n \times \mathbf{Z}/n$$

4. There are only finitely many fields satisfying 2. and 3. so $\text{im } \psi$ is finite.

To do descent, need more than ψ (i.e. injection).

Definition 1.8 Let G be a group and M a G -module then let

$$H^0(G, M) = M^G = \{m \in M : gm = m \forall g \in G\}$$

$$H^1(G, M) = \{\text{skew homs } G \rightarrow M\} / \{\text{skew homs } G \rightarrow M \text{ of the form } g \mapsto g(t) - t, t \in M\}.$$

◇

Remark 1.9 If G acts trivially on M then

$$H^0(G, M) = M$$

$$H^1(G, M) = \text{Hom}(G, M).$$

When G is profinite then we want that the skew homomorphisms factor through finite Galois groups. We will prove that

$$E(K)/nE(K) \hookrightarrow H^1(G_K, E[n]).$$

Theorem 1.10 If

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

is an exact sequence of G -modules then

$$0 \rightarrow H^0(G, A) \rightarrow H^0(G, B) \rightarrow H^0(G, C) \rightarrow H^1(G, A) \rightarrow H^1(G, B) \rightarrow H^1(G, C).$$

Lemma 1.11

1. ψ is finite-to-one (gives Mordell-Weil)

2. Let

$$\phi_P: G_K \rightarrow E[n]$$

$$\phi_P(gh) = \phi_P(g) + g\phi_P(h)$$

is a skew (or crossed) homomorphism. If $(\frac{1}{n}P)'$ is another choice of $\frac{1}{n}P$ and ϕ'_P is the corresponding skew homomorphism, then

$$\phi_P - \phi'_P$$

is of the form

$$g \mapsto T \ominus gT$$

where $T \in E[n]$.

3. ϕ_P factors through

$$\text{Gal}(K(\frac{1}{n}P, E[n])/K).$$

4.

$$\phi: E(K)/nE(K) \rightarrow Z/B$$

$$P \mapsto \phi_P$$

is an injective homomorphism. Where

$$Z = \{\text{skew homs } G_K \rightarrow E[n]\}$$

$$B = \{\text{skew homs } G_K \rightarrow E[n] \text{ of the form } g \mapsto T \ominus gT, T \in E[n]\}.$$

Proof.

1. There are finitely many skew homomorphisms

$$\text{Gal}(K(\frac{1}{n}P, E[n])/K) \rightarrow E[n]$$

and by 4.

$$P \mapsto \{\phi_P, K(\frac{1}{n}P, E[n])\}$$

is injective. So $\psi: P \mapsto K(\frac{1}{n}P, E[n])$ is finite to one by 3.

2.

$$\begin{aligned} \phi_P(gh) &= \frac{1}{n}P \ominus gh \frac{1}{n}P \\ &= \left(\left(\frac{1}{n}P \right) \ominus g \left(\frac{1}{n}P \right) \right) \oplus \left(g \left(\frac{1}{n}P \right) \ominus g(h \left(\frac{1}{n}P \right)) \right) \\ &= \phi_P \oplus g(\phi_P(h)). \end{aligned}$$

Remark: If $E[n] \subseteq E(K)$ then ϕ_P is a homomorphism. Recall for $n = 2$

$$\begin{aligned} \phi_P(gh) &= \frac{1}{2}P \ominus gh \left(\frac{1}{2}P \right) \\ &= \frac{1}{2}P \ominus h \left(\frac{1}{2}P \right) \oplus h \left(\frac{1}{2}P \right) \ominus g(h \left(\frac{1}{2}P \right)) \\ &= \phi_P(h) \oplus \phi_P(g) \end{aligned}$$

since $2h(\frac{1}{2}P) = h(P) = P$. Consider now

$$\frac{1}{n}P = \frac{1}{n}P' \oplus T$$

for some $T \in E[n]$

$$\begin{aligned} (\phi_P \ominus \phi'_P)(g) &= \phi_P(g) - \phi'_P(g) = \frac{1}{n}P \ominus g \left(\frac{1}{n}P \right) - \left[\left(\frac{1}{n}P \right) \oplus T \ominus g \left(\frac{1}{n}P \right) \oplus gT \right] \\ &= T \ominus gT. \end{aligned}$$

■

Take $G = G_K, B = E(\overline{K}), A = E[n], C = E(\overline{K})$ to get

$$0 \rightarrow E[n] \rightarrow E(\overline{K}) \xrightarrow{\cdot n} E(\overline{K}) \rightarrow 0$$

which gives the long exact sequence

$$\begin{aligned} 0 \rightarrow E(K)[n] \rightarrow E(K) \xrightarrow{\cdot n} E(K) \xrightarrow{\delta} H^1(G_K, E[n]) \rightarrow H^1(G_K, E(\bar{K})) \rightarrow \\ \implies E(K)/nE(K) \hookrightarrow H^1(G_K, E[n]). \end{aligned}$$

Problem:

$$H^1(G_K, E[n])$$

is infinite. What subgroup of

$$H^1(G_K, E[n])$$

do we land in?

Notation: When v is a place of K we have $G_{K_v} \subseteq G_K$, for any module M have $M^{G_K} \subseteq M^{G_{K_v}}$ and

$$\text{Res}: H^1(G_K, E[n]) \rightarrow H^1(G_{K_v}, E[n]).$$

We have from the theorem

$$\begin{array}{ccccccc} 0 & \longrightarrow & E(K)/nE(K) & \xrightarrow{\delta} & H^1(G_K, E[n]) & \longrightarrow & H^1(G_K, E(\bar{K})) \longrightarrow \\ & & & & \downarrow \text{Res} & & \downarrow \text{Res} \\ 0 & \longrightarrow & \prod_v E(K_v)/nE(K_v) & \xrightarrow{\delta} & \prod_v H^1(G_{K_v}, E[n]) & \longrightarrow & \prod_v H^1(G_{K_v}, E(\bar{K})) \longrightarrow \end{array}$$

we want to understand $\text{im } \delta$ i.e. the subgroup

$$\ker\{H^1(G_K, E[n]) \rightarrow H^1(G_K, E(\bar{K}))\}$$

this is as hard as finding $E(K)$, here is why:

Claim 1.12

$$H^1(G_K, E(\bar{K}))$$

corresponding to principal homogeneous spaces for E (genus 1 curves whose jacobian is E)

Finding

$$\ker\{H^1(G_K, E[n]) \rightarrow H^1(G_K, E(\bar{K}))\}$$

is equivalent to finding which PHS coming from H^1 have a rational point. ???

Hensels lemma

Let C be a curve

$$\text{Isom}(C) \hookrightarrow C(\bar{K}) \times \text{Aut}(c)$$

$$\tau_P \circ \alpha \hookrightarrow (P, \sigma)$$

$$\text{Twist}(E/K) \hookrightarrow H^1(G_K, \text{Isom}(C))$$

$$C \simeq_{\bar{K}} E$$

$$\text{PHS} \hookrightarrow H^1(G_K, E(\bar{K}))$$

C is a PHS for E iff E is the jacobian of C .