Singular Moduli

Alex J. Best

WIMP 2014

29/11/2014

### In this talk:

- Introduction
- 2 Background
- The Hilbert class field
- 4 Singular moduli
- Modern work
- 6 Conclusion

 $e^{\pi\sqrt{43}} \approx 884736743.999777466$ 

$$e^{\pi\sqrt{43}} \approx 884736743.999777466$$

$$e^{\pi\sqrt{67}} \approx 147197952743.999998662454$$

$$e^{\pi\sqrt{43}} \approx 884736743.999777466$$

$$e^{\pi\sqrt{67}} \approx 147197952743.999998662454$$

$$e^{\pi\sqrt{43}} \approx 884736743.999777466$$
  
 $\approx 12^3(9^2-1)^3+744-10^{-4}\cdot 2.225\dots$   
 $e^{\pi\sqrt{67}} \approx 147197952743.999998662454$   
 $\approx 12^3(21^2-1)^3+744-10^{-6}\cdot 1.337\dots$   
 $e^{\pi\sqrt{163}} \approx 262537412640768743.999999999999925007$   
 $\approx 12^3(231^2-1)^3+744-10^{-13}\cdot 7.499\dots$ 

# Some definitions

#### Definition

A finite Galois extension L|K is **abelian** extension if Gal(L|K) is abelian.

## Some definitions

#### Definition

A finite Galois extension L|K is **abelian** extension if Gal(L|K) is abelian.

Examples:

## Some definitions

#### Definition

A finite Galois extension L|K is **abelian** extension if Gal(L|K) is abelian.

Examples:

Non-examples:

# The ideal class group

#### Definition

The **ideal class group** of a number field K is the quotient

$$cl(\mathbf{Z}_K) = I(\mathbf{Z}_K)/P(\mathbf{Z}_K).$$

## The ideal class group

#### Definition

The ideal class group of a number field K is the quotient

$$cl(\mathbf{Z}_K) = I(\mathbf{Z}_K)/P(\mathbf{Z}_K).$$

 $cl(\mathbf{Z}_L)$  measures how far  $\mathbf{Z}_K$  is from having unique factorisation.

Let K be an **imaginary quadratic** number field, i.e.  $K = \mathbb{Q}(\sqrt{-n})$  for some  $n \in \mathbb{Z}_{\geq 1}$ .

Let K be an **imaginary quadratic** number field, i.e.  $K = \mathbb{Q}(\sqrt{-n})$  for some  $n \in \mathbb{Z}_{\geq 1}$ .

#### Definition

An extension L|K is **unramified** if for all prime ideals  $\mathfrak{p}$  of  $\mathbf{Z}_K$  we have a factorisation

$$\mathfrak{p} \, \mathsf{Z}_L = \mathfrak{P}_1 \, \mathfrak{P}_2 \cdots \mathfrak{P}_n$$

into distinct prime ideals  $\mathfrak{P}_i$  of  $\mathbf{Z}_L$ .

Let K be an **imaginary quadratic** number field, i.e.  $K = \mathbb{Q}(\sqrt{-n})$  for some  $n \in \mathbb{Z}_{\geq 1}$ .

#### Definition

An extension L|K is **unramified** if for all prime ideals  $\mathfrak p$  of  $\mathbf Z_K$  we have a factorisation

$$\mathfrak{p}\,\mathsf{Z}_L=\mathfrak{P}_1\,\mathfrak{P}_2\cdots\mathfrak{P}_n$$

into **distinct** prime ideals  $\mathfrak{P}_i$  of  $\mathbf{Z}_L$ .

#### Definition

The **Hilbert class field** of K is the maximal unramified abelian extension of K.

#### Definition

The **Hilbert class field** of K is the maximal unramified abelian extension of K.

#### **Examples**

# The Artin reciprocity theorem for the Hilbert class field

#### Theorem

If K is a number field and L is its Hilbert class field then

$$cl(\mathbf{Z}_K) \cong Gal(L|K).$$

# The j-invariant

Letting  $q = e^{2\pi iz}$  we have

$$j(z) = \frac{1}{q} + 744 + 196884q + 21493760q^{2} + 864299970q^{3} + 20245856256q^{4} + \cdots$$

# Closing remarks

• Singular moduli are not particularly complex objects in and of themselves.

# Closing remarks

- Singular moduli are not particularly complex objects in and of themselves.
- But their relation between different areas of mathematics ensures that they are still a research topic to this day.

#### Sources

I used some of the following when preparing this talk, and so they are probably good places to look to learn more about the topic:

• "Primes of the form  $x^2 + ny^2$ " – David A. Cox