# The (inescapable) *p*-adics

Alex J. Best

10/10/22

VU Master Seminar - Algebra

### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

### Example: Fibonacci

$$a_0 = 0, a_1 = 1 \text{ and } a_n = a_{n-1} + a_{n-2} \text{ for } n \ge k = 2$$
:

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377, 610, 987, 1597, 2584, 41

### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

### Example: Fibonacci

$$a_0 = 0$$
,  $a_1 = 1$  and  $a_n = a_{n-1} + a_{n-2}$  for  $n \ge k = 2$ :

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377, 610, 987, 1597, 2584, 4187, 1487

 $a_n$  grows exponentially.

### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

### Example: A periodic sequence

$$a_0 = 1, a_1 = 0$$
 with  $a_n = -a_{n-1} - a_{n-2}$ 

$$1, 0, -1, 1, 0$$

#### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

### Example: A periodic sequence

$$a_0 = 1, a_1 = 0$$
 with  $a_n = -a_{n-1} - a_{n-2}$ 

$$1, 0, -1, 1, 0$$

 $a_n$  is periodic now.

### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

#### Example: Natural numbers interlaced with zeroes

$$a_0 = 1, a_1 = 0, a_2 = 2, a_3 = 0$$
 with  $a_n = 2a_{n-2} - a_{n-4}$ 

1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13, 0, 14, 0, 12, 0, 13, 0, 14

#### Definition: Linear recurrence sequence

A linear recurrence sequence, is a sequence whose nth term is a linear combination of the previous k terms (for all  $n \ge k$ ).

#### Example: Natural numbers interlaced with zeroes

$$a_0 = 1$$
,  $a_1 = 0$ ,  $a_2 = 2$ ,  $a_3 = 0$  with  $a_n = 2a_{n-2} - a_{n-4}$ 

not periodic but the zeroes *do* have a regular repeating pattern.

## The ultimate question

### Question

What possible patterns are there for the zeroes of a linear recurrence sequence?

## The ultimate question

#### Question

What possible patterns are there for the zeroes of a linear recurrence sequence?

#### Observation

A linear recurrence sequence is the Taylor expansion around 0 of a rational function

$$\frac{a_1 + a_2 x + \dots + a_\ell x^\ell}{b_1 + b_2 x \dots + b_k x^k}$$

with  $b_1 \neq 0$  (so that the expansion makes sense).

### Example

$$\frac{x}{1-x-x^2}$$
.  $\leftrightarrow$  Fibonacci

### Example

$$\frac{x}{1-x-x^2}$$
.  $\leftrightarrow$  Fibonacci

$$\frac{1}{1+x+x^2}. \leftrightarrow 1, 0, -1, 1, 0, -$$

#### Example

$$\frac{x}{1-x-x^2}$$
.  $\leftrightarrow$  Fibonacci

$$\frac{1 - x - x^{2}}{1 + x + x^{2}} \longleftrightarrow 1, 0, -1,$$

$$\frac{1}{(1-x^2)^2}. \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0$$

#### Example

$$\frac{x}{1-x-x^2}. \leftrightarrow \mathsf{Fibonacci}$$
 
$$\frac{1}{1+x+x^2}. \leftrightarrow 1, 0, -1, 1, 0,$$

$$\frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0$$

 $\frac{(1+x)^3 - x^3}{(1+x)^5 - x^5} \leftrightarrow 1, -2, 3, -5, 10, -20, 35, -50, 50, 0, -175, 625, -1625, 3625, -7250, 13125, -21250, 29375, -29375, 0, 106250, -384375, 1006250, -2250000, 4500000,$ 

0, 106250, -384375, 1006250, -2250000, 4500000, 3  $-8140625, 13171875, -18203125, 18203125, \emptyset, -656$ 

# Consequences

#### Observation

The set of all linear recurrence sequences is a vector space! Hard to tell how the rule changes.

### Consequences

#### Observation

The set of all linear recurrence sequences is a vector space! Hard to tell how the rule changes.

We can always mess up a finite amount of behaviour. So assume  $a_n$  has infinitely many zeroes, what is the structure of the zero set?

#### Example

$$\frac{1}{(1-x^2)^2} - (1-x+2x^2+3x^4+4x^6) \leftrightarrow 0, 1, 0, 0, 0, 0, 0, 0, 5, 0, 6, 0, 7, 0, 8$$

#### Example

$$\frac{1}{(1-x^2)^2} - (1-x+2x^2+3x^4+4x^6) \leftrightarrow 0, 1, 0, 0, 0, 0, 0, 0, 5, 0, 6, 0, 7, 0, 8$$

#### Example

$$\frac{1}{(1-x^2)^2} - (1-x+2x^2+3x^4+4x^6) \leftrightarrow 0, 1, 0, 0, 0, 0, 0, 0, 5, 0, 6, 0, 7, 0, 8$$

#### Example

$$\frac{1}{(1-x^2)^2} - (1-x+2x^2+3x^4+4x^6) \leftrightarrow 0, 1, 0, 0, 0, 0, 0, 0, 5, 0, 6, 0, 7, 0, 8$$

$$\frac{1}{(1-x)^2} \leftrightarrow 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21$$

$$\frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13$$

$$\frac{1}{(1-x)^2} \leftrightarrow 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, \frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13$$

$$\frac{1}{(1-x)^2} \leftrightarrow 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, \frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13, \frac{1}{(1-x^4)^2} \leftrightarrow 1, 0, 0, 0, 2, 0, 0, 0, 3, 0, 0, 0, 4, 0, 0, 0, 5, 0, 0, 0, 6, 0, 0, 7, 0, 0$$

 $\frac{x}{(1-x^4)^2} \leftrightarrow 0, 1, 0, 0, 0, 2, 0, 0, 0, 3, 0, 0, 0, 4, 0, 0, 0, 5, 0, 0, 0, 6, 0, 0, 7, 0$ 

$$\frac{1}{(1-x)^2} \leftrightarrow 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21$$

$$\frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13$$

$$\frac{1+2x}{(1-x^4)^2} \leftrightarrow 1, 2, 0, 0, 2, 4, 0, 0, 3, 6, 0, 0, 4, 8, 0, 0, 5, 10, 0, 0, 6, 12, 0, 0, 7, 1$$

$$\frac{1}{(1-x)^2} \leftrightarrow 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21$$

$$\frac{1}{(1-x^2)^2} \leftrightarrow 1, 0, 2, 0, 3, 0, 4, 0, 5, 0, 6, 0, 7, 0, 8, 0, 9, 0, 10, 0, 11, 0, 12, 0, 13$$

$$\frac{1+2x}{(1-x^4)^2} \leftrightarrow 1, 2, 0, 0, 2, 4, 0, 0, 3, 6, 0, 0, 4, 8, 0, 0, 5, 10, 0, 0, 6, 12, 0, 0, 7, 1$$

Still has periodic zero set, all n congruent to 2,3 modulo 4.

Expand into partial fractions

$$\frac{p(x)}{q(x)} = \sum_{i=1}^{m} \sum_{j=1}^{n_j} \frac{r_{ij}}{(1 - \alpha_i x)^j}$$

Expand into partial fractions

$$\frac{p(x)}{q(x)} = \sum_{i=1}^{m} \sum_{j=1}^{n_j} \frac{r_{ij}}{(1 - \alpha_i x)^j}$$

do some math:

$$\sum_{n=0}^{\infty} \left( \sum_{i=1}^{m} \sum_{j=1}^{n_j} r_{ij} \binom{n+j-1}{j-1} \alpha_i^n \right) x^n$$

Expand into partial fractions

$$\frac{p(x)}{q(x)} = \sum_{i=1}^{m} \sum_{j=1}^{n_j} \frac{r_{ij}}{(1 - \alpha_i x)^j}$$

do some math:

$$\sum_{n=0}^{\infty} \left( \sum_{i=1}^{m} \sum_{j=1}^{n_j} r_{ij} \binom{n+j-1}{j-1} \alpha_i^n \right) x^n$$

Upshot: there are polynomials  $A_i(n)$  such that

$$a_n = \sum_{i=1}^m A_i(n)\alpha_i^n.$$

Like that formula for Fibonacci with the golden ratio in.

So  $a_n$  is an analytic function of n which has zeroes for infinitely many integer values.

So  $a_n$  is an analytic function of n which has zeroes for infinitely many integer values.

Like

$$sin(\pi x)!$$

So  $a_n$  is an analytic function of n which has zeroes for infinitely many integer values.

Like

$$sin(\pi x)!$$

#### Ridiculous suggestion

What if the integers were bounded? In that case infinitely many zeroes  $\implies$  the function is zero!

### What is bounded?

What if the integers were bounded?

How do we define boundedness?

#### **Definition: Absolute Values**

Let C be a commutative ring, an **absolute value** on C, is a function  $|\cdot|\colon R\to\mathbb{R}_{\geq 0}$  satisfying for all  $x,y\in C$ 

$$|x| = 0 \iff x = 0$$
  
 $|xy| = |x||y|$   
 $|x + y| \le |x| + |y|$ 

C

# Are there other absolute values for the integers?

$$|x| = 0 \iff x = 0$$
$$|xy| = |x||y|$$
$$|x + y| \le |x| + |y|$$

Property 2 implies that |1|=1 and  $|-1|^2=1$  so |-1|=1 also. So it remains to decide what happens for all primes  $p\in\mathbb{Z}$ .

We could set |x| = 1 for all  $x \neq 0$ , this is the **trivial absolute** value.

Or |x| = x for all positive x, this gives the usual absolute value.

## A strange absolute value

We can in fact define another absolute value  $|\cdot|_p$  for each prime p.

Pick a value  $\alpha = |p|_p < 1$ , and let  $|q|_p = 1$  for all other primes q.

Now we have that

$$|x+y| \le \max(|x|,|y|) \le |x|+|y|$$

#### Theorem: Ostrowski

The only nontrivial absolute values on  $\mathbb Q$  are

$$x \mapsto \operatorname{sgn}(x)x$$
 and  $|\cdot|_p$  for some prime p

With  $|\cdot|_p$  the integers are bounded!

With  $|\cdot|_p$  the integers are bounded! Are the functions

$$\sum_{i=1}^m A_i(n)\alpha_i^n$$

*p*-adic analytic functions of *n*?

With  $|\cdot|_p$  the integers are bounded! Are the functions

$$\sum_{i=1}^m A_i(n)\alpha_i^n$$

p-adic analytic functions of n?

#### **Problem**

The *p*-adic exponential function has finite radius of convergence.

With  $|\cdot|_p$  the integers are bounded! Are the functions

$$\sum_{i=1}^m A_i(n)\alpha_i^n$$

p-adic analytic functions of n?

#### **Problem**

The *p*-adic exponential function has finite radius of convergence.

#### The fix

Choose p so that  $|\alpha_i|_p = 1$  for all i, then  $\alpha_i^{p-1} = 1 + \lambda_i$  with  $|\lambda_i|_p \leq \frac{1}{p}$ . Now  $(\alpha_i^{p-1})^n$  is analytic!

Write n as r + (p-1)n' with  $0 \le r < p-1$ 

Write n as r + (p-1)n' with  $0 \le r < p-1$ , then

$$a_{n} = \sum_{i=1}^{m} A_{i}(n)\alpha_{i}^{n} = \sum_{i=1}^{m} A_{i}(r + (p-1)n')\alpha_{i}^{r+(p-1)n'}$$
$$= \sum_{i=1}^{m} A_{i}(r + (p-1)n')\alpha_{i}^{r}(\alpha_{i}^{(p-1)})^{n'}$$

for each fixed r this function of n' is analytic.

Write n as r + (p-1)n' with  $0 \le r < p-1$ , then

$$a_{n} = \sum_{i=1}^{m} A_{i}(n)\alpha_{i}^{n} = \sum_{i=1}^{m} A_{i}(r + (p-1)n')\alpha_{i}^{r+(p-1)n'}$$
$$= \sum_{i=1}^{m} A_{i}(r + (p-1)n')\alpha_{i}^{r}(\alpha_{i}^{(p-1)})^{n'}$$

for each fixed r this function of n' is analytic. Infinitely many zeroes for integer n means  $\exists r$  with infinitely many zeroes of the form r + (p-1)n'. So the function

$$\sum_{i=1}^{m} A_{i}(r+(p-1)n')\alpha_{i}^{r}(\alpha_{i}^{(p-1)})^{n'}$$

is identically zero, and all these  $a_n = 0$  when  $n \equiv r \pmod{p-1}$ .

#### **Finale**

#### Theorem: Skolem → Mahler → Lech

All except finitely many indices of the zeroes of a linear recurrence lie in a finite union of arithmetic progressions, i.e. they are all of the form nM+b for some  $b\in B\subset \{0,\ldots,M-1\}$ ,  $n\in \mathbb{N}$ .

#### **Finale**

#### Theorem: Skolem → Mahler → Lech

All except finitely many indices of the zeroes of a linear recurrence lie in a finite union of arithmetic progressions, i.e. they are all of the form nM+b for some  $b\in B\subset \{0,\ldots,M-1\}$ ,  $n\in \mathbb{N}$ .

