

Hi! 😁

Alex J. Best

5/28/2018

Boston University

Coleman integration

If C/\mathbf{R} is a curve, $P, Q \in C$, $\omega \in \Omega_C^1$ (e.g. $\frac{x dx}{y}$), we have a path integral

$$\int_P^Q \omega \in \mathbf{R}.$$

Coleman integration

If C/\mathbf{R} is a curve, $P, Q \in C$, $\omega \in \Omega_C^1$ (e.g. $\frac{x dx}{y}$), we have a path integral

$$\int_P^Q \omega \in \mathbf{R}.$$

What about if C/\mathbf{Q}_p ?

Coleman defined

$$\int_P^Q \omega \in \mathbf{Q}_p \quad \text{💣}$$

Coleman integration

If C/\mathbf{R} is a curve, $P, Q \in C$, $\omega \in \Omega_C^1$ (e.g. $\frac{x dx}{y}$), we have a path integral

$$\int_P^Q \omega \in \mathbf{R}.$$

What about if C/\mathbf{Q}_p ?

Coleman defined

$$\int_P^Q \omega \in \mathbf{Q}_p$$



a “path” integral, with cool properties



Coleman integration

If C/\mathbf{R} is a curve, $P, Q \in C$, $\omega \in \Omega_C^1$ (e.g. $\frac{xdx}{y}$), we have a path integral

$$\int_P^Q \omega \in \mathbf{R}.$$

What about if C/\mathbf{Q}_p ?

Coleman defined

$$\int_P^Q \omega \in \mathbf{Q}_p$$



a “path” integral, with cool properties



These can be explicitly computed in many cases!



Applications

Rational points: We can sometimes find ω so that

$$\text{Zeroes} \left(\int_{p_0}^x \omega \right) \supseteq C(\mathbf{Q})$$



Applications

Rational points: We can sometimes find ω so that

$$\text{Zeroes} \left(\int_{p_0}^x \omega \right) \supseteq C(Q) \quad \text{🕵️}$$

Heights:

Coleman-Gross introduced a height pairing on abelian varieties, it be decomposed as a sum of local terms, one of which is

$$h_p(D_1, D_2) = \int_{D_2} \omega_{D_1} \quad \text{🍏}$$

Applications

Rational points: We can sometimes find ω so that

$$\text{Zeroes} \left(\int_{p_0}^x \omega \right) \supseteq C(\mathbf{Q})$$



Heights:

Coleman-Gross introduced a height pairing on abelian varieties, it be decomposed as a sum of local terms, one of which is

$$h_p(D_1, D_2) = \int_{D_2} \omega_{D_1}$$



p -adic BSD:

Using the above height pairing one can define a p -adic regulator so that for a modular abelian variety A/\mathbf{Q} conjecturally

$$\mathcal{L}^*(A, 0) = \epsilon_p(A) \frac{|\text{III}(A/\mathbf{Q})| \text{Reg}_\gamma(A/\mathbf{Q}) \prod_v c_v}{|A(\mathbf{Q})_{\text{tors}}| |A^\vee(\mathbf{Q})_{\text{tors}}|}$$

