

Gamma Cross Sections

Aman LaChapelle

University of Chicago

October 21, 2015

Gamma Cross Sections

PHA spectra

- Gaussian vs Lorentzian

- Distribution Width

- Error Propagation

Countrates

- Na

- Ba

- Comparison to Literature

- Error Propagation


References

Gaussian Fits

$$f(x) = \frac{N}{\sigma\sqrt{2\pi}} e^{\frac{-(x-\mu)^2}{2\sigma^2}} + Ax + B$$

We fit all the peaks to this function because we didn't trust the ROI settings and their consistency.

Gaussian Fits



An example fit.

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na

Ba

Comparison to Literature

Error Propagation

References

Gaussian vs Lorentzian

Why don't we fit to

$$L(x) = \frac{1}{\pi} \frac{\frac{1}{2}\Gamma}{(x - x_0)^2 + \frac{1}{2}\Gamma^2}?$$

Because the errors are counting errors - they act on \bar{x} like $\sigma_{\bar{x}}$, the standard deviation of the mean. And since they are counting errors, they go like \sqrt{N} which is the SDOM.

They would both fit, but the Gaussian fits better.

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na


Ba

Comparison to Literature

Error Propagation

References

Distribution Width



The distribution should be a delta function - it isn't because of the detector resolution

Distribution Width

Energy (keV)	Γ (channels)	Γ/E
31	8.36	26.9%
81	10.18	12.6%
356	30.25	8.49%
511	10.68	2.09%
1270	16.69	1.31%

Not perfect resolution - this is our σ (up to $\Gamma/\sqrt{2}\sqrt{\ln(2)}$).

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na

Ba

Comparison to Literature

Error Propagation

References

Error Propagation of the PHA fits

Since we performed fits, the error of the i th parameter is just:

$$\sigma_i = \sqrt{\text{cov}(i, i)}$$

Because our largest δt was less than 0.001% the uncertainty in parameter i is just σ_i

Covariance Matrix

Scipy and Numpy return the covariance matrix, but here is some information on what it is:

$$X_i = (X_i - E(X_i))(X_i - E(X_i))^T$$

$$\Sigma = \frac{1}{N} \sum_i X_i$$

$$\Sigma_{ij} \equiv \text{cov}(i, j)$$

with each X_i being the squared difference between the datum and the model-estimated parameter - each parameter is expressed as a vector here.

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na

Ba

Comparison to Literature

Error Propagation

References



Fitting to find λ

We fit to the function

$$R(x) = R_0 e^{-\lambda x} + C$$

to extract our values for λ

Fitting to find λ



Energy	λ	$\delta\lambda$
511 keV	0.19 cm^{-1}	0.03 cm^{-1}
1.27 MeV	0.09 cm^{-1}	0.05 cm^{-1}

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na

Ba

Comparison to Literature

Error Propagation

References


Fitting to find λ

We fit to the function

$$R(x) = R_0 e^{-\lambda x} + R'_0 e^{-\tau x} + C$$

with λ being the compton-like attenuation and τ being the photoelectric-like attenuation.

Fitting to find λ



Energy	λ	$\delta\lambda$
31 keV	2.9 cm^{-1}	0.1 cm^{-1}
81 keV	0.513 cm^{-1}	0.06 cm^{-1}
356 keV	0.24 cm^{-1}	0.04 cm^{-1}

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na


Ba

Comparison to Literature

Error Propagation

References

Comparison to Literature



Energy	$\lambda \pm \delta\lambda \text{ (cm}^{-1}\text{)}$	$\lambda_{calc} \pm \delta\lambda_{calc} \text{ (cm}^{-1}\text{)}$	Confidence Level
31 keV	2.9 ± 0.1	2.98 ± 0.03	57.6%
81 keV	0.513 ± 0.06	0.533 ± 0.03	66.8%
356 keV	0.24 ± 0.04	0.3 ± 0.1	79.0%
511 keV	0.19 ± 0.03	0.22 ± 0.05	42.1%
1.27 MeV	0.09 ± 0.05	0.15 ± 0.07	33.8%

So we have pretty good confidence in our data, I think we did ok. The NIST values have uncertainties calculated based on how far the tabulated energy was from our energy and estimating the slope of the area immediately around the point of interest.

Table of Contents

PHA spectra

Gaussian vs Lorentzian

Distribution Width

Error Propagation

Countrates

Na

Ba

Comparison to Literature

Error Propagation

References

Error Propagation of the Countrate Fits

We use the same technique as we did with the PHA spectra to calculate the uncertainty in our parameters. A sample of the data used to calculate the countrates:

Ba-133 Countrates and Uncertainty

Thickness (mm)	N1 (count/s)	N2 (count/s)	N3 (count/s)	$\delta N1$	$\delta N2$	$\delta N3$
56	9.044	9.910	39.361	1.915%	4.832%	1.959%
40	13.480	20.943	57.620	2.632%	3.389%	2.156%
...
2	303.314	145.682	156.056	1.094%	1.302%	2.127%
1	391.492	152.985	155.885	1.032%	1.320%	2.074%

Our error bars in the countrate data are exactly the percent error you see above.

Systematic Uncertainty

We noticed that our data (for λ) was consistently lower than the NIST values, so we tried to fit it and the NIST values to:*

$$f(x) = Ae^{Bx} + Ce^{Dx} + E.$$

We found that the difference between our values and the NIST values was in E - it was just an offset. Furthermore we got a value for E of 0.020 ± 0.003 and so we added that 0.02 to all our uncertainties throughout for λ .

* not shown

References



Physics.NIST.gov - Table of XRay Mass Attenuation Coefficients
<http://physics.nist.gov/PhysRefData/XrayMassCoef/ElemTab/z13.html>



University of Chicago, PHYS 211 Lab Manual - P211 Wiki



An Introduction to Error Analysis - John Taylor