


# Hardy-Muckenhoupt Bounds for Laplacian Eigenvalues

Gary L. Miller   Noel J. Walkington   Alex L. Wang

Carnegie Mellon University


APPROX 2019



- 1 Introduction: Laplacians, eigenvalues, why do we care?
- 2 Bounding  $\lambda_2$  in terms of graph structure
- 3 What is  $\Psi_2$ ?
- 4 Summary


- 1 Introduction: Laplacians, eigenvalues, why do we care?
- 2 Bounding  $\lambda_2$  in terms of graph structure
- 3 What is  $\Psi_2$ ?
- 4 Summary

# Graphs and the Laplacian




- Weighted connected graph,  $G = (V, E, \mu, \kappa)$

# Graphs and the Laplacian



- Weighted connected graph,  $G = (V, E, \mu, \kappa)$
- Mass,  $\mu \in \mathbb{R}_{>0}^V$

# Graphs and the Laplacian



- Weighted connected graph,  $G = (V, E, \mu, \kappa)$
- Mass,  $\mu \in \mathbb{R}_{>0}^V$
- Edge weights,  $\kappa \in \mathbb{R}_{>0}^E$

# Graphs and the Laplacian

- Laplacian,  $L = D - A$

# Graphs and the Laplacian

- Laplacian,  $L = D - A$
- Degree matrix,  $D = \text{diag}(d_v)$



# Graphs and the Laplacian

- Laplacian,  $L = D - A$
- Degree matrix,  $D = \text{diag}(d_v)$
- Adjacency matrix,  $A_{u,v} = \kappa_{u,v}$

# Graphs and the Laplacian

- Laplacian,  $L = D - A$
- Degree matrix,  $D = \text{diag}(d_v)$  ,
- Adjacency matrix,  $A_{u,v} =$

$$d_v = \sum_{u \sim v} \kappa_{u,v}$$

$$\kappa_{u,v}$$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$(Lx)_v$$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$(Lx)_v = (Dx)_v - (Ax)_v$$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$\begin{aligned}(Lx)_v &= (Dx)_v - (Ax)_v \\ &= \left( \sum_{u \sim v} \kappa_{u,v} \right) x_v - \sum_{u \sim v} \kappa_{u,v} x_u\end{aligned}$$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$\begin{aligned}(Lx)_v &= (Dx)_v - (Ax)_v \\&= \left( \sum_{u \sim v} \kappa_{u,v} \right) x_v - \sum_{u \sim v} \kappa_{u,v} x_u \\&= \sum_{u \sim v} \kappa_{u,v} (x_v - x_u)\end{aligned}$$

# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$\begin{aligned}(Lx)_v &= (Dx)_v - (Ax)_v \\&= \left( \sum_{u \sim v} \kappa_{u,v} \right) x_v - \sum_{u \sim v} \kappa_{u,v} x_u \\&= \sum_{u \sim v} \kappa_{u,v} (x_v - x_u)\end{aligned}$$

- As a quadratic form

$$x^\top Lx$$



# Understanding the Laplacian

- Let  $x \in \mathbb{R}^V$
- As a linear map

$$\begin{aligned}(Lx)_v &= (Dx)_v - (Ax)_v \\&= \left( \sum_{u \sim v} \kappa_{u,v} \right) x_v - \sum_{u \sim v} \kappa_{u,v} x_u \\&= \sum_{u \sim v} \kappa_{u,v} (x_v - x_u)\end{aligned}$$

- As a quadratic form

$$\begin{aligned}x^\top Lx &= \dots \\&= \sum_{(u,v) \in E} \kappa_{u,v} (x_u - x_v)^2\end{aligned}$$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

- $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{|V|}$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

- $0 \leq \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{|V|}$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

- $0 = \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{|V|}$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

- $0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{|V|}$

# Eigenvalues associated with $L$

- Mass matrix,  $M = \text{diag}(\mu_v)$
- Generalized eigenvalue equation

$$Lx = \lambda Mx$$

- $0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \leq \lambda_{|V|}$
- The Neumann Eigenvalue



# Eigenvalues associated with $L$

- Setting  $\mu_v = d_v$  gives normalized Laplacian,  $\lambda_2$  controls mixing rate of random walks

# Eigenvalues associated with $L$

- Setting  $\mu_v = d_v$  gives normalized Laplacian,  $\lambda_2$  controls mixing rate of random walks
- Numerical linear algebra?

# Eigenvalues associated with $L$


- Setting  $\mu_v = d_v$  gives normalized Laplacian,  $\lambda_2$  controls mixing rate of random walks
- Numerical linear algebra?
- Goal: give good bounds for  $\lambda_2$  in terms of graph structure

- 1 Introduction: Laplacians, eigenvalues, why do we care?
- 2 Bounding  $\lambda_2$  in terms of graph structure
- 3 What is  $\Psi_2$ ?
- 4 Summary

# Cuts and Cheeger's inequality

- Sparsest cut,  $\Phi$


$$\Phi(G) = \min_A \left\{ \frac{\sum_{e \in E(A, \bar{A})} \kappa_e}{\min(\mu(A), \mu(\bar{A}))} \mid A, \bar{A} \neq \emptyset \right\}$$



# Cuts and Cheeger's inequality

- Sparsest cut,  $\Phi$

$$\Phi(G) = \min_A \left\{ \frac{\sum_{e \in E(A, \bar{A})} \kappa_e}{\min(\mu(A), \mu(\bar{A}))} \mid A, \bar{A} \neq \emptyset \right\}$$



## Theorem (Cheeger's Inequality)

If  $\mu_v = d_v$ ,

$$\frac{\Phi^2}{2} \leq \lambda_2 \leq 2\Phi.$$

# Our work

- Neumann content,  $\Psi_2$


$$\Psi_2(G) = \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A,B)}{(\mu(A)^{-1} + \mu(B)^{-1})^{-1}} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$



# Our work

- Neumann content,  $\Psi_2$

$$\Psi_2(G) \approx \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A,B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$





# Our work

- Neumann content,  $\Psi_2$

$$\Psi_2(G) \approx \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$



- $\kappa_{\text{eff}}(A, \bar{A}) = \sum_{e \in E(A, \bar{A})} \kappa_e$

# Our work

- Neumann content,  $\Psi_2$

$$\Psi_2(G) \approx \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$



- $\kappa_{\text{eff}}(A, \bar{A}) = \sum_{e \in E(A, \bar{A})} \kappa_e$

## Theorem ([MWW19])


*Let  $G$  be a weighted connected graph. Then*

$$\frac{\Psi_2}{4} \leq \lambda_2 \leq \Psi_2.$$

*Furthermore, all constants are tight.*

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$



# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$



- 

$$\Psi_2 \approx \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A,B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- 

$$\Psi_2 \approx \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A,B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\Psi_2 \approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right)\end{aligned}$$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right) \\ &= \min_{1 \leq a \leq n} \left( \frac{1}{a(2n + 1 - 2a)} \right)\end{aligned}$$



# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right) \\ &= \min_{1 \leq a \leq n} \left( \frac{1}{a(2n + 1 - 2a)} \right) \approx \frac{1}{n^2}.\end{aligned}$$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$



- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right) \\ &= \min_{1 \leq a \leq n} \left( \frac{1}{a(2n + 1 - 2a)} \right) \approx \frac{1}{n^2}.\end{aligned}$$

- [MWW19]  $\implies \lambda_2 \approx \frac{1}{n^2}$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$




- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right) \\ &= \min_{1 \leq a \leq n} \left( \frac{1}{a(2n + 1 - 2a)} \right) \approx \frac{1}{n^2}.\end{aligned}$$

- [MWW19]  $\implies \lambda_2 \approx \frac{1}{n^2}$
- $\Phi \approx 1/n$

# A first example

- Let  $G$  be a path graph on  $2n$  vertices with  $\kappa = \mu = 1$



- $\kappa_{\text{eff}}(a, b) = 1/(b - a)$ , can rewrite

$$\begin{aligned}\Psi_2 &\approx \min_{A, B} \left\{ \frac{\kappa_{\text{eff}}(A, B)}{\min(\mu(A), \mu(B))} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\} \\ &= \min_{1 \leq a < b \leq 2n} \left( \frac{1/(b - a)}{\min(a, 2n + 1 - b)} \right) \\ &= \min_{1 \leq a \leq n} \left( \frac{1}{a(2n + 1 - 2a)} \right) \approx \frac{1}{n^2}.\end{aligned}$$

- [MWW19]  $\implies \lambda_2 \approx \frac{1}{n^2}$
- $\Phi \approx 1/n$
- Cheeger's inequality  $\implies \frac{1}{n^2} \lesssim \lambda_2 \lesssim \frac{1}{n}$

## Related work

- Miclo [Mic99] studied  $\Psi$  for weighted path graphs with Dirichlet boundary conditions

$$\frac{\Psi}{4} \leq \lambda \leq 2\Psi$$

## Related work

- Miclo [Mic99] studied  $\Psi$  for weighted path graphs with Dirichlet boundary conditions

$$\frac{\Psi}{4} \leq \lambda \leq 2\Psi$$

- Schild [Sch18] studied  $\rho$  in the context of the normalized Laplacian

$$\frac{\rho}{25600} \leq \lambda_2 \leq 2\rho$$

## Related work

- Miclo [Mic99] studied  $\Psi$  for weighted path graphs with Dirichlet boundary conditions

$$\frac{\Psi}{4} \leq \lambda \leq 2\Psi$$

- Schild [Sch18] studied  $\rho$  in the context of the normalized Laplacian

$$\frac{\rho}{25600} \leq \lambda_2 \leq 2\rho$$

### Theorem ([MWW19])

Let  $G$  be a weighted connected graph. Then


$$\frac{\Psi_2}{4} \leq \lambda_2 \leq \Psi_2.$$

Furthermore, all constants are tight.

- 1 Introduction: Laplacians, eigenvalues, why do we care?
- 2 Bounding  $\lambda_2$  in terms of graph structure
- 3 What is  $\Psi_2$ ?
- 4 Summary




# Graphs with two vertices




- Can compute

$$\lambda_2 = \frac{\kappa}{(\mu_A^{-1} + \mu_B^{-1})^{-1}} = \Psi_2.$$

# Graphs with more vertices




# Graphs with more vertices




- Pick  $1 \leq a < b \leq n$

# Graphs with more vertices




- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$


# Graphs with more vertices




- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$
- Consider graph  $G_{a,b}$  with two vertices:



# Graphs with more vertices




- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$
- Consider graph  $G_{a,b}$  with two vertices:




- $\lambda_2(G_{a,b}) = \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}}$

# Graphs with more vertices




- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$
- Consider graph  $G_{a,b}$  with two vertices:




- $\lambda_2(G_{a,b}) = \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}}$
- Then,

$$\Psi_2 = \min_{a < b} \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}}$$

# Graphs with more vertices



- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$
- Consider graph  $G_{a,b}$  with two vertices:




- $\lambda_2(G_{a,b}) = \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}}$
- Then,


$$\Psi_2 = \min_{a < b} \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}} = \min_{a < b} \lambda_2(G_{a,b})$$



# Graphs with more vertices



- Pick  $1 \leq a < b \leq n$
- $A = \{1, \dots, a\}$ ,  $B = \{b, \dots, n\}$
- Consider graph  $G_{a,b}$  with two vertices:



- $\lambda_2(G_{a,b}) = \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}}$
- Then,

$$\Psi_2 = \min_{a < b} \frac{\kappa_{\text{eff}}(A, B)}{(\mu_A^{-1} + \mu_B^{-1})^{-1}} = \min_{a < b} \lambda_2(G_{a,b})$$

$\Psi_2$  is  $\lambda_2$  for the best two-vertex approximation of  $G$

- 1 Introduction: Laplacians, eigenvalues, why do we care?
- 2 Bounding  $\lambda_2$  in terms of graph structure
- 3 What is  $\Psi_2$ ?
- 4 Summary

# Summary

- Neumann content

$$\Psi_2 = \min_{A,B} \left\{ \frac{\kappa_{\text{eff}}(A,B)}{(\mu(A)^{-1} + \mu(B)^{-1})^{-1}} \mid A, B \neq \emptyset, A \cap B = \emptyset \right\}$$

- Showed

$$\frac{\Psi_2}{4} \leq \lambda_2 \leq \Psi_2$$

- Generalizations
- Future work: approximation algorithms?

Thanks for listening!

Questions?

Gary L. Miller, Noel J. Walkington, and Alex L. Wang.  
“Hardy-Muckenhoupt Bounds for Laplacian Eigenvalues”. In:  
*Approximation, Randomization, and Combinatorial Optimization.  
Algorithms and Techniques (APPROX/RANDOM 2019)*. Vol. 145.  
2019, 8:1–8:19

# References I



Laurent Miclo. “An example of application of discrete Hardy’s inequalities”. In: *Markov Process. Related Fields* 5.3 (1999), pp. 319–330.



Aaron Schild. “A Schur Complement Cheeger Inequality”. In: *arXiv preprint arXiv:1811.10834* (2018).