


New notions of simultaneous diagonalizability of quadratic forms with applications to QCQPs

Alex L. Wang, CMU Theory Lunch, Apr. 21



Joint work with Rujun Jiang, Fudan University

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- $$\begin{aligned} \inf_x \quad & q_1(x) \\ \text{s.t.} \quad & q_i(x) = 0, \forall i = 2, \dots, m \end{aligned}$$

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- $$\begin{aligned} \inf_x \quad & q_1(x) \\ \text{s.t.} \quad & q_i(x) = 0, \forall i = 2, \dots, m \end{aligned}$$
- Max-Cut

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- $$\begin{aligned} \inf_x \quad & q_1(x) \\ \text{s.t.} \quad & q_i(x) = 0, \forall i = 2, \dots, m \end{aligned}$$
- Max-Cut, Max-Clique

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- $$\begin{aligned} \inf_x \quad & q_1(x) \\ \text{s.t.} \quad & q_i(x) = 0, \forall i = 2, \dots, m \end{aligned}$$
- Max-Cut, Max-Clique, binary programming

Quadratically constrained quadratic programs (QCQPs)

- $q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$ quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- $$\begin{aligned} \inf_x \quad & q_1(x) \\ \text{s.t.} \quad & q_i(x) = 0, \forall i = 2, \dots, m \end{aligned}$$

- Max-Cut, Max-Clique, binary programming, polynomial optimization

Diagonal QCQPs

- QCQPs where $\{A_i\}$ are diagonal matrices

$$x^\top A_i x = \sum_{j=1}^n (A_i)_{j,j} x_j^2$$

Diagonal QCQPs

- QCQPs where $\{A_i\}$ are diagonal matrices

$$x^\top A_i x = \sum_{j=1}^n (A_i)_{j,j} x_j^2$$

- Nicer?

Diagonal QCQPs

- QCQPs where $\{A_i\}$ are diagonal matrices

$$x^\top A_i x = \sum_{j=1}^n (A_i)_{j,j} x_j^2$$

- Nicer?
- SDP relaxations more tractable¹

¹[Ben-Tal, den Hertog 14], [Jiang, Li 16], [Le, Nguyen 20]

Diagonal QCQPs

- QCQPs where $\{A_i\}$ are diagonal matrices

$$x^\top A_i x = \sum_{j=1}^n (A_i)_{j,j} x_j^2$$

- Nicer?
- SDP relaxations more tractable¹
- Better understanding of exactness of relaxations²

¹[Ben-Tal, den Hertog 14], [Jiang, Li 16], [Le, Nguyen 20]

²[Burer, Ye 19], [W and Kılınç-Karzan 21]

Diagonal QCQPs

- QCQPs where $\{A_i\}$ are diagonal matrices

$$x^\top A_i x = \sum_{j=1}^n (A_i)_{j,j} x_j^2$$

- Nicer?
- SDP relaxations more tractable¹
- Better understanding of exactness of relaxations²
- Black-box global solvers seem to perform better

¹[Ben-Tal, den Hertog 14], [Jiang, Li 16], [Le, Nguyen 20]

²[Burer, Ye 19], [W and Kılınç-Karzan 21]

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist invertible $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist **invertible** $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist **invertible** $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

- $x = Py$

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist **invertible** $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

- $x = Py \implies q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist **invertible** $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

- $x = Py \implies q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$
 $= y^\top (P^\top A_i P) y + 2b_i^\top P y + c_i$

From a general QCQP to a diagonal QCQP

- Given a QCQP, can we rewrite it as a diagonal QCQP?
- Given $\{A_i\} \subseteq \mathbb{S}^n$, does there exist **invertible** $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

- $x = Py \implies q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$
 $= y^\top (P^\top A_i P) y + 2b_i^\top P y + c_i$

Definition

Such sets $\{A_i\} \subseteq \mathbb{S}^n$ are **simultaneously diagonalizable via congruence** (SDC)

Outline

1 Introduction: QCQPs and diagonalization

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC?

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments
- 5 Conclusion: additional work, future directions

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments
- 5 Conclusion: additional work, future directions

SDC: What is known?

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is SDC if there exists invertible $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

[Horn, Johnson 12]

SDC: What is known?

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is SDC if there exists invertible $P \in \mathbb{R}^{n \times n}$:

$$P^\top A_i P = D_i, \quad \forall i$$

Theorem

If A is invertible. Then

$$\{A, B\} \text{ SDC} \iff A^{-1}B \text{ diagonalizable, real spectrum}$$

[Horn, Johnson 12]

SDC: First examples

- $\{A\}$ is SDC

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^T A P = \Lambda$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top AP = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top AP = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$A^{-1/2}BA^{-1/2}$$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top AP = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$U^\top \left(A^{-1/2} B A^{-1/2} \right) U = \Lambda$$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top A P = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$\left(A^{-1/2}U\right)^\top B \left(A^{-1/2}U\right) = \Lambda$$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top A P = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$\left(A^{-1/2}U\right)^\top B \left(A^{-1/2}U\right) = \Lambda$$

$$\left(A^{-1/2}U\right)^\top A \left(A^{-1/2}U\right) = U^\top U = I$$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top A P = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$\left(A^{-1/2}U\right)^\top B \left(A^{-1/2}U\right) = \Lambda$$

$$\left(A^{-1/2}U\right)^\top A \left(A^{-1/2}U\right) = U^\top U = I$$

- Let $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top A P = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$\left(A^{-1/2}U\right)^\top B \left(A^{-1/2}U\right) = \Lambda$$

$$\left(A^{-1/2}U\right)^\top A \left(A^{-1/2}U\right) = U^\top U = I$$

- Let
$$A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$$
$$A^{-1}B = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}$$

SDC: First examples

- $\{A\}$ is SDC \iff Spectral theorem $P^\top A P = \Lambda$
- If $A \succ 0$, then $\{A, B\}$ is SDC:

$$\left(A^{-1/2}U\right)^\top B \left(A^{-1/2}U\right) = \Lambda$$

$$\left(A^{-1/2}U\right)^\top A \left(A^{-1/2}U\right) = U^\top U = I$$

- Let $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$

$$A^{-1}B = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}$$

eigenvalues of $A^{-1}B = \{\pm i\} \implies \{A, B\}$ not SDC

SDC: Revisited

- $\{A_i\}$ is SDC $\iff \exists \{\ell_1, \dots, \ell_n\} \subseteq \mathbb{R}^n :$
basis
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$

SDC: Revisited

- $\{A_i\}$ is SDC $\iff \exists \{\ell_1, \dots, \ell_n\} \subseteq \mathbb{R}^n :$
basis
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$

- Given $\{A, B\}$, how many $\{\ell_1, \ell_2, \dots\}$ do we need

$$A = \sum_j \mu_j \ell_j \ell_j^\top, \quad B = \sum_j \lambda_j \ell_j \ell_j^\top?$$

SDC: Revisited

- $\{A_i\}$ is SDC $\iff \exists \{\ell_1, \dots, \ell_n\} \subseteq \mathbb{R}^n :$
basis
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$

- Given $\{A, B\}$, how many $\{\ell_1, \ell_2, \dots\}$ do we need

$$A = \sum_j \mu_j \ell_j \ell_j^\top, \quad B = \sum_j \lambda_j \ell_j \ell_j^\top?$$

Naively: $2n$

SDC: Revisited

- $\{A_i\}$ is SDC $\iff \exists \{\ell_1, \dots, \ell_n\} \subseteq \mathbb{R}^n :$
basis
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$

- Given $\{A, B\}$, how many $\{\ell_1, \ell_2, \dots\}$ do we need

$$A = \sum_j \mu_j \ell_j \ell_j^\top, \quad B = \sum_j \lambda_j \ell_j \ell_j^\top?$$

Naively: $2n$

Sneak peek: $n + 1$

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments
- 5 Conclusion: additional work, future directions

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is d -Restricted SDC if there exists $\{\bar{A}_i\} \subseteq \mathbb{S}^{n+d}$ SDC

$$\bar{A}_i = \begin{pmatrix} A_i & * \\ * & * \end{pmatrix}$$

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is d -Restricted SDC if there exists $\{\bar{A}_i\} \subseteq \mathbb{S}^{n+d}$ SDC

$$\bar{A}_i = \begin{pmatrix} A_i & * \\ * & * \end{pmatrix}$$

-

$$x^\top A_i x = \begin{pmatrix} x \\ 0 \end{pmatrix}^\top \begin{pmatrix} A_i & * \\ * & * \end{pmatrix} \begin{pmatrix} x \\ 0 \end{pmatrix}$$

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is d -Restricted SDC if there exists $\{\bar{A}_i\} \subseteq \mathbb{S}^{n+d}$ SDC

$$\bar{A}_i = \begin{pmatrix} A_i & * \\ * & * \end{pmatrix}$$

- $$x^\top A_i x = \begin{pmatrix} x \\ 0 \end{pmatrix}^\top \begin{pmatrix} A_i & * \\ * & * \end{pmatrix} \begin{pmatrix} x \\ 0 \end{pmatrix}$$
- $$\inf_x \begin{pmatrix} x \\ 0 \end{pmatrix}^\top \bar{A}_1 \begin{pmatrix} x \\ 0 \end{pmatrix} + \dots$$

s.t.
$$\begin{pmatrix} x \\ 0 \end{pmatrix}^\top \bar{A}_i \begin{pmatrix} x \\ 0 \end{pmatrix} + \dots = 0, \forall i = 2, \dots, m$$

Definition

$\{A_i\} \subseteq \mathbb{S}^n$ is d -Restricted SDC if there exists $\{\bar{A}_i\} \subseteq \mathbb{S}^{n+d}$ SDC

$$\bar{A}_i = \begin{pmatrix} A_i & * \\ * & * \end{pmatrix}$$

- $$x^\top A_i x = \begin{pmatrix} x \\ 0 \end{pmatrix}^\top \begin{pmatrix} A_i & * \\ * & * \end{pmatrix} \begin{pmatrix} x \\ 0 \end{pmatrix}$$
- $$\inf_{x,w} \begin{pmatrix} x \\ w \end{pmatrix}^\top \bar{A}_1 \begin{pmatrix} x \\ w \end{pmatrix} + \dots$$

s.t.
$$\begin{pmatrix} x \\ w \end{pmatrix}^\top \bar{A}_i \begin{pmatrix} x \\ w \end{pmatrix} + \dots = 0, \forall i = 2, \dots, m$$

$$w = 0$$

d -RSDC

- $\{A_i\} \subseteq \mathbb{S}^n$ is d -RSDC $\iff \exists \{\ell_1, \dots, \ell_{n+d}\} \subseteq \mathbb{R}^n$:
spanning \mathbb{R}^n
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$

d -RSDC

- $\{A_i\} \subseteq \mathbb{S}^n$ is d -RSDC $\iff \exists \{\ell_1, \dots, \ell_{n+d}\} \subseteq \mathbb{R}^n$:
spanning \mathbb{R}^n
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$
- $\{A, B\} \subseteq \mathbb{S}^n$ is naively n -RSDC

d -RSDC

- $\{A_i\} \subseteq \mathbb{S}^n$ is d -RSDC $\iff \exists \{\ell_1, \dots, \ell_{n+d}\} \subseteq \mathbb{R}^n$:
spanning \mathbb{R}^n
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$
- $\{A, B\} \subseteq \mathbb{S}^n$ is naively n -RSDC

Theorem ([W and Jiang 21])

Let $\{A, B\} \subseteq \mathbb{S}^n$. Suppose $A^{-1}B$ has only simple eigenvalues. Then $\{A, B\}$ is 1-RSDC.

d -RSDC

- $\{A_i\} \subseteq \mathbb{S}^n$ is d -RSDC $\iff \exists \{\ell_1, \dots, \ell_{n+d}\} \subseteq \mathbb{R}^n$:
spanning \mathbb{R}^n
$$A_i = \sum_j \mu_j^{(i)} \ell_j \ell_j^\top, \quad \forall i$$
- $\{A, B\} \subseteq \mathbb{S}^n$ is naively n -RSDC

Theorem ([W and Jiang 21])

Let $\{A, B\} \subseteq \mathbb{S}^n$. Suppose $A^{-1}B$ has only simple eigenvalues. Then $\{A, B\}$ is 1-RSDC.

- Tools: canonical form for pairs of symmetric matrices²

² [Uhlig 76], [Lancaster, Rodman 05]

Main idea



$$A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$$

Main idea

- $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$
- $\bar{A} = \left(\begin{array}{cc|c} & 1 & * \\ 1 & & * \\ \hline * & * & * \end{array} \right), \quad \bar{B} = \left(\begin{array}{cc|c} 1 & & * \\ & -1 & * \\ \hline * & * & * \end{array} \right)$

Main idea

- $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$
- $\bar{A} = \left(\begin{array}{cc|c} & 1 & * \\ 1 & & * \\ \hline * & * & * \end{array} \right), \quad \bar{B} = \left(\begin{array}{cc|c} 1 & & * \\ & -1 & * \\ \hline * & * & * \end{array} \right)$
- Want $\bar{A}^{-1}\bar{B}$ diagonalizable, real spectrum

Main idea

- $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$
- $\bar{A} = \left(\begin{array}{c|c} 1 & * \\ \hline 1 & * \\ * & * \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & * \\ \hline -1 & * \\ * & * \end{array} \right)$
- Want $\bar{A}^{-1}\bar{B}$ diagonalizable, real spectrum
- Suffices $\bar{A}^{-1}\bar{B}$ real simple eigenvalues

Main idea

- $A = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad B = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$
- $\bar{A} = \left(\begin{array}{c|c} 1 & * \\ \hline 1 & * \\ * & * \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & * \\ \hline -1 & * \\ * & * \end{array} \right)$
- Want $\bar{A}^{-1}\bar{B}$ diagonalizable, real spectrum
- Suffices $\bar{A}^{-1}\bar{B}$ **real simple eigenvalues**
- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & \\ \hline & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ \hline -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$

Main Idea



$$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta \\ \beta & \gamma \end{array} \right)$$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta \\ \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$


Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta \\ \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ \hline -1 & \beta \\ \hline \alpha & \beta \\ & \gamma \end{array} \right)$
 - $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$
- Basis for deg. 2 polynomials in z
-

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta \\ & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$ 

Basis for deg. 2 polynomials in z

Surjective function

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta \\ \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$
- Pick $\{\lambda_1, \lambda_2, \lambda_3\} \subseteq \mathbb{R}$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$

$$\gamma(\lambda_1^2 + 1) + (2\alpha\beta)\lambda_1 + (\beta^2 - \alpha^2)1 = \lambda_1(\lambda_1^2 + 1)$$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$
- $\det(\bar{A}^{-1}\bar{B} - zI)$
 $= \dots$
 $= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1)$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$

$$\gamma(2) + (2\alpha\beta)(-1) + (\beta^2 - \alpha^2)1 = -2$$

Main Idea

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ \hline -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$$
- $$\begin{aligned} \det(\bar{A}^{-1}\bar{B} - zI) \\ &= \dots \\ &= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1) \end{aligned}$$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$

$$\begin{matrix} & z^2+1 & z & 1 \\ \lambda_1 & \begin{pmatrix} 2 & -1 & 1 \end{pmatrix} \\ \lambda_2 & \begin{pmatrix} 1 & 0 & 1 \end{pmatrix} \\ \lambda_3 & \begin{pmatrix} 2 & 1 & 1 \end{pmatrix} \end{matrix} \begin{pmatrix} \gamma \\ 2\alpha\beta \\ \beta^2 - \alpha^2 \end{pmatrix} = \begin{matrix} & z(z^2+1) \\ \lambda_1 & \begin{pmatrix} -2 \\ 0 \\ 2 \end{pmatrix} \\ \lambda_2 & \\ \lambda_3 & \end{matrix}$$

Main Idea

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ \hline -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$$
- $$\begin{aligned} \det(\bar{A}^{-1}\bar{B} - zI) \\ &= \dots \\ &= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1) \end{aligned}$$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$

$$\begin{matrix} & z^2+1 & z & 1 \\ \lambda_1 & \begin{pmatrix} 2 & -1 & 1 \end{pmatrix} \\ \lambda_2 & \begin{pmatrix} 1 & 0 & 1 \end{pmatrix} \\ \lambda_3 & \begin{pmatrix} 2 & 1 & 1 \end{pmatrix} \end{matrix} \begin{pmatrix} \gamma \\ 2\alpha\beta \\ \beta^2 - \alpha^2 \end{pmatrix} = \begin{matrix} & z(z^2+1) \\ \lambda_1 & \begin{pmatrix} -2 \\ 0 \\ 2 \end{pmatrix} \\ \lambda_2 & \\ \lambda_3 & \end{matrix}$$

Main Idea

- $\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & \alpha \\ \hline -1 & \beta \\ \hline \alpha & \beta & \gamma \end{array} \right)$
- $$\begin{aligned} \det(\bar{A}^{-1}\bar{B} - zI) \\ &= \dots \\ &= \gamma(z^2 + 1) + (2\alpha\beta)z + (\beta^2 - \alpha^2)1 - z(z^2 + 1) \end{aligned}$$
- Pick $\{\lambda_1 = -1, \lambda_2 = 0, \lambda_3 = 1\} \subseteq \mathbb{R}$
$$\begin{array}{c} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{array} \begin{array}{c} z^2+1 \quad z \quad 1 \\ \left(\begin{array}{ccc} 2 & -1 & 1 \\ 1 & 0 & 1 \\ 2 & 1 & 1 \end{array} \right) \left(\begin{array}{c} \gamma \\ 2\alpha\beta \\ \beta^2 - \alpha^2 \end{array} \right) = \begin{array}{c} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{array} \begin{array}{c} z(z^2+1) \\ \left(\begin{array}{c} -2 \\ 0 \\ 2 \end{array} \right) \end{array}$$
- $\alpha = 1, \beta = 1, \gamma = 0$

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & 1 \\ -1 & 1 \\ \hline 1 & 1 \\ 0 & \end{array} \right)$$

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$
- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & 1 \\ \hline -1 & 1 \\ \hline 1 & 1 & 0 \end{array} \right)$$
- $\det(\bar{A}^{-1}\bar{B} - zI) = 0$ **at** $z = -1, 0, 1$

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & 1 \\ -1 & 1 \\ \hline 1 & 1 \\ 0 & \end{array} \right)$$

- $\det(\bar{A}^{-1}\bar{B} - zI) = 0$ at $z = -1, 0, 1$
- $\bar{A}^{-1}\bar{B}$ has real simple eigenvalues $\{-1, 0, 1\}$

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & 1 \\ -1 & 1 \\ \hline 1 & 1 \end{array} \middle| \begin{array}{c} 1 \\ 0 \end{array} \right)$$

- $\det(\bar{A}^{-1}\bar{B} - zI) = 0$ at $z = -1, 0, 1$
- $\bar{A}^{-1}\bar{B}$ has real simple eigenvalues $\{-1, 0, 1\}$
- $\{\bar{A}, \bar{B}\}$ is SDC

Main Idea: Recap

- $\alpha = 1, \beta = 1, \gamma = 0$

- $$\bar{A} = \left(\begin{array}{c|c} 1 & \\ \hline 1 & 1 \end{array} \right), \quad \bar{B} = \left(\begin{array}{c|c} 1 & 1 \\ \hline -1 & 1 \\ \hline 1 & 1 & 0 \end{array} \right)$$

- $\det(\bar{A}^{-1}\bar{B} - zI) = 0$ at $z = -1, 0, 1$
- $\bar{A}^{-1}\bar{B}$ has real simple eigenvalues $\{-1, 0, 1\}$
- $\{\bar{A}, \bar{B}\}$ is SDC
- Similar calculations generalize to (almost every) pair $\{A, B\} \subseteq \mathbb{S}^n$

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments
- 5 Conclusion: additional work, future directions

Setup



$$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$

Note: Slightly different setup than in paper

Setup

- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form

Note: Slightly different setup than in paper

Setup

- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form
- k is number of pairs of complex eigenvalues of $A_1^{-1}A_2$
“How far $\{A_1, A_2\}$ is from being SDC”

Note: Slightly different setup than in paper

Setup

- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form
- k is number of pairs of complex eigenvalues of $A_1^{-1}A_2$
“How far $\{A_1, A_2\}$ is from being SDC”
- Tested: As-is

Note: Slightly different setup than in paper

Setup

- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form
- k is number of pairs of complex eigenvalues of $A_1^{-1}A_2$
“How far $\{A_1, A_2\}$ is from being SDC”
- Tested: As-is, 1-RSDC

Note: Slightly different setup than in paper

Setup

- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form
- k is number of pairs of complex eigenvalues of $A_1^{-1}A_2$
“How far $\{A_1, A_2\}$ is from being SDC”
- Tested: As-is, 1-RSDC, 2-RSDC


Note: Slightly different setup than in paper

Setup


- $$\begin{aligned} \inf_{x \in \mathbb{R}^n} \quad & x^\top A_1 x \\ \text{s.t.} \quad & x^\top A_2 x \leq 0 \\ & Lx \leq 1 \end{aligned}$$
- A_1, A_2 generated randomly in canonical form
- k is number of pairs of complex eigenvalues of $A_1^{-1}A_2$
“How far $\{A_1, A_2\}$ is from being SDC”
- Tested: As-is, 1-RSDC, 2-RSDC, n -RSDC

Note: Slightly different setup than in paper

Results for $n = 15$




Results for $n = 15$




- *-RSDC outperforms As-is on every instance

Results for $n = 15$



- *-RSDC outperforms As-is on every instance
- Condition number blows up with k

Results for $n = 15$



- *-RSDC outperforms As-is on every instance
- Condition number blows up with k
 - $k = 3$: 1-RSDC ($\sim 10^3$), 2-RSDC ($\sim 10^2$), n -RSDC (1)

Outline

- 1 Introduction: QCQPs and diagonalization
- 2 Prior work: SDC, first examples
 - When is $\{A, B\}$ SDC?
- 3 New notions of simultaneous diagonalizability
 - d -Restricted SDC
 - When is $\{A, B\}$ 1-RSDC? Almost everywhere!
- 4 Experiments
- 5 Conclusion: additional work, future directions

Summary, future directions

- New notions: d -RSDC

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices
- Future directions:

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices
- Future directions: Need better definitions of SDC

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices
- Future directions: Need better definitions of SDC
 - What if $\text{cond}(P)$ must be bounded?






Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices
- Future directions: Need better definitions of SDC
 - What if $\text{cond}(P)$ must be bounded?
 - Parameterized constructions of d -RSDC?

Summary, future directions

- New notions: d -RSDC
- 1-RSDC holds a.e. for pairs
- Additional results:
 - Almost SDC (ASDC)
 - Complete characterization of ASDC for pairs
 - Necessary and/or sufficient conditions for ASDC and d -RSDC for small numbers of matrices
- Future directions: Need better definitions of SDC
 - What if $\text{cond}(P)$ must be bounded?
 - Parameterized constructions of d -RSDC?
- Thank you. Questions?

References I

-  [A. Ben-Tal and D. den Hertog.](#) “Hidden conic quadratic representation of some nonconvex quadratic optimization problems”. In: *Math. Program.* 143 (2014), pp. 1–29.
-  [S. Burer and Y. Ye.](#) “Exact semidefinite formulations for a class of (random and non-random) nonconvex quadratic programs”. In: *Math. Program.* 181 (2019), pp. 1–17.
-  [R. A. Horn and C. R. Johnson.](#) *Matrix analysis*. Cambridge University Press, 2012.
-  [R. Jiang and D. Li.](#) “Simultaneous diagonalization of matrices and its applications in quadratically constrained quadratic programming”. In: *SIAM J. Optim.* 26.3 (2016), pp. 1649–1668.
-  [P. Lancaster and L. Rodman.](#) “Canonical forms for Hermitian matrix pairs under strict equivalence and congruence”. In: *SIAM Review* 47.3 (2005), pp. 407–443.

References II



T. H. Le and T. N. Nguyen. “Simultaneous diagonalization via congruence of Hermitian matrices: some equivalent conditions and a numerical solution”. In: *arXiv preprint arXiv:2007.14034* (2020).



T. Nguyen et al. “On simultaneous diagonalization via congruence of real symmetric matrices”. In: *arXiv preprint arXiv:2004.06360* (2020).



K. O’meara and C. Vinsonhaler. “On approximately simultaneously diagonalizable matrices”. In: *Linear Algebra Appl.* 412 (2006), pp. 39–74.



F. Uhlig. “A canonical form for a pair of real symmetric matrices that generate a nonsingular pencil”. In: *Linear Algebra Appl.* 14.3 (1976), pp. 189–209.

References III



A. L. Wang and R. Jiang. “New notions of simultaneous diagonalizability of quadratic forms with applications to QCQPs”. In: *arXiv preprint 2101.12141* (2021).



A. L. Wang and F. Kılınç-Karzan. “On the tightness of SDP relaxations of QCQPs”. In: *Math. Program.* (2021). Forthcoming. DOI: 10.1007/s10107-020-01589-9.

Additional results

Definition

$\{A_i\}$ is almost SDC (ASDC) if for all $\epsilon > 0$, there exists $\{A'_i\}$ SDC, $\max_i \|A_i - A'_i\| \leq \epsilon$

Theorem ([W and Jiang 21])

Let $\{A, B\} \subseteq \mathbb{S}^n$

- If A invertible, then

$$\{A, B\} \text{ ASDC} \iff A^{-1}B \text{ has real spectrum}$$

- If $\text{span}(\{A, B\})$ does not contain invertible matrix, then

$$\{A, B\} \text{ ASDC}$$

Related: [O'meara, Vinsonhaler 06]

Additional results

Theorem

$A \in \mathbb{S}^n$ invertible. Then,

$$\{A, B, C\} \text{ ASDC} \iff \{A^{-1}B, A^{-1}C\} \text{ commute,} \\ \text{real spectrum}$$

Theorem

$\{A = I_n, B, C\} \subseteq \mathbb{S}^n$. If $d \leq \text{rank}([B, C])/2$, then

- $\{A, B, C\}$ is not d -RSDC
- $\left\{ \begin{pmatrix} A & \\ & 0_d \end{pmatrix}, \begin{pmatrix} B & \\ & 0_d \end{pmatrix}, \begin{pmatrix} C & \\ & 0_d \end{pmatrix} \right\}$ is not ASDC

Additional results

Theorem

There exists $\{A_1, \dots, A_7\} \subseteq \mathbb{S}^6$ such that

- A_1 invertible,
- $\{A_1^{-1}A_2, \dots, A_1^{-1}A_7\}$ commute, real spectrum,
- not ASDC

Theorem

There exists $\{A_1, \dots, A_5\} \subseteq \mathbb{H}^4$ such that

- A_1 invertible,
- $\{A_1^{-1}A_2, \dots, A_1^{-1}A_5\}$ commute, real spectrum,
- not ASDC

[W and Jiang 21]