


# Exactness in SDP relaxations of QCQPs

\*now with 50% more pictures!

Alex L. Wang , PSE Seminar, Oct. 20



Supported in part by NSF grant CMMI 1454548 and ONR grant N00014-19-1-2321

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms
- 4 Some results
- 5 Application: robust least squares
- 6 Conclusion

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms
- 4 Some results
- 5 Application: robust least squares
- 6 Conclusion

# Quadratically constrained quadratic programs (QCQPs)

- Family of highly expressive optimization problems

# Quadratically constrained quadratic programs (QCQPs)

- Family of highly expressive optimization problems
- Computer science  
MAX-CUT, MAX-CLIQUE

# Quadratically constrained quadratic programs (QCQPs)

- Family of highly expressive optimization problems
- Computer science  
MAX-CUT, MAX-CLIQUE
- Operations research  
Facility location, production planning

# Quadratically constrained quadratic programs (QCQPs)

- Family of highly expressive optimization problems
- Computer science  
MAX-CUT, MAX-CLIQUE
- Operations research  
Facility location, production planning
- Engineering  
Pooling problem, truss design problem

# Quadratically constrained quadratic programs (QCQPs)

- Family of highly expressive optimization problems
- Computer science  
MAX-CUT, MAX-CLIQUE
- Operations research  
Facility location, production planning
- Engineering  
Pooling problem, truss design problem
- More generally  
Binary programming, polynomial programming



# From linear programs to quadratic programs

- Linear programs (LPs)

# From linear programs to quadratic programs

- Linear programs (LPs)
- $\ell_0, \ell_1, \dots, \ell_m : \mathbb{R}^n \rightarrow \mathbb{R}$  linear functions
$$\ell_i(x) = b_i^\top x + c_i$$

# From linear programs to quadratic programs

- Linear programs (LPs)
- $\ell_0, \ell_1, \dots, \ell_m : \mathbb{R}^n \rightarrow \mathbb{R}$  linear functions  
 $\ell_i(x) = b_i^\top x + c_i$

- Want to find


$$\text{Opt} := \min_{x \in \mathbb{R}^n} \left\{ \ell_0(x) : \begin{array}{l} \ell_1(x) \leq 0 \\ \vdots \\ \ell_m(x) \leq 0 \end{array} \right\}$$

# From linear programs to quadratic programs

- Linear programs (LPs)
- $\ell_0, \ell_1, \dots, \ell_m : \mathbb{R}^n \rightarrow \mathbb{R}$  linear functions  
 $\ell_i(x) = b_i^\top x + c_i$

- Want to find

$$\text{Opt} := \min_{x \in \mathbb{R}^n} \left\{ \ell_0(x) : \begin{array}{l} \ell_1(x) \leq 0 \\ \vdots \\ \ell_m(x) \leq 0 \end{array} \right\}$$



# From linear programs to quadratic programs

- Quadratically constrained quadratic programs (QCQPs)

# From linear programs to quadratic programs

- Quadratically constrained quadratic programs (QCQPs)
- $q_0, q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$  quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

# From linear programs to quadratic programs

- Quadratically constrained quadratic programs (QCQPs)
- $q_0, q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$  quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- Want to find

$$\text{Opt} := \min_{x \in \mathbb{R}^n} \left\{ q_0(x) : \begin{array}{l} q_1(x) \leq 0 \\ \vdots \\ q_m(x) \leq 0 \end{array} \right\}$$


# From linear programs to quadratic programs

- Quadratically constrained quadratic programs (QCQPs)
- $q_0, q_1, \dots, q_m : \mathbb{R}^n \rightarrow \mathbb{R}$  quadratic functions

$$q_i(x) = x^\top A_i x + 2b_i^\top x + c_i$$

- Want to find


$$\text{Opt} := \min_{x \in \mathbb{R}^n} \left\{ q_0(x) : \begin{array}{l} q_1(x) \leq 0 \\ \vdots \\ q_m(x) \leq 0 \end{array} \right\}$$






# The QCQP Epigraph

$$\text{Opt} = \min_{x \in \mathbb{R}^n} \{q_0(x) : q_i(x) \leq 0, \forall i \in [m]\}$$





# The QCQP Epigraph

$$\begin{aligned}\text{Opt} &= \min_{x \in \mathbb{R}^n} \{q_0(x) : q_i(x) \leq 0, \forall i \in [m]\} \\ &= \min_{x, t} \left\{ t : \begin{array}{l} q_0(x) \leq t \\ q_i(x) \leq 0, \forall i \in [m] \end{array} \right\}\end{aligned}$$






# The QCQP Epigraph

$$\begin{aligned}\text{Opt} &= \min_{x \in \mathbb{R}^n} \{q_0(x) : q_i(x) \leq 0, \forall i \in [m]\} \\ &= \min_{x,t} \left\{ t : \begin{array}{l} q_0(x) \leq t \\ q_i(x) \leq 0, \forall i \in [m] \end{array} \right\} =: \min_{x,t} \left\{ t : (x,t) \in \mathcal{E} \right\}\end{aligned}$$



# The QCQP Epigraph

$$\begin{aligned}\text{Opt} &= \min_{x \in \mathbb{R}^n} \{q_0(x) : q_i(x) \leq 0, \forall i \in [m]\} \\ &= \min_{x,t} \left\{ t : \begin{array}{l} q_0(x) \leq t \\ q_i(x) \leq 0, \forall i \in [m] \end{array} \right\} =: \min_{x,t} \left\{ t : (x,t) \in \mathcal{E} \right\} \\ &= \min_{x,t} \{t : (x,t) \in \text{conv}(\mathcal{E})\}\end{aligned}$$



# Convex relaxations

- QCQPs are NP-hard in general 😞

# Convex relaxations

- QCQPs are NP-hard in general 😞
- One issue with QCQPs is nonconvexity!

# Convex relaxations

- QCQPs are NP-hard in general ☹️
- One issue with QCQPs is nonconvexity!
- Will look for a convex *relaxation*



# Convex relaxations

- QCQPs are NP-hard in general ☹️
- One issue with QCQPs is nonconvexity!
- Will look for a convex *relaxation*




- SDP relaxation can be solved efficiently




# Convex relaxations

- Properties you might want for a convex relaxation





# Convex relaxations

- Properties you might want for a convex relaxation
- Types of “exactness”




# Convex relaxations

- Properties you might want for a convex relaxation
- Types of “exactness”





Objective value




Convex hull

# Convex relaxations

- Properties you might want for a convex relaxation
- Types of “exactness”



Objective value



Convex hull

- **Q:** When do these properties hold?

# The standard SDP relaxation of QCQP

Standard semidefinite program (SDP) relaxation

$$\begin{aligned}\text{Opt} &= \min_x \left\{ q_0(x) : q_i(x) \leq 0, \forall i \right\} \\ &= \min_x \left\{ x^\top A_0 x + 2b_0^\top x + c_0 : x^\top A_i x + 2b_i^\top x + c_i \leq 0, \forall i \right\}\end{aligned}$$

# The standard SDP relaxation of QCQP

Standard semidefinite program (SDP) relaxation

$$\begin{aligned}\text{Opt} &= \min_x \left\{ q_0(x) : q_i(x) \leq 0, \forall i \right\} \\ &= \min_x \left\{ x^\top A_0 x + 2b_0^\top x + c_0 : x^\top A_i x + 2b_i^\top x + c_i \leq 0, \forall i \right\} \\ &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} Y = xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}\end{aligned}$$

# The standard SDP relaxation of QCQP

Standard semidefinite program (SDP) relaxation

$$\begin{aligned}\text{Opt} &= \min_x \left\{ q_0(x) : q_i(x) \leq 0, \forall i \right\} \\ &= \min_x \left\{ x^\top A_0 x + 2b_0^\top x + c_0 : x^\top A_i x + 2b_i^\top x + c_i \leq 0, \forall i \right\} \\ &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} Y = xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &\geq \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}\end{aligned}$$

# The standard SDP relaxation of QCQP

Standard semidefinite program (SDP) relaxation

$$\begin{aligned}\text{Opt} &= \min_x \left\{ q_0(x) : q_i(x) \leq 0, \forall i \right\} \\ &= \min_x \left\{ x^\top A_0 x + 2b_0^\top x + c_0 : x^\top A_i x + 2b_i^\top x + c_i \leq 0, \forall i \right\} \\ &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} Y = xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &\geq \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &=: \text{Opt}_{\text{SDP}}\end{aligned}$$



# SDP epigraph

$$\text{Opt}_{\text{SDP}} = \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}$$

# SDP epigraph

$$\begin{aligned}\text{Opt}_{\text{SDP}} &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &= \min_{x, t} \left\{ t : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_0, Y \rangle + 2b_0^\top x + c_0 \leq t \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}\end{aligned}$$

# SDP epigraph


$$\begin{aligned}\text{Opt}_{\text{SDP}} &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &= \min_{x, t} \left\{ t : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_0, Y \rangle + 2b_0^\top x + c_0 \leq t \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}\end{aligned}$$

- Let  $\mathcal{E}_{\text{SDP}}$  denote the SDP epigraph

# SDP epigraph

$$\begin{aligned}\text{Opt}_{\text{SDP}} &= \min_x \left\{ \langle A_0, Y \rangle + 2b_0^\top x + c_0 : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\} \\ &= \min_{x, t} \left\{ t : \begin{array}{l} \exists Y \succeq xx^\top \\ \langle A_0, Y \rangle + 2b_0^\top x + c_0 \leq t \\ \langle A_i, Y \rangle + 2b_i^\top x + c_i \leq 0, \forall i \end{array} \right\}\end{aligned}$$

- Let  $\mathcal{E}_{\text{SDP}}$  denote the SDP epigraph



# Recap

- QCQPs are highly expressive but NP-hard in general

# Recap


- QCQPs are highly expressive but NP-hard in general
- Solve SDP relaxation instead

# Recap

- QCQPs are highly expressive but NP-hard in general
- Solve SDP relaxation instead
- **Q:** What are sufficient conditions for

# Recap


- QCQPs are highly expressive but NP-hard in general
- Solve SDP relaxation instead
- **Q:** What are sufficient conditions for
  - Convex hull exactness:  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$ ?





# Recap

- QCQPs are highly expressive but NP-hard in general
- Solve SDP relaxation instead
- **Q:** What are sufficient conditions for
  - Convex hull exactness:  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$ ?



- Objective value exactness:  $\text{Opt} = \text{Opt}_{\text{SDP}}$ ?

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers
- The quadratic eigenvalue multiplicity

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers
  - The quadratic eigenvalue multiplicity
- Example application: robust least squares

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers
  - The quadratic eigenvalue multiplicity
- Example application: robust least squares
- Additional work in this area

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers
  - The quadratic eigenvalue multiplicity
- Example application: robust least squares
- Additional work in this area
- Future directions

---

[W and Kılınç-Karzan 19]

# Outline

- Main result for today:

If the quadratic forms “interact nicely” and each have “large amounts of symmetry”, then convex hull exactness holds

- The set of convex Lagrange multipliers
  - The quadratic eigenvalue multiplicity
- Example application: robust least squares
- Additional work in this area
- Future directions
- Sneak peek of (very) recent work

---

[W and Kılınç-Karzan 19]



- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers**
- 3 Symmetries in quadratic forms
- 4 Some results
- 5 Application: robust least squares
- 6 Conclusion

# Aggregation

- For  $\gamma \in \mathbb{R}_+^m$ , define



$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

# Aggregation

- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

- For all  $\gamma \in \mathbb{R}_+^m$ ,  $(x, t) \in \mathcal{E} \implies q_\gamma(x) \leq t$






# Aggregation

- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

- For all  $\gamma \in \mathbb{R}_+^m$ ,  $(x, t) \in \mathcal{E} \implies q_\gamma(x) \leq t$



# Aggregation

- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$


# Aggregation

- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

- Define

$$\Gamma := \{ \gamma \in \mathbb{R}_+^m : q_\gamma(x) \text{ is convex} \}$$



# Aggregation



- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

- Define

$$\Gamma := \{\gamma \in \mathbb{R}_+^m : q_\gamma(x) \text{ is convex}\}$$

- For all  $\gamma \in \Gamma$ ,  $(x, t) \in \text{conv}(\mathcal{E}) \implies q_\gamma(x) \leq t$



# Aggregation


- For  $\gamma \in \mathbb{R}_+^m$ , define

$$q_\gamma(x) := q_0(x) + \sum_{i=1}^m \gamma_i q_i(x)$$

- Define

$$\Gamma := \{ \gamma \in \mathbb{R}_+^m : q_\gamma(x) \text{ is convex} \}$$

- For all  $\gamma \in \Gamma$ ,  $(x, t) \in \text{conv}(\mathcal{E}) \implies q_\gamma(x) \leq t$







# Rewriting the SDP in terms of $\Gamma$

## Lemma

Suppose primal feasibility and dual strict feasibility, then

$$\mathcal{E}_{\text{SDP}} = \left\{ (x, t) : \max_{\gamma \in \Gamma} q_{\gamma}(x) \leq t \right\}$$



---


Related: [Fujie and Kojima 97]

# Rewriting the SDP in terms of $\Gamma$

## Lemma

Suppose primal feasibility and dual strict feasibility, then

$$\mathcal{E}_{\text{SDP}} = \left\{ (x, t) : \max_{\gamma \in \Gamma} q_{\gamma}(x) \leq t \right\}$$




Related: [Fujie and Kojima 97]

# Rewriting the SDP in terms of $\Gamma$

## Lemma

Suppose primal feasibility and dual strict feasibility, then

$$\mathcal{E}_{\text{SDP}} = \left\{ (x, t) : \max_{\gamma \in \Gamma} q_{\gamma}(x) \leq t \right\}$$




Related: [Fujie and Kojima 97]

# Rewriting the SDP in terms of $\Gamma$

## Lemma

Suppose primal feasibility and dual strict feasibility, then

$$\mathcal{E}_{\text{SDP}} = \left\{ (x, t) : \max_{\gamma \in \Gamma} q_{\gamma}(x) \leq t \right\}$$




Related: [Fujie and Kojima 97]

# Rewriting the SDP in terms of $\Gamma$

## Lemma

Suppose primal feasibility and dual strict feasibility, then

$$\mathcal{E}_{\text{SDP}} = \left\{ (x, t) : \max_{\gamma \in \Gamma} q_{\gamma}(x) \leq t \right\}$$



Related: [Fujie and Kojima 97]

# What does $\Gamma$ look like?


- Recall

$$\Gamma = \left\{ \gamma \in \mathbb{R}_+^m : A_0 + \sum_{i=1}^m \gamma_i A_i \succeq 0 \right\}$$

# What does $\Gamma$ look like?

- Recall



$$\Gamma = \left\{ \gamma \in \mathbb{R}_+^m : A_0 + \sum_{i=1}^m \gamma_i A_i \succeq 0 \right\}$$



# What does $\Gamma$ look like?

- Recall

$$\Gamma = \left\{ \gamma \in \mathbb{R}_+^m : A_0 + \sum_{i=1}^m \gamma_i A_i \succeq 0 \right\}$$







# What does $\Gamma$ look like?

- Recall

$$\Gamma = \left\{ \gamma \in \mathbb{R}_+^m : A_0 + \sum_{i=1}^m \gamma_i A_i \succeq 0 \right\}$$



- When  $A_i$ s diagonal  $\implies \Gamma$  is polyhedral

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms**
- 4 Some results
- 5 Application: robust least squares
- 6 Conclusion

# Quadratic eigenvalue multiplicity

## Definition

Let  $1 \leq k \leq n$  be the largest integer such that for each  $i = 0, \dots, m$ , the matrix  $A_i \in \mathbb{S}^n$  has the following block form

$$A_i = \hat{A}_i \otimes I_k = \begin{pmatrix} \hat{A}_i & & & \\ & \hat{A}_i & & \\ & & \ddots & \\ & & & \hat{A}_i \end{pmatrix}$$

where  $\hat{A}_i \in \mathbb{S}^{n/k}$

# Quadratic eigenvalue multiplicity

- $A_i = \hat{A}_i \otimes I_k$

# Quadratic eigenvalue multiplicity

- $A_i = \hat{A}_i \otimes I_k$
- Suppose  $n = 4$ , i.e.  $x \in \mathbb{R}^4$

---

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 \quad \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \quad k = 4$$

# Quadratic eigenvalue multiplicity

- $A_i = \hat{A}_i \otimes I_k$
- Suppose  $n = 4$ , i.e.  $x \in \mathbb{R}^4$

---

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 \quad \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \quad k = 4$$

$$(x_1 - x_2)^2 + (x_3 - x_4)^2 \quad \begin{pmatrix} 1 & -1 & & \\ -1 & 1 & & \\ & & 1 & -1 \\ & & -1 & 1 \end{pmatrix} \quad k = 2$$

---

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms
- 4 Some results**
- 5 Application: robust least squares
- 6 Conclusion

# Some results

## Corollary

Suppose primal feasibility and dual strict feasibility. If  $\Gamma$  is polyhedral and

$$k \geq \min(m, |\{b_i \neq 0\}_{i=1}^m| + 1),$$

then  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$  and  $\text{Opt} = \text{Opt}_{\text{SDP}}$ .

---

Related: [Yakubovich 79], [Burer and Ye 19]



# Some results

## Corollary

Suppose primal feasibility and dual strict feasibility. If  $\Gamma$  is polyhedral and

$$k \geq \min(m, |\{b_i \neq 0\}_{i=1}^m| + 1),$$

then  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$  and  $\text{Opt} = \text{Opt}_{\text{SDP}}$ .

- $m = 1$

---

Related: [Yakubovich 79], [Burer and Ye 19]

# Some results

## Corollary

Suppose primal feasibility and dual strict feasibility. If  $\Gamma$  is polyhedral and

$$k \geq \min(m, |\{b_i \neq 0\}_{i=1}^m| + 1),$$

then  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$  and  $\text{Opt} = \text{Opt}_{\text{SDP}}$ .

- $m = 1$
- $A_i$ s diagonal and  $b_1 = b_2 = \dots = b_m = 0$

---

Related: [Yakubovich 79], [Burer and Ye 19]

# Some results

## Corollary

Suppose primal feasibility and dual strict feasibility. If  $\Gamma$  is polyhedral and

$$k \geq \min(m, |\{b_i \neq 0\}_{i=1}^m| + 1),$$

then  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$  and  $\text{Opt} = \text{Opt}_{\text{SDP}}$ .

- $m = 1$
- $A_i$ s diagonal and  $b_1 = b_2 = \dots = b_m = 0$
- $A_i = \alpha_i I_n$  for all  $i$  and  $n \geq m$


---

Related: [Yakubovich 79], [Burer and Ye 19]

# Example: Swiss cheese

- Minimizing distance to a piece of Swiss cheese


$$\min_{x \in \mathbb{R}^n} \left\{ \|x\|^2 : \begin{array}{l} \text{inside ball constraints} \\ \text{outside ball constraints} \\ \text{linear constraints} \end{array} \right\}$$



# Example: Swiss cheese

- Minimizing distance to a piece of Swiss cheese

$$\min_{x \in \mathbb{R}^n} \left\{ \|x\|^2 : \begin{array}{l} \text{inside ball constraints} \\ \text{outside ball constraints} \\ \text{linear constraints} \end{array} \right\}$$




- inside ball  $\mapsto I$ , outside ball  $\mapsto -I$ , linear constraints  $\mapsto 0$

# Example: Swiss cheese

- Minimizing distance to a piece of Swiss cheese

$$\min_{x \in \mathbb{R}^n} \left\{ \|x\|^2 : \begin{array}{l} \text{inside ball constraints} \\ \text{outside ball constraints} \\ \text{linear constraints} \end{array} \right\}$$



- inside ball  $\mapsto I$ , outside ball  $\mapsto -I$ , linear constraints  $\mapsto 0$
- If nonempty and  $n \geq m$ , then the standard SDP relaxation is tight for this QCQP

# Some results

## Corollary

Suppose primal feasibility and dual strict feasibility.

- If  $k \geq m + 2$ , then  $\text{conv}(\mathcal{E}) = \mathcal{E}_{\text{SDP}}$
- If  $k \geq m + 1$ , then  $\text{Opt} = \text{Opt}_{\text{SDP}}$


---

Related: [Beck 07]

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms
- 4 Some results
- 5 Application: robust least squares**
- 6 Conclusion




# Least squares



- Input, output pairs

$$Y^* = \begin{pmatrix} (y_1^*)^\top \\ \vdots \\ (y_n^*)^\top \end{pmatrix} \in \mathbb{R}^{n \times k}, \quad z = \begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix} \in \mathbb{R}^n$$

# Least squares




- Input, output pairs

$$Y^* = \begin{pmatrix} (y_1^*)^\top \\ \vdots \\ (y_n^*)^\top \end{pmatrix} \in \mathbb{R}^{n \times k}, \quad z = \begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix} \in \mathbb{R}^n$$

- Want  $Y^* \ell \approx z$

$$\min_{\ell \in \mathbb{R}^k} \|Y^* \ell - z\|_2^2$$


# Robust least squares



- Empirical measurement  $\hat{Y}$  and uncertainty  $\mathcal{U}$ , i.e.,

$$Y^* \in \hat{Y} + \mathcal{U}$$


# Robust least squares



- Empirical measurement  $\hat{Y}$  and uncertainty  $\mathcal{U}$ , i.e.,  
 $Y^* \in \hat{Y} + \mathcal{U}$
- Want to minimize

$$\min_{\ell \in \mathbb{R}^k} \max_{X \in \mathcal{U}} \left\| (\hat{Y} + X)\ell - z \right\|_2^2$$

# Robust least squares



- Empirical measurement  $\hat{Y}$  and uncertainty  $\mathcal{U}$ , i.e.,  
 $Y^* \in \hat{Y} + \mathcal{U}$
- Want to minimize

$$\min_{\ell \in \mathbb{R}^k} \max_{X \in \mathcal{U}} \left\| (\hat{Y} + X)\ell - z \right\|_2^2$$

- When  $\mathcal{U}$  is defined by quadratics, can apply our theory!

# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

- Consider

$$\max_{X \in \mathbb{R}^{n \times k}} \left\{ \left\| (\hat{Y} + X)\ell - z \right\|_2^2 : X \in \mathcal{U} \right\}$$

# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

- Consider

$$\max_{X \in \mathbb{R}^{n \times k}} \left\{ \left\| (\hat{Y} + X)\ell - z \right\|_2^2 : X \in \mathcal{U} \right\}$$

- Write as a QCQP in the variable  $x \in \mathbb{R}^{nk}$



# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

- Consider

$$\max_{X \in \mathbb{R}^{n \times k}} \left\{ \left\| (\hat{Y} + X)\ell - z \right\|_2^2 : X \in \mathcal{U} \right\}$$

- Write as a QCQP in the variable  $x \in \mathbb{R}^{nk}$
- Quadratic forms

$$L_i^\top L_i \otimes I_k$$

# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

- Consider

$$\max_{X \in \mathbb{R}^{n \times k}} \left\{ \left\| (\hat{Y} + X)\ell - z \right\|_2^2 : X \in \mathcal{U} \right\}$$

- Write as a QCQP in the variable  $x \in \mathbb{R}^{nk}$
- Quadratic forms

$$L_i^\top L_i \otimes I_k$$

- $k \geq m + 2$  implies convex hull exactness

# Robust Least Squares

- Suppose

$$\mathcal{U} = \left\{ X \in \mathbb{R}^{n \times k} : \|L_i X\|_F^2 \leq c_i, \forall i \in [m] \right\}$$

- Consider

$$\max_{X \in \mathbb{R}^{n \times k}} \left\{ \left\| (\hat{Y} + X)\ell - z \right\|_2^2 : X \in \mathcal{U} \right\}$$

- Write as a QCQP in the variable  $x \in \mathbb{R}^{nk}$
- Quadratic forms

$$L_i^\top L_i \otimes I_k$$

- $k \geq m + 2$  implies convex hull exactness
- $k \geq m + 1$  implies objective value exactness

- 1 Introduction: QCQPs and SDPs
- 2 SDP relaxations and convex Lagrange multipliers
- 3 Symmetries in quadratic forms
- 4 Some results
- 5 Application: robust least squares
- 6 Conclusion**

# Recap, additional work, future work

- QCQPs are NP-hard

---

Related: [Argue, Kılınç-Karzan, and [W](#) 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation

---

Related: [Argue, Kılınç-Karzan, and [W](#) 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,

---

Related: [Argue, Kılınç-Karzan, and [W](#) 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$

---

Related: [Argue, Kılınç-Karzan, and W 20]



# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation exact?

| Polyhedral $\Gamma$ | General $\Gamma$ |
|---------------------|------------------|
| Cvx. hull           |                  |
| Obj. val.           |                  |

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ |                  |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ |                  |

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 2$   |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 1$   |

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 2$   |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 1$   |

- Current and future work:

---

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 2$   |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 1$   |

- Current and future work:
  - Is the polyhedral case fundamentally different from the general case?

---

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 2$   |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 1$   |

- Current and future work:
  - Is the polyhedral case fundamentally different from the general case?
  - Conditions that only depend on the constraints?

Related: [Argue, Kılınç-Karzan, and W 20]

# Recap, additional work, future work

- QCQPs are NP-hard
- Analyzed SDP relaxation
  - Convex Lagrange multipliers,
  - “Amount of symmetry”  $k$
- **Q:** When is the SDP relaxation **exact**?

|           | Polyhedral $\Gamma$                            | General $\Gamma$ |
|-----------|--|------------------|
| Cvx. hull | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 2$   |
| Obj. val. | $k \geq \min(m,  \{b_i \neq 0\}_{i=1}^m  + 1)$ | $k \geq m + 1$   |

- Current and future work:
  - Is the polyhedral case fundamentally different from the general case?
  - Conditions that only depend on the constraints?
  - Can **approximation results** be explained in this framework?

Related: [Argue, Kılınç-Karzan, and W 20]

Thank you. Questions?

Slides

[cs.cmu.edu/~alw1](https://cs.cmu.edu/~alw1)


Full version

[A. L. Wang and F. Kılınç-Karzan](#). “On the tightness of SDP relaxations of QCQPs”. In: *arXiv preprint arXiv:1911.09195* (2019)




# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$




# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$
- Rounding scheme: Given  $x \in \overline{\mathcal{S}}$ ,




# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$
- Rounding scheme: Given  $x \in \overline{\mathcal{S}}$ ,
  - Either output convex decomposition for  $x \in \text{conv}(\mathcal{S})$




# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$
- Rounding scheme: Given  $x \in \overline{\mathcal{S}}$ ,
  - Either output convex decomposition for  $x \in \text{conv}(\mathcal{S})$
  - Or, claim  $x' \in \overline{\mathcal{S}} \setminus \text{conv}(\mathcal{S})$ .




# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$
- Rounding scheme: Given  $x \in \overline{\mathcal{S}}$ ,
  - Either output convex decomposition for  $x \in \text{conv}(\mathcal{S})$
  - Or, claim  $x' \in \overline{\mathcal{S}} \setminus \text{conv}(\mathcal{S})$ .
- Correct when  $\Gamma^\circ$  is facially exposed








# When $\Gamma$ is non-polyhedral

- The geometry of  $\Gamma$  and  $\Gamma^\circ$
- Rounding scheme: Given  $x \in \overline{\mathcal{S}}$ ,
  - Either output convex decomposition for  $x \in \text{conv}(\mathcal{S})$
  - Or, claim  $x' \in \overline{\mathcal{S}} \setminus \text{conv}(\mathcal{S})$ .
- Correct when  $\Gamma^\circ$  is facially exposed
- Extends framework to handle complementarity constraints  
→ sparse regression



# References I

-  C.J. Argue, F. Kılınç-Karzan, and A. L. Wang. “Necessary and sufficient conditions for rank-one generated cones”. In: *arXiv preprint arXiv:2007.07433* (2020).
-  A. Beck. “Quadratic matrix programming”. In: *SIAM J. Optim.* 17.4 (2007), pp. 1224–1238.
-  S. Burer and Y. Ye. “Exact semidefinite formulations for a class of (random and non-random) nonconvex quadratic programs”. In: *Math. Program.* 181 (2019), pp. 1–17.
-  T. Fujie and M. Kojima. “Semidefinite programming relaxation for nonconvex quadratic programs”. In: *J. Global Optim.* 10.4 (1997), pp. 367–380.
-  V. A. Yakubovich. “S-procedure in nonlinear control theory”. In: *Vestnik Leningrad Univ. Math.* (1971), pp. 73–93.