Asian option call price

Table of Contents

Parameters	1
Integral approximation	
Output results	2
Function	2

Consider the following integral, which arises in the pricing of an Asian arithmetic mean call option

```
Error updating Text.
```

```
String scalar or character vector must have valid interpreter syntax: 
 \pi_{\alpha} = \pi_{\alpha} \cdot \pi_{\alpha
```

Notice the exponent part $(e^{g(z)})$ coresponds to the probability distribution function of a normally distributed random vector with covariance matrix Σ . That's the main reason we will generate IID standard normal random variables. We omit the mathematical justification here.

Parameters

Integral approximation

Monte Carlo methods can be used to estimate expectations, the mean value of a random variable's function. For continuous variables, expectations take the form of integrals.

Therefore, if we can express an integral as an expectation, it is possible to estimate its value using Monte Carlo methods.

```
A = chol(Sigma); % Sigma = A'*A

s_1 = @(z_1)100*exp(-0.0225 + 0.3*z_1);
s_2 = @(z_2)100*exp(-0.045 + 0.3*z_2);
z = @(n)randn(n, 2)*A;

g = @(n)gsub(n, z, s_1, s_2); % encapsulate inner function

tic;
[muhat, out] = meanMC_g(g, abstol, reltol);
toc;
```

Output results

```
disp(['The estimated fair price is ' num2str(muhat)])
The estimated fair price is 9.445
```

Function

```
function [price] = gsub(n, z, s_1, s_2)
%GSUB Pricing of an Asian arithmetic mean call option
zn = z(n);
s1 = s_1(zn(:, 1));
s2 = s_2(zn(:, 2));
asianCallOpt = .5*(s1 + s2) - 100;
price = max([asianCallOpt zeros(n, 1)], [], 2);
end

Elapsed time is 0.690454 seconds.

Author: Alejandro Madriñán Fernández
```

Published with MATLAB® R2021a