

COMP27112


Computer
Graphics
and
Image Processing

7: Rendering (1)


Toby.Howard@manchester.ac.uk



Local and global illumination



Local illumination
model



Global illumination
model

2

Local and global models

- We can make models of light-matter interaction in two ways
 - **Locally**: we treat each object in a scene separately from any other object
 - **Globally**: we treat all objects together, and model the interactions between objects
- we'll develop a simple local illumination model in detail
- we'll look at global models in COMP37111 next year

3

Illumination models

- The interaction of light and matter is an extremely complex process
- In computer graphics we try to **model** this process. In other words, we **approximate** it
- Computer pictures are digital, with finite precision. We can only ever approximate.




4

Local illumination: elements


- We'll develop a model step-by-step, to include the following:
 - Ambient illumination
 - Diffuse reflection
 - Positional light source
 - Specular reflection
 - Colour of lights and surfaces

5

Reflectivity

- There are three kinds of reflection:
 - Perfect diffuse reflection — 
 - Perfect specular reflection — 
 - Imperfect specular reflection — 

6




MANCHESTER
1824

Diffuse and specular reflection

- **Diffuse** reflection is absorption and uniform re-radiation
 - Some wavelengths are absorbed, some are reflected
 - a green object looks green because it only reflects green
- **Specular** reflection is reflection at the air/surface interface
 - To a first approximation, the colour of the specular reflection is that of the light source

8

Perfect diffuse reflection




- Incident rays of light are reflected in all directions from the surface

9

Perfect diffuse reflection

- A perfect diffuse surface reflects an incoming ray across all angles



- The surface looks dull, or matte

10

MANCHESTER
1824

Some diffuse reflectors



- Sponge



- Brick



- Carpet





- Felt

11

MANCHESTER
1824

Perfect specular reflection


- A perfect specular surface reflects an incoming ray like a perfect mirror



12

Imperfect specular reflection

- A imperfect specular surface reflects an incoming ray across a small range of angles



- The surface looks shiny, with highlights

13

Some imperfect specular reflectors



- Lacquer-coated aluminium



- Glazed ceramic




- Stainless steel

14

Diffuse/specular surfaces

- Most surfaces exhibit a combination of diffuse and specular reflection



- We can model these effects with varying degrees of realism

15

Illumination sources

- We begin our development of the local illumination model by considering **diffuse** reflection, and **sources** of illumination:
 - Ambient illumination
 - Point illumination source at infinity (directional illumination)
 - Point illumination source in the scene

16

Ambient illumination

- Consider an environment with a light source
- Multiple reflections cause a general level of illumination in the scene



17

Ambient illumination

- If intensity of ambient light is I_a
- Amount of light diffusely reflected from a surface is

$$I_{\text{ambient}} = k_a I_a$$

- Where k_a is the **ambient reflection coefficient** of the surface

$$0 \leq k_a \leq 1$$

18

Local illumination model v1


- We now have the first term in the model we're developing

- $I = \text{ambient}$

- $I = k_a I_a$


19

Wireframe versus ambient



Wireframe rendering


Rendering with
ambient illumination



20

Effect of ambient illumination

- Each object is uniformly illuminated
- Lose 3D information



21

Varying k_a

I_a is constant

$k_a = 0.1$  $k_a = 1.0$



22

True ambient lighting

Note: the

$$I = k_a I_a$$

term is a gross simplification of true ambient illumination, which is **not** constant in a scene



Advanced Interfaces Group

23

True ambient lighting



Real scene (photograph)




Scene modelled and rendered with accurate global illumination model

Cornell University

24

Directional lighting




Ken Brodie

- We need to model the effects of
 - different angles of incidence
 - different distances from the light source

25


Directional lighting

- Let's consider light from a point source at infinity – so we're only concerned with the direction



26


Describing surface orientation



- \hat{N} is surface normal
- \hat{L} is direction of light source
- θ is angle of incidence
- Note the vectors are **normalised**

27

Diffuse reflection: Lambert's Law



- Light source of intensity I_p
- Effective intensity received is I_e
- Lambert's Law: $I_e = I_p \cos \theta$




Johann Heinrich
Lambert
(1728-1777)

28

Lambert's Law derived

- Consider light of intensity I and cross-sectional width x falling on a surface:




- So width x on surface receives all of intensity I

29

Lambert's Law derived

- Now consider the light inclined at θ
- $x' = x / \cos \theta$, so $x = x' \cos \theta$
- So width x receives intensity $I \cos \theta$



30

Diffuse reflectivity

- We express how good a diffuse reflector a surface is using k_d
- k_d is the **diffuse reflection coefficient** of the surface, $0 \leq k_d \leq 1$
- Amount of diffusely reflected light is
 - $I_{\text{diffuse}} = I_p k_d \cos \theta$
 - $I_{\text{diffuse}} = I_p k_d (\hat{\mathbf{N}} \cdot \hat{\mathbf{L}})$

31


Local illumination model v2

- $I = \text{ambient} + \text{diffuse}$
- $$I = k_a I_a + I_p k_d (\hat{\mathbf{N}} \cdot \hat{\mathbf{L}})$$

32


MANCHESTER
1824

Ambient versus point source



Rendering with ambient illumination

Rendering with a point light source and diffuse reflection




Ken Brodie

33


MANCHESTER
1824

Ambient versus point source



Rendering with ambient illumination

Rendering with a point light source and diffuse reflection



34

Effect of k_d

- Varying k_d from 0 to 1



35

Light source distance

- Physically, light intensity falls off with the square of distance travelled
- After travelling d , original intensity I_p is now I_e

- $$I_e = \frac{I_p}{4\pi d^2}$$

36

Light source distance

- While this is physically correct, it doesn't always work well for computer graphics
- We only have a limited number of pixel intensities, and often the d^2 term changes too rapidly, so instead we use:

$$I_e = \frac{I_p}{k_c + k_l d + k_q d^2}$$

We can choose k_c, k_l, k_q for the best results

37

Local illumination model v3

- I = ambient + distance (diffuse)


$$I = k_a I_a + \frac{I_p}{d'} k_d (\hat{\mathbf{N}} \cdot \hat{\mathbf{L}})$$

- Where $d' = k_c + k_l d + k_q d^2$

38

MANCHESTER
1824

Modelling specular reflection




- $\hat{\mathbf{R}}$ is a vector giving the direction of maximum specular reflection.
- $\hat{\mathbf{R}}$ makes an angle of θ with $\hat{\mathbf{N}}$, as does $\hat{\mathbf{L}}$
 - (for a perfect mirror, angle of incidence == angle of reflection)
- $\hat{\mathbf{V}}$ is a vector pointing to the observer's position

39

MANCHESTER
1824

Modelling specular reflection




- Experiment shows specular reflection varies with:
 - Angle ϕ between $\hat{\mathbf{R}}$ and $\hat{\mathbf{V}}$
 - Incident angle θ and light wavelength λ
 - So we seek a function $I_{\text{specular}} = S(\phi, \theta, \lambda)$

40


Modelling specular reflection

- What is this function $S(\phi, \theta, \lambda)$?
- We'll look at the effects of ϕ, θ, λ in turn
- First, ϕ



41

Modelling specular reflection




Observed specular = 100%

- Specular reflection spread out, around $\hat{\mathbf{R}}$
- Viewer sees maximum specular reflection when $\hat{\mathbf{V}}$ coincides with $\hat{\mathbf{R}}$

42

MANCHESTER
1824

Modelling specular reflection




$\phi = 0^\circ$
Observed specular = 100%

- As $\hat{\mathbf{V}}$ diverges from $\hat{\mathbf{R}}$ by angle ϕ , viewer sees less specular reflection

43

MANCHESTER
1824

Modelling specular reflection




$\phi = 20^\circ$
Observed specular = 60%

- As $\hat{\mathbf{V}}$ diverges from $\hat{\mathbf{R}}$ by angle ϕ , viewer sees less specular reflection

44

MANCHESTER
1824

Modelling specular reflection



Observed specular = 5%

- As $\hat{\mathbf{V}}$ diverges from $\hat{\mathbf{R}}$ by angle ϕ , viewer sees less specular reflection


45

MANCHESTER
1824

Modelling specular reflection

- Variation of observed specular = $F(\phi)$
- But what is the function F ?
- Bui-Tuong Phong (1942-1975) proposed using the function $\cos^n \phi$





46



MANCHESTER
1824

Phong's specular function

- So we now have
 - $I_{\text{specular}} = I_p \cos^n \phi$
- Which we can rewrite using vectors as
 - $I_{\text{specular}} = I_p (\hat{\mathbf{R}} \cdot \hat{\mathbf{V}})^n$
- Normally we use $1 \leq n \leq 200$

$n=10$ $n=20$ $n=80$ $n=200$


Richard Lobb

Incident angle and wavelength

- Recall $I_{\text{specular}} = S(\phi, \theta, \lambda)$
- We've accounted for ϕ
- Now we look at the effects of θ and λ



49

Modelling specular reflection




- Experiment shows specular reflection varies with:
 - Angle ϕ between \hat{R} and \hat{V}
 - Incident angle θ and light wavelength λ
 - So we seek a function $I_{\text{specular}} = S(\phi, \theta, \lambda)$

50



Reflectance of a non-conductor



53

Incident angle and wavelength

- This complex variation is expressed by the Fresnel equation

(Augustin-Jean Fresnel, 1788-1827)

- A founder of the wave theory of light.
- Developed theory of diffraction of light.
- Obtained circularly polarised light
- Developed the use of compound lenses instead of mirrors for lighthouses.




54

Incident angle and wavelength

- This complex variation is expressed by the Fresnel equation (Augustin-Jean Fresnel, 1788-1827)

$$F = \frac{1}{2} \left[\frac{\sin^2(\phi - \theta)}{\sin^2(\phi + \theta)} + \frac{\tan^2(\phi - \theta)}{\tan^2(\phi + \theta)} \right]$$

- F is the fraction of light reflected
- $\sin \theta = \sin \phi / \mu$
- μ is the refractive index of the material (λ dependent)



55

Incident angle and wavelength

- In practice, we usually approximate F with a single constant k_s
- k_s is the **specular reflection coefficient** of the surface, $0 \leq k_s \leq 1$
- $I_{\text{specular}} = I_p k_s (\mathbf{R} \cdot \mathbf{V})^n$
- But, we sacrifice accuracy for efficiency

56

Local illumination model v4


- I = ambient + distance (diffuse + specular)

$$I = k_a I_a + \frac{I_p}{d'} \left[k_d (\hat{\mathbf{N}} \cdot \hat{\mathbf{L}}) + k_s (\hat{\mathbf{R}} \cdot \hat{\mathbf{V}})^n \right]$$

- Where $d' = k_c + k_l d + k_q d^2$


57

Diffuse and specular rendering



Rendering with
a point light source
and diffuse reflection


Rendering with
a point light source
and diffuse and
specular reflection



Ken Brodlie

58

Diffuse and specular rendering




Rendering with
a point light source
and diffuse reflection



Rendering with
a point light source
and diffuse and
specular reflection

59

Varying k_d and k_s



Alan Watt

60

MANCHESTER
1824

Varying k_s and exponent n

k_s increasing

Exponent n increasing

Alan Watt

61

MANCHESTER
1824

Phong applet

Richard Lobb

- <http://www.cs.auckland.ac.nz/~richard/research-topics/PhongApplet/PhongDemoApplet.html>

62

Phong demo




<http://learningwebgl.com/lessons/lesson14/>


63

Inaccuracy of Phong model

- For a large angle of incidence, the Phong \cos^n term is incorrect



Phong \cos^n approximation



Accurate Fresnel-based reflection

64

Incorporating colour

- So far, we've only considered light **intensity**, not **colour**
- It's easy to incorporate – we express light colour as a triple of RGB intensities:
 - I_{pR}, I_{pG}, I_{pB}
- And correspondingly we express surface colour using
 - k_{aR}, k_{aG}, k_{aB}
 - k_{dR}, k_{dG}, k_{dB}

65

Local illumination model v5

- We now have a separate expression for each colour component. For example, for red:
- $I_R = \text{ambient}_{\text{Red}} +$
distance (diffuse_{Red} + specular)

$$I_R = k_{aR} I_{aR} + \frac{I_{pR}}{d'} \left[k_{dR} (\hat{\mathbf{N}} \cdot \hat{\mathbf{L}}) + k_s (\hat{\mathbf{R}} \cdot \hat{\mathbf{V}})^n \right]$$

66

Multiple lights


- Finally, what if we have more than one light?
- Easy, compute illumination separately for each and sum. For M lights:

- $$I = ambient + \sum_{i=1}^M (diffuse_i + specular_i)$$


67

Multiple lights

One light



Two lights




Ken Brodlie


68

MANCHESTER
1824


Examples of the model in use




Lex Lemnings




AIG



James Sinnott



Alan Watt



nVIDIA

69

MANCHESTER
1824

The “standard” model

- The local illumination model we’ve developed is the “standard” model in use today
- Implemented in OpenGL
- Implemented in hardware on consumer 3D graphics cards

70